## 1 A meta-analysis of genome-wide association studies of childhood wheezing

# 2 phenotypes identifies ANXA1 as a susceptibility locus for persistent wheezing

- 3 Granell R<sup>1\*</sup>, Curtin JA<sup>2</sup>, Haider S<sup>3</sup>, Kitaba N<sup>4</sup>, Mathie S<sup>3</sup>, Gregory L<sup>3</sup>, Yates LL<sup>3</sup>, Tutino M<sup>2</sup>, Hankinson
- 4 J<sup>2</sup>, Perretti M<sup>5</sup>, Vonk JM<sup>6,7</sup>, Arshad SH<sup>8,9,10</sup>, Cullinan P<sup>3</sup>, Fontanella S<sup>3</sup>, Roberts G<sup>4,8,9</sup>, Koppelman
- 5 GH<sup>7,11</sup>, Simpson A<sup>2</sup>, Turner S<sup>12</sup>, Murray CS<sup>2</sup>, Lloyd CM<sup>3§</sup>, Holloway JW<sup>4,8,§</sup>, Custovic A<sup>3,§</sup>

# 6 on behalf of UNICORN<sup>13</sup> and Breathing Together<sup>14</sup> Investigators

- 7 §Equal contribution, joint senior authors
- 8 1 MRC Integrative Epidemiology Unit, Department of Population Health Sciences, Bristol Medical School,
- 9 University of Bristol, Bristol, United Kingdom;
- 10 2 Division of Infection, Immunity and Respiratory Medicine, School of Biological Sciences, The University of
- 11 Manchester, Manchester Academic Health Science Centre, and Manchester University NHS Foundation
- 12 Trust;
- 13 3 National Heart and Lung Institute, Imperial College London, London, United Kingdom;
- 4 Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, United
- 15 Kingdom;
- 16 5 William Harvey Research Institute, Barts and The London School of Medicine Queen Mary University of
- 17 London, London, United Kingdom;
- 18 6 University of Groningen, University Medical Center Groningen, Department of Epidemiology, Groningen,
- 19 the Netherlands.
- 20 7 University of Groningen, University Medical Center Groningen, Groningen Research Institute for Asthma
- 21 and COPD (GRIAC), Groningen, the Netherlands.
- 22 8 NIHR Southampton Biomedical Research Centre, University Hospitals Southampton NHS Foundation
- 23 Trust, Southampton, United Kingdom;
- 24 9 David Hide Asthma and Allergy Research Centre, Isle of Wight, United Kingdom;
- 25 10 Clinical and Experimental Sciences, Faculty of Medicine, University of Southampton, Southampton,
- 26 United Kingdom;
- 27 11 University of Groningen, University Medical Center Groningen, Beatrix Children's Hospital, Department
- of Pediatric Pulmonology and Pediatric Allergology, Groningen, the Netherlands.
- 29 12 Child Health, University of Aberdeen, Aberdeen, United Kingdom

- \*Correspondence and requests for reprints should be addressed to Dr Raquel Granell, Population
   Health Sciences, Bristol Medical School, University of Bristol, BRISTOL BS8 2BN, UK
   E-mail Raquel.Granell@bristol.ac.uk
- 33 This article is dedicated to the memory of our wonderful colleague and friend Professor John
- 34 Henderson (1958-2019), whose contribution to the understanding of the heterogeneity of
- 35 childhood asthma cannot be overstated. Rainbow-chasers and UNICORN riders forever.
- 36 <sup>13</sup>STELAR/UNICORN investigators
- 37 Professor Graham Devereux
- 38 Dr Dimitrios Charalampopoulos
- 39 <sup>41</sup>Breathing Together investigators
- 40 Prof Andrew Bush
- 41 Prof Sejal Saglani
- 42 Prof Benjamin Marsland
- 43 Prof Jonathan Grigg
- 44 Prof Jurgen Schwarze
- 45 Prof Mike Shields
- 46 Prof Peter Ghazal
- 47 Prof Ultan Power
- 48 Dr Ceyda Oksel
- 49 **Keywords:** GWAS, asthma, wheeze phenotypes, Annexin 1
- Data availability: The informed consent obtained from all included participants does not allow the
- data to be made freely available through any third party maintained public repository. However,
- 52 data used for this submission can be made available on request to the corresponding cohort
- 53 Executive. The ALSPAC website provides information on how to request and access its data (

- 54 <a href="http://www.bristol.ac.uk/alspac/researchers/access/">http://www.bristol.ac.uk/alspac/researchers/access/</a>). For queries regarding access of data from
- 55 MAAS, IoW, SEATON or Ashford please contact Philip Couch <a href="mailto:philip.couch@manchester.ac.uk">philip.couch@manchester.ac.uk</a>).
- 56 All summary data used to plot the figures in our manuscript has been deposited in Dryad.
- 57 **Code availability:** Code used for this submission has been deposited in Dryad.
- 58 Competing Interests statement: Graham Roberts: MRC grant to my institution President of the
- 59 British Society of Allergy and Clinical Immunology. Gerard Koppelman: Dutch Lung Foundation,
- 60 Ubbo Emmius Foundation (Money to insititution) Dutch Lung Foundation, Vertex, TEVA the
- 61 Netherlands, GSK, ZON-MW (VICI grant), European Union (Money to institution) Astra Zeneca,
- 62 Pure IMS, GSK (Money to institution) Sanofi, Boehringer Ingelheim (Money to institution). Angela
- 63 Simpson: Medical research council Research grant JP Moulton Charitable Foundation Research
- 64 grant Asthma UK Research grant. Clare Murray: Asthma Uk National Institute for Health Research
- 65 Moulton Charitable Foundation North West Lung Centre Charity GSK (Lecture fees) Novartis
- 66 (Lecture fees). Clare Lloyd: Wellcome Trust 107059/Z/15/Z. John Holloway: Medial Research
- 67 Council grant MR/S025340/1 (to institution) American Academy of Allergy Asthma and
- 68 Immunology (AAAI) (Support for speaker travel to AAAAI annual congress). Adnan Custovic: MRC
- 69 (research grants) EPSRC (research grant) Wellcome Trust (research grant) Worg Pharmaceoticals
- 70 (Personal payment <US\$5000) GSK Honorarium for lecture, personal, <US\$ 5000 AstraZeneca
- 71 Honorarium for lecture, personal, <US\$ 5000 Sanofi Honorarium for lecture, personal, <US\$ 5000
- 72 Stallergens-Greer Honorarium for lecture, personal, <US\$ 5000 WAO (Board of officers, unpaid).
- 73 Funding: Supported by the UK Medical Research Council (MRC) Programme Grant MR/S025340/1,
- 74 the Wellcome Trust (WT) Strategic Award (108,818/15/Z) and a WT Senior Fellowship to CML
- 75 (107059/Z/15/Z). The MRC and Wellcome (Grant ref: 217065/Z/19/Z) and the University of Bristol
- 76 provide core support for ALSPAC. ALSPAC GWAS data was generated by Sample Logistics and

- 77 Genotyping Facilities at Wellcome Sanger Institute and LabCorp (Laboratory Corporation of
- America) using support from 23andMe. A comprehensive list of grants funding is available on the
- 79 ALSPAC website (http://www.bristol.ac.uk/alspac/external/documents/grant-
- 80 <u>acknowledgements.pdf</u>).
- PIAMA was funded by the Netherlands Lung Foundation (grant 3.4.01.26, 3.2.06.022, 3.4.09.081
- 82 and 3.2.10.085CO), the ZON-MW Netherlands Organization for Health Research and Development
- 83 (grant 912-03-031), the Stichting Astmabestrijding and the Ministry of the Environment. Genome-
- 84 wide genotyping was funded by the European Commission as part of GABRIEL (grant number
- 85 018996) and a grant from BBMRI-NL (CP 29). GHK is supported by a ZON-MW VICI grant.
- 86 For the purpose of Open Access, the author has applied a CC BY public copyright license to any
- 87 Author Accepted Manuscript version arising from this submission.

## **ABSTRACT**

**Background:** Many genes associated with asthma explain only a fraction of its heritability. Most genome-wide association studies (GWASs) used a broad definition of "doctor-diagnosed asthma", thereby diluting genetic signals by not considering asthma heterogeneity. The objective of our study was to identify genetic associates of childhood wheezing phenotypes.

**Methods:** We conducted a novel multivariate GWAS meta-analysis of wheezing phenotypes jointly derived using unbiased analysis of data collected from birth to 18 years in 9,568 individuals from five UK birth-cohorts.

Results: 44 independent SNPs were associated with early-onset persistent, 25 with preschool remitting, 33 with mid-childhood remitting and 32 with late-onset wheeze. We identified a novel locus on chr9q21.13 (close to annexin 1 (*ANXA1*), p<6.7×10<sup>-9</sup>), associated exclusively with early-onset persistent wheeze. We identified rs75260654 as the most likely causative single nucleotide polymorphism (SNP) using Promoter Capture Hi-C loops, and then showed that the risk allele (T) confers a reduction in *ANXA1* expression. Finally, in a murine model of house dust mite (HDM)-induced allergic airway disease, we demonstrated that anxa1 protein expression increased and anxa1 mRNA was significantly induced in lung tissue following HDM exposure. Using anxa1<sup>-/-</sup> deficient mice, we showed that loss of anxa1 results in heightened airway hyperreactivity and Th2 inflammation upon allergen challenge.

**Conclusions:** Targeting this pathway in persistent disease may represent an exciting therapeutic prospect.

**Funding**: UK Medical Research Council Programme Grant MR/S025340/1 and the Wellcome Trust Strategic Award (108,818/15/Z) provided most of the funding for this study.

## **Abstract word count: 240**

#### INTRODUCTION

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

Asthma is a complex disorder caused by a variety of mechanisms which result in multiple clinical phenotypes(1). It has a strong genetic component, and twin studies estimate its heritability to be ~ 60-70%(2). "Asthma genes" have been identified through a range of approaches, from candidate gene association studies(3) and family-based genome-wide linkage analyses(4) to genome-wide association studies (GWASs)(5-7). The first asthma GWAS (2007) identified multiple markers on chromosome 17q21 associated with childhood-onset asthma(5). A comprehensive review summarising the results of 42 GWASs of asthma and asthma-related traits has been published recently(8). The most widely replicated locus is 17q12-21, followed by 6p21 (HLA region), 2q12 (IL1RL1/IL18R1), 5q22 (TSLP) and 9p24 (IL33)(9). Overall, the evidence suggests that multiple genes are underlying the association peaks(9). However, despite undeniable successes, genetic studies of asthma have produced relatively heterogeneous results, and only a small proportion of the heritability is accounted for (10). One part of the explanation for the paucity of precise replication are numerous gene-environment interactions(11). Another important consideration is asthma heterogeneity, in that asthma diagnosis comprises several conditions with distinct pathophysiology(12, 13), each potentially underpinned by different genetic associations(14). However, in order to maximise sample size, most GWASs used a definition of "doctor-diagnosed asthma" (15). Such aggregated outcome definitions are imprecise (16) and phenotypically and mechanistically heterogeneous (17), and this heterogeneity may dilute important genetic signals(14). One way of disaggregating asthma diagnosis is to use data-driven methods to derive subtypes in a hypothesis-neutral way(18). For example, we jointly modelled data on wheezing from birth to adolescence in five UK population-based birth cohorts and identified five distinct phenotypes (19).

However, although latent modelling approaches have been instrumental in elucidating the heterogenous nature of childhood asthma diagnosis(13), there has been little research into the genetic associations of phenotypes derived using data-driven methods. This is the first study to investigate the genetic architecture of wheezing phenotypes from infancy to adolescence, to identify genes specific to each phenotype and better understand the genetic heterogeneity between the disease class profiles.

#### **MATERIALS & METHODS**

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

Study design, setting, participants and data sources/measurement The Study Team for Early Life Asthma Research (STELAR) consortium(20) brings together five UK population-based birth cohorts: Avon Longitudinal Study of Parents and Children (ALSPAC)(21), Ashford(22) and Isle of Wight (IOW)(23) cohorts, Manchester Asthma and Allergy Study (MAAS)(24) and the Aberdeen Study of Eczema and Asthma to Observe the Effects of Nutrition (SEATON)(25). All studies were approved by research ethics committees. See Appendix 1: Description of cohorts for more details. Informed consent was obtained from parents, and study subjects gave their assent/consent when applicable. Validated questionnaires were completed on multiple occasions from infancy to adolescence (19). A list of variables, per cohort, is shown in **Appendix 1-Table 1**, and the cohort–specific time points and sample sizes in Appendix 1-Table 2. Data were harmonised and imported into Asthma eLab web-based knowledge management platform to facilitate joint analyses (20). Definition of primary outcome (wheeze phenotypes from infancy to adolescence) In the pooled analysis among 15,941 subjects with at least two observations on current wheeze, we used latent class analysis (LCA) to derive wheeze phenotypes from birth to age 18 years (19). A detailed description of the analysis is presented in(19) and in Appendix 1: Definition of variables. A five-class solution was selected as the optimal model(19), and the classes (wheeze phenotypes) were labeled as: (1) never/infrequent wheeze (52.4%); (2) early-onset pre-school remitting wheeze (18.6%); (3) early-onset middle-childhood remitting wheeze (9.8%); (4) early-onset persistent wheeze (10.4%); and (5) late-onset wheeze (8.8%). These latent classes were used in the subsequent GWAS.

## **Genotyping, imputation and GWAS Meta-Analysis**

Genotyping, quality control, imputation and exclusions are described in **Appendix 2: Genotyping** and imputation. Analyses were performed independently in ALSPAC, MAAS and the combined IOW-SEATON-ASHFORD (genotyped on the same platform, at the same time, and imputed together). We used SNPTEST v2.5.2(26) with a frequentist additive multinomial logistic regression model (-method newml), using the never/infrequent wheeze as the reference and without including any covariates. A meta-analysis of the three GWASs was performed using METAL(27) with a total of 8,057,852 single nucleotide polymorphism (SNPs). See **Appendix 2: LD clumping**, **pre-Selection and Gene Annotation** for more details.

#### **Post-GWAS studies**

Our GWAS identified a novel locus in chr9q21 nearby *Annexin A1* (*ANXA1*), exclusively associated with early-onset persistent wheeze (see results section). We therefore proceeded with studies to identify causal variants and explore the biological mechanisms underlying this locus (see **Appendix 3: Post-GWAS: rs75260654** (*ANXA1*) for more details). To this end, we firstly identified the most likely causative SNP using Promoter Capture Hi-C loops. We then ascertained genotype effect on gene expression and assessed the potential biological function of *ANXA1* in asthma. Finally, we used a murine model of house dust mite (HDM)-induced allergic airway disease to investigate whether *ANXA1* was important in regulating immune responses to a clinically relevant aeroallergen and used knock-out mice to derive further *in vivo* functional data to support our GWAS-finding.

## **RESULTS**

## Participants and descriptive data

We included a total of 9,568 subjects with European ancestry: ALSPAC, n=6833; MAAS, n=887; SEATON, n=548; ASHFORD, n=348; and IOW, n=952. Demographic characteristics of the participants in STELAR cohorts included in this analysis and a flow chart are shown in **Appendix 1-Table 3** and **Appendix 1-Figure 1**. Cohorts contain similar proportions of males (range 48%-54%), maternal history of asthma (11%-14%), maternal smoking (14%-23%), (doctor-diagnosed) asthma ever during mid-childhood (16%-24%) and adolescence (20%-30%), current wheeze (12%-20% mid-childhood, 9%-25% adolescence) and current use of asthma medication (12%-17% mid-childhood, 11%-17% adolescence). Individuals with missing genetic data, as well as related and non-European individuals were excluded. Comparison of included vs. excluded individuals across cohorts (per cohort and time point) is in **Appendix 1** and **Appendix 1-Table 4**.

#### **GWAS Meta-Analysis**

We conducted three GWASs (ALSPAC, MAAS, IOW-SEATON-Ashford) in parallel and results were meta-analyzed. The distribution of the minor allele frequencies was consistent across genotyped datasets (mean SD 0.01). A circular Manhattan plot and a QQ-plot are shown in **Figure 1-figure supplement 1**. Some observed p-values were clearly more significant than expected under the null hypothesis, particularly for Eearly-onset persistent wheeze, without an early separation of the expected from the observed which indicates low evidence of population stratification. We observed slight deflation of the meta-analysis pvalues in our summary statistics. Genomic Inflation Factor ( $\lambda$ ) for Early-onset Pre-school Remitting= 0.96, Early-onset Mid-childhood Remitting = 0.94, Late-onset= 0.96, and Early-onset Persistent wheezing= 0.97. . A total of 589 SNPs were associated with at least one phenotype with p<10<sup>-5</sup>. After clumping, we identified 134

independent SNPs uniquely associated with different phenotypes (p<10<sup>-5</sup>): of these, 44 were exclusively associated with early-onset persistent, 25 with early-onset preschool remitting, 33 with early-onset mid-childhood remitting and 32 with late-onset wheeze (Appendix 2-Table 1). Scatter plots in Figure 2- figure supplement 1 show the heterogeneity in the genetic profile of the wheeze phenotypes. The plots show that all signals were phenotype-specific at p<10<sup>-5</sup> and only nominal associations were shared across wheezing phenotypes. More details on how these plots were derived can be found in Appendix 2: Heterogeneity scatter plots. For example, chr17q21 was identified as a top locus for early-onset persistent wheeze (p=5.42×10<sup>-9</sup>), but some of the SNPs in this region were also associated with the early-onset mid-childhood remitting phenotype (p<10<sup>-4</sup>). To help identify functional elements located near the GWAS-associated variants (potential causal variants), we used locus zoom plots (LZP) for the 134 independent SNPs (p<10<sup>-5</sup>). Following close inspection of all plots, we short-listed 85 independent SNPs (Appendix 2-Table 1) for which the LZPs potentially indicated more than one causal variant (Appendix 2-Figures 1-4) and followed them up for further annotation. The results of GWAS meta-analysis for these 85 SNPs with main associations across the four wheeze phenotypes are presented in Table 1. Previously associated traits for each region/gene associated with the different wheeze phenotypes are shown in Appendix 4-Tables 1-4 and results are summarized in Appendix 4: Results in context of literature. Briefly, one region (6q27) among the top hits for early-onset preschool remitting wheeze was previously associated with asthma, but in the context of obesity with a nominal association with asthma and BMI(28). Another region/gene (3q26.31/NAALADL2) identified as top hit for earlyonset preschool remitting wheeze, was reported as an associate of severe asthma exacerbations, but only at nominal level (29). No regions/genes identified as top hits for early-onset midchildhood remitting wheeze were found to have previous associations with asthma. Several genes/loci identified as top hits for late-onset wheeze were previously associated with asthma:

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

asthma susceptibility(31)), CD200 3q13.2 (adult onset non-allergic asthma(32)), GIMAP family 7q36.1 (autoimmune diabetes, asthma, allergy(33)), 9p22.3 (asthma in <16 year-old (34)) and 16p12.1 (asthma and rhino-conjunctivitis at 10-15 years(35)). We identified two GWAS-significant loci for early-onset persistent wheeze: 17q21, p<5.5×10<sup>-9</sup>, and a novel locus on 9q21.13 (ANXA1), p<6.7×10<sup>-9</sup>. The ANXA1 locus was the only GWAS-significant locus that had not previously been associated with asthma or atopic traits, with one previous study showing an association with FEV<sub>1</sub>/FVC and bronchodilator response in smokers(36). ANXA1 is strongly expressed in bronchial mast cells and has anti-inflammatory properties (37), and may be involved in epithelial airway repair (38) (Appendix 4-Table 1). We therefore followed up top SNPs from this locus. ANXA1 locus and persistent wheeze Two SNPs (rs75260654, the lead SNP, and rs116849664 located downstream of ANXA1) were associated with early-onset persistent wheeze at genome-wide significance (GWS), with an additional SNP rs78320984 almost reaching GWS (Appendix 5-Table 1). These SNPs are in LD with each other (**Appendix 5-figure 1**), but not with any other SNPs. Promoter Capture identifies rs75260654 as the most likely causative variant To identify the most likely causative variant, we investigated the overlap of the SNPs with Promoter Capture Hi-C interactions involving the ANXA1 promoter in CD4+ cells in MAAS cohort subjects. Of the three SNPs, only rs75260654 overlapped a region interacting with the ANXA1

ACOXL chr2q13 (later onset asthma and obesity(30)), PRKAA2 chr1p32.2 (lymphocyte count and

promoter (Figure 3). Moreover, rs75260654 overlapped a POLR2A ChIP-seq peaks and an ATAC-

seq peak and active enhancer in the type II pneumocyte derived A549 cell line. This shows that

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

252 transcriptionally active in relevant cell types. 253 Allele Frequencies of rs75260654 (MAF=0.02) across wheeze phenotypes are shown in **Appendix** 254 5-Table 2. Two individuals (one in MAAS and one in ALSPAC) were homozygote for the minor allele 255 (T), and both were in the early-onset persistent wheeze class. One subject reported current 256 wheeze and asthma through childhood, with hospitalizations for lower respiratory tract infection in the 1<sup>st</sup> year of life confirmed in healthcare records. The second individual reported current 257 258 wheezing at 1.5, 2.5 and 8-9 years and doctor-diagnosed asthma and the use of asthma 259 medication at 8-9 years. 260 Rs75260654: Effect on Genomic Features 261 VEP prediction shows the SNP rs75260654 (C changed to T) to be located downstream of three 262 protein-coding transcripts of AXNA1 and overlapping the known regulatory region ID 263 ENSR00000882742 on Chromosome 9: 73,173,001-73,173,200. This region is active in the GI tract, 264 M2 macrophages, neural progenitor cells, and trophoblasts, but is repressed in Tlymphocytes 265 including CD4+ CD25+, Treg, and CD8+ cells. 266 Rs75260654: Effect on gene expression 267 The effect of rs75260654 on the expression of nearby genes was investigated by browsing the 268 eQTL GTEX data available in Ensembl. Compared to C, the T allele was found to reduce the 269 expression of ANXA1 in naïve B-cells (effect size=-2.36795, p=0.01) and to increase expression in 270 Lymphoblasoid Cell Lines (LCL) (effect size=0.848856, pe=0.001) (Figure 4). This SNP affects 271 expression of the neighboring gene ALDH1A1 (aldehyde dehydrogenase-1 family member A1)

rs75260654 is located in a region directly interacting with the ANXA1 promoter and is

(effect size=-2.40446, p=0.0039 in macrophages infected with Salmonella). In the eQTL catalogue,

rs75260654 is identified as an eQTL of ANXA1 in various immune cells (at nominal significance)

272

273

including T cells, monocytes, fibroblasts, whole blood, Th2 memory cells, naïve B cells. rs75260654 is also an eQTL of ANXA1 in monocytes that were stimulated with R848 (agonist of TLRs 7 and 8) and a human seasonal influenza A virus (39) (at nominal significance) (Appendix 5-Table 3). In the lung rs116849664 and rs78320984 (both in LD with rs75260654) were eQTLs of ANXA1 (Appendix **5-Table 4**) as well as LINC01474 at nominal significance levels. Additional supporting evidence regarding the significance of the T-allele on the expression of these genes was provided using eQTLGene Consortium meta-analysis of 24 Cohorts and 24331 samples(40). This method reproduced the previous modest results showing a cis-eQTL effect of rs75260654 on both the ANXA1 ( $p=6.02\times10^{-23}$ ) and ALDH1A1 ( $p=1.11\times10^{-19}$ ) at FDR=0. No significant trans-eQTLs were observed. Potential biological function of ANXA1 in asthma Protein-protein network analysis demonstrated that ANXA1 interacts directly with genes enriched for asthma (including IL4 and IL13) and inflammatory regulation (NR3C1, Glucocorticoid receptor) showing its significance in dysregulation of the immune response (see Appendix 5-figure 2 and Appendix 5-Table 5). Functional studies of anxa1 in a murine model Pulmonary expression of anxa1 is modulated by aeroallergen exposure We first analysed expression of anxa1 using a model of HDM-induced allergic airway disease (Figure 5A)(41). Consistently, immunohistochemistry analysis revealed anxa1 protein expression increased following HDM challenge (Figure 5B,C). Anxa1 mRNA was significantly induced in lung tissue following HDM exposure (Figure 5D). This increase suggests that the pro-resolving anxa1

may play a role in regulating the pulmonary immune response to allergen.

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

Anxa1 suppresses allergen-induced airway hyperresponsiveness (AHR) and type-2 inflammation

To confirm a functional role for anxa1 in allergic airway disease, we exposed anxa1<sup>-/-</sup> mice to
intranasal HDM. Wildtype mice given HDM over 3 weeks developed significant AHR compared to
PBS control mice. Mice deficient in anxa1 had significantly worse lung function (greater airway
resistance) compared to WT treated mice (Figure 5E). Anxa1<sup>-/-</sup> mice exhibited significantly
increased airway eosinophilia and elevated numbers of Th2 lymphocytes (Figure 5F,G). Lung tissue
cytokine levels reflected the exacerbated airway Th2 inflammation, with elevation in IL-4, and
significant induction of IL-5 and IL-13 (Figure 5H,J). Thus, anxa1 deficiency results in an alteration
of the pulmonary immune response, with uncontrolled eosinophilia and an exacerbation of type-2
inflammation and AHR in response to allergen. More details in Appendix 6: Functional mouse
experiments.

## **DISCUSSION**

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

Herein, we present a comprehensive description of the genetic architecture of childhood wheezing disorders. Using a novel approach applied to a unique dataset from five UK birth cohorts, we identified subsets of SNPs differentially associated across four wheezing phenotypes: early-onset persistent (44 SNPs, 19 loci), early-onset preschool remitting (25 SNPs, 10 loci), early-onset midchildhood remitting (33 SNPs, 9 loci) and late-onset (32 SNPs, 20 loci). We found little evidence of genetic associations spanning across different phenotypes. This suggests that genetic architecture of different wheeze phenotypes comprises a limited number of variants likely underpinning mechanisms which are shared across phenotypes, but that each phenotype is also characterized by unique phenotype-specific genetic associations. Importantly, we identified a novel locus in chr9q21 nearby ANXA1 exclusively associated with early-onset persistent wheeze (p<6.7×10<sup>-9</sup>). To identify the most likely causative variant, we investigated the overlap of the associated SNPs with Promoter Capture Hi-C interactions to demonstrate that SNP rs75260654 overlapped a region interacting with the ANXA1 promoter. Using eQTL data, we identified that the risk allele (T) of rs75260654 associated with early-onset persistent wheeze is also associated with ANXA1 expression. Further investigation of the biological function of ANXA1 revealed that it interacts with genes enriched for asthma (including IL4 and IL13) and inflammatory regulation (NR3C1, Glucocorticoid receptor). In functional mouse experiments, anxa1 protein expression increased and anxa1 mRNA was significantly induced in lung tissue following HDM exposure, suggesting that the pro-resolving anxa1 may play a role in regulating the pulmonary immune response to allergen. Concurrently, by utilizing anxa1<sup>-/-</sup> deficient mice we demonstrated that loss of anxa1 results in heightened AHR and Th2 inflammation upon allergen challenge, providing important in vivo functional data to support our GWAS-finding.

ANXA1 is a 37-kDa glycoprotein with potent anti-inflammatory and pro-resolving properties that are mediated by interaction with a specific G-protein coupled receptor FPR2(42). This axis represents an important resolution pathway in chronic inflammatory settings such as those of rheumatoid arthritis(43) and ulcerative colitis(44). ANXA1 belongs to the annexin family of Ca<sup>2+</sup>dependent phospholipid-binding proteins, and through inhibition of phospholipase A2, it reduces eicosanoid production, which also contributes to its anti-inflammatory activities. Modulation of M2 macrophage phenotype is also promoted by ANXA1 to attenuate tissue inflammation(45). Corticosteroids (a mainstay of asthma treatment) increase the synthesis of ANXA1(46). Plasma ANXA1 levels are significantly lower in asthmatic patients with frequent exacerbations compared to those with stable disease, suggesting a link between this mediator and disease state(47). Furthermore, children with wheeze have reduced airway levels of ANXA1(48). Previous functional studies using anxa1<sup>-/-</sup> deficient mice challenged with ovalbumin showed anxa1-deficient mice to have elevated AHR compared to WT mice(49). Ng et al. demonstrated that untreated anxa1-deficient mice have spontaneous AHR that predisposes them to exacerbated response to allergen(49). In the current study, we demonstrated in the murine lung the induction of Anxa1 in response to HDM exposure. In addition, genetic deletion of anxa1 potentiated the development of AHR and enhanced eosinophilia and markers of Th2 inflammation in mice treated with HDM, which is consistent with and extends previous reports. Of interest, in mice, anxa1 expression was recently found to be characteristic of a novel cell-type called the Hillock cell, which may be involved in squamous barrier function and immunomodulation (50). These data identify the ANXA1/FPR2 signaling axis as an important regulator of allergic disease, that could be manipulated for therapeutic benefit.

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

Our study has several limitations. By GWAS standards, our study is comparatively small and may be considered to be underpowered. The sample size may be an issue when using an aggregated definition (such as "doctor-diagnosed asthma") but is less likely to be an issue when primary outcome is determined by deep phenotyping. This is indirectly confirmed in our analyses. Our primary outcome was derived through careful phenotyping over a period of more than two decades in five independent birth cohorts, and although comparatively smaller than some asthma GWASs, our study proved to be powered enough to detect previously identified key associations (e.g. chr17q21 locus). Precise phenotyping has the potential to identify new risk loci. For example, a comparatively small GWAS (1,173 cases and 2,522 controls) which used a specific subtype of early-onset childhood asthma with recurrent severe exacerbations as an outcome, identified a functional variant in a novel susceptibility gene CDHR3 (SNP rs6967330) as an associate of this disease subtype, but not of doctor-diagnosed asthma(51). This important discovery was made with a considerably smaller sample size but using a more precise asthma subtype. In contrast, the largest asthma GWAS to date had a ~40-fold higher sample size(7), but reported no significant association between CDHR3 and aggregated asthma diagnosis. Therefore, with careful phenotyping, smaller sample sizes may be adequately powered to identify larger effect sizes than those in large GWASs with broader outcome definitions(52). The importance of the precise outcome definition was highlighted in our previous studies in ALSPAC which explored genetic associates of wheeze phenotypes derived by LCA(53, 54). Our current findings are consistent with our earlier report suggesting that 17q21 SNPs are associated with early-onset persistent, but not with early transient or late-onset wheeze (53). Further analysis using genetic prediction scores based on 10-200,000 SNPs ranked according to their associations with physician-diagnosed asthma found that the 46 highest ranked SNPs predicted persistent wheeze more strongly than doctor-diagnosed asthma(54). Finally, a candidate gene study

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

combining data from ALSPAC and PIAMA found different associations of IL33-IL1RL1 pathway polymorphisms with different phenotypes(<u>55</u>).

We are cognisant that there may be a perception of the lack of replication of our GWAS findings. We would argue that direct replication is almost certainly not possible in other cohorts, as phenotypes for replication studies should be homogenous (56). However, there is a considerable heterogeneity in LCA-derived wheeze phenotypes between studies, and although phenotypes in different studies are usually designated with the same names, they differ between studies in temporal trajectories, distributions within a population, and associated risk factors(57). This heterogeneity is in part consequent on the number and the non-uniformity of the timepoints used, and is likely one of the factors responsible for the lack of consistent associations of discovered phenotypes with risk factors reported in previous studies (58). This will also adversely impact the ability to identify phenotype-specific genetic associates. For example, we have previously shown that less distinct wheeze phenotypes in PIAMA were identified compared to those derived in ALSPAC(59). Thus, phenotypes that are homogeneous to those in our study almost certainly cannot readily be derived in available populations. This is exemplified in our attempted replication of ANXA1 findings in PIAMA cohort (see Appendix 5: Replication of ANXA1 top hits in PIAMA cohort and Appendix 5-Table 6). In this analysis, the number of individuals assigned to persistent wheezing in PIAMA was small (40), associates of this phenotype differed to those in STELAR cohorts, and the SNPs' imputation scores were low (<0.60), which meant the conditions for replication were not met.

Our study population is of European descent, and we cannot generalize the results to different ethnicities or environments. It is important to highlight the under-representation of ethnically diverse populations in most GWASs(9). To mitigate against this, large consortia have been formed,

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

which combine the results of multiple ethnically diverse GWASs to increase the overall power to identify asthma-susceptibility loci. Examples include the GABRIEL(6), EVE(60) and TAGC(7) consortia, and the value of diverse, multi-ethnic participants in large-scale genomic studies has recently been shown(61). However, such consortia do not have the depth of longitudinal data to allow the type of analyses which we carried out to derive a multivariable primary outcome. Finally, the manual and visual inspection of Locus Zoom plots for the refinement of association signals and identification of functional elements was an objective approach which might have undermined the findings. One strength of our approach is that we used data from five birth cohorts with detailed and lifelong phenotyping, which were harmonised in a common knowledge management platform(20), allowing joint analyses. We performed three parallel GWASs that produced estimates with remarkably consistent directions of effects. In conclusion, using unique data from five UK birth cohorts, we identified subsets of SNPs differentially associated across four wheezing phenotypes from infancy to adolescence. We found little evidence of genetic associations spanning across different phenotypes. We discovered a novel locus in chr9q21 uniquely associated with early-onset persistent wheeze (p<6.7×10<sup>-9</sup>), identified SNP rs75260654 as the most likely causative variant, and demonstrated that the risk allele (T) confers a reduction in ANXA1 expression. In mouse experiments, ANXA1 expression increased in lung tissue following allergen exposure, suggesting that the pro-resolving ANXA1 may play a role in regulating the pulmonary immune response to allergen. Using ANXA1-deficient mice, we demonstrated that loss of ANXA1 results in heightened AHR and Th2 inflammation upon allergen challenge, providing important in vivo functional data to support our GWAS finding. Targeting these pathways to promote the clearance of chronic inflammation in persistent disease

may represent an exciting therapeutic prospect

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

# 422 TABLES AND FIGURES

Table 1. GWAS Meta-analysis: Short-listed 85 top independent SNPs across the four wheezing phenotypes

Early-ons	Early-onset Persistent Wheezing								
Locus	Independent SNPs	Nearby Genes (SNPnexus)	Effect allele (freq)/ other allele	Beta	SE	P-value	Effect Direction (3 GWAS)	min_pval _other*	Previous relevant associations†
1q43	rs4620530	CHRM3	g(0.56)/t	0.25	0.05	2.45E-06	+++	0.79	FEV <sub>1</sub> , FEV <sub>1</sub> /FVC, asthma- high priority drug target
2p25.1	rs13398488	RNF144A	g(0.29)/a	0.25	0.05	2.18E-06	+	0.13	asthma, allergy, childhood onset asthma, allergic rhinitis
2q12.2	rs6543291	FHL2	c(0.4)/t	0.23	0.05	6.97E-06	+++	0.10	bronchial hyper-responsiveness, airway inflammation; novel gene associated with asthma severity in human
3q21.3	rs77655717	EFCC1, RAB43, RAB7A	c(0.05)/t	0.47	0.10	6.40E-06	+++	0.39	<i>RAB43</i> : response to bronchodilator, FEV/FEC ratio; <i>RAB7A</i> : eosinophil count
4m1C 2	rs7680608 <sup>eQTL</sup>	RNF212, IDUA,	g(0.93)/c	-0.42	0.09	1.31E-06		0.15	4p16: asthma
4p16.3	rs77822621 <sup>eQTL</sup>	DGKQ, SLC26A1	c(0.96)/t	-0.50	0.11	7.16E-06		0.01	
4q31.21	rs115228498	INPP4B	c(0.02)/t	0.79	0.17	2.70E-06	+++	0.02	atopic asthma
5p15.31	rs116494115	ADCY2	g(0.01)/a	0.75	0.17	6.49E-06	+++	0.09	asthma $\times$ air pollution, childhood asthma
7q22.3	rs76871421	CDHR3	c(0.12)/t	0.37	0.07	5.71E-07	+++	0.22	childhood asthma
9q21.13	rs75260654 rs116849664	<b>ANXA1</b> , TMC1, LOC101927258, ALDH1A1	c(0.98)/t c(0.98)/t	-0.90	0.16	6.66E-09 1.99E-08		0.05	<b>ANXA1</b> : FEV <sub>1</sub> /FVC, response to bronchodilators in smokers, with anti-inflammatory properties, strongly expressed in bronchial mast cells and potentially involved in epithelial airway

									repair
10q24.2	rs7088157	LOXL4, R3HCC1L	g(0.5)/a	-0.23	0.05	7.34E-06		0.26	R3HCC1L: eosinophil count, atopic eczema, psoriasis, BMI
11p15.4	rs112474574	TRIM5, TRIM6, TRIM22	c(0.96)/t	-0.55	0.12	2.29E-06		0.14	severe asthma and insulin resistance
11q23.3	rs116861530 <sup>eQTL</sup>	SIK3	g(0.94)/a	-0.42	0.09	9.07E-06		0.01	triglycerides, glucose metabolism, eosinophil count
14q22.1	rs1105683	KTN1	c(0.07)/t	0.41	0.09	9.15E-06	+++	0.24	severe asthma
5q13.3	rs2202714 <sup>eQTL</sup>	FAM227B	g(0.36)/a	0.23	0.05	8.71E-06	+++	0.01	$rs35251997$ and $FEV_1$ ; $FEV_1/FVC$
15q25.2	rs117540214 <sup>eQTL</sup>	ADAMTSL3	g(0.06)/a	0.42	0.10	9.82E-06	+++	3.91E-03	FEV <sub>1</sub> /FVC
17q12	rs17676191	11/752	g(0.10)/a	0.36	0.08	2.18E-06	+++	3.06E-03	
	rs79026872	IKZF3	c(0.03)/t	0.64	0.13	2.08E-06	+++	2.56E-03	
	rs4795400		c(0.53)/t	0.30	0.05	5.42E-09	+++	1.96E-04	
	rs1031460	GSDMB	g(0.50)/t	0.27	0.05	8.71E-08	+++	1.87E-04	
	rs56199421		c(0.45)/t	-0.23	0.05	4.50E-06		9.61E-04	early-onset asthma, persistent wheezing (chr17q12-q21)
17q21	rs4795406	LRRC3C	g(0.55)/c	-0.24	0.05	9.91E-07		1.51E-03	Wilestin's (om 1, 411 411)
	rs72832972		c(0.92)/t	-0.38	0.08	8.91E-06		0.01	
	rs4794821	GSDMA	c(0.47)/t	0.27	0.05	9.43E-08	+++	1.07E-03	
	rs59843584		c(0.78)/a	-0.31	0.06	6.38E-08		6.63E-03	
	rs4804311		g(0.08)/a	0.42	0.09	9.65E-07	+-+	0.05	
19p13.2	rs2013694	MARCH2,	c(0.89)/t	-0.38	0.08	8.29E-07	+	0.39	triglycerides, HDL-cholesterol, metabolic syndrome; <i>MYO1F</i> :
19μ13.2	rs73501545	HNRNPM, MYO1F	g(0.16)/a	0.31	0.07	8.39E-06	+++	0.29	FEV <sub>1</sub> and FVC
	rs111644945		g(0.9)/a	-0.41	0.08	4.01E-07		0.02	
22q11.1	rs5994170	CECR5	g(0.4)/a	0.23	0.05	4.95E-06	+++	0.58	triglycerides, eosinophil count
22411.1	rs34902370	CECNS	c(0.75)/t	-0.25	0.06	6.80E-06		0.41	and body height

# **Early-onset Pre-school Remitting Wheezing**

Locus	SNP	Nearby Genes (SNPnexus)	coded(freq)/ other allele	Beta	SE	P-value	Direction	min_pval _other	Previous relevant associations
1q32.3	rs12730098 <sup>eQTL</sup>	PPP2R5A	c(0.79)/t	-0.22	0.05	8.44E-06		0.53	waist circumference & obesity
2,24.2	rs2880066	FAM49A or	t(0.09)/a	0.32	0.07	4.34E-06	+++	0.20	
2p24.2	rs10180268	CYRIA	c(0.06)/t	0.43	0.09	6.56E-07	+++	0.19	airway repair in non-atopic asthma
	rs3861377	NLGN1	g(0.89)/a	-0.28	0.06	7.75E-06		0.28	smoking
3q26.31	rs10513743	NAALADL2	c(0.84)/t	-0.25	0.06	4.97E-06	-+-	0.06	Exacerbations requiring hospitalisation in asthma-suggestive pvalue
5q13.3	rs10075253	SV2C	c(0.85)/t	-0.27	0.06	1.20E-06		0.17	ВМІ
6q27	rs2453395	PDE10A	g(0.33)/a	0.19	0.04	9.51E-06	+++	0.01	Birthweight; asthma and BMI
	rs4730561		g(0.36)/a	-0.20	0.04	6.78E-06		0.13	
7q21.11	rs73144976	MAGI2	g(0.97)/a	-0.47	0.11	9.41E-06		0.26	allergic diseases & atopy, smoking, BMI, airway wall thickness
•	rs67259321		c(0.06)/t	0.43	0.08	1.65E-07	+-+	0.76	,
9p13.3	rs10758259 <sup>eQTL</sup>	C9orf24	g(0.17)/a	-0.27	0.06	4.64E-06		0.01	airway repair
11q22.3	rs72994149	GUCY1A2	c(0.84)/t	-0.24	0.05	8.33E-06	-+-	0.06	systolic blood pressure
13q21.1	rs2872948	PRR20A/B/C/D/E	t(0.96)/a	-0.54	0.10	5.93E-08		0.27	systolic blood pressure
13421.1	rs73527654	FMM2UA/B/C/D/E	g(0.08)/a	0.34	0.07	2.85E-06	+++	0.41	systolic blood pressure
15a21 1	rs116966886	SEMA6D	g(0.99)/a	-0.82	0.18	7.55E-06	-+-	0.57	smoking
13921.1	<i>15q21.1</i> rs117565527	SEMA6D	g(0.99)/a	-0.87	0.17	2.38E-07	-+-	0.43	SHOKING

# **Early-onset Mid-childhood Remitting Wheezing**

Locus	SNP	Nearby Genes (SNPnexus)	coded(freq)/ other allele	Beta	SE	P-value	direction	min_pval _other	Previous relevant associations
	rs35725789	CADM3,	c(0.95)/a	-0.56	0.12	5.42E-06	-+-	0.01	neutrophil count, CRP
1q23.2	rs146141555	FCER1A, MPTX1,	c(0.98)/t	-0.89	0.17	2.04E-07	-+-	0.08	
	rs146575092	OR10J1	g(0.98)/a	-0.85	0.17	8.73E-07	-+-	0.07	
2p22.3	rs7595553	MRPL50P1	g(0.16)/c	-0.46	0.10	3.26E-06		0.12	PM 2.5 exposure level and global DNA methylation level
3p25.3	rs34315999 <sup>eQTL</sup>	RAD18	c(0.03)/t	0.69	0.14	1.11E-06	+++	0.14	atopy/SPT
		MRPL50P1,							<i>3q29:</i> BMI
3q29	rs146961758	LSG1, TMEM44-AS1, TMEM44, ATP13A3	t(0.05)/a	0.57	0.12	6.01E-06	+-+	0.11	TMEM44-AS1, TMEM44, ATP13A3: diastolic blood pressure; LSG1: BMI, eosinophil count
4q24	rs138794367	SLC9B1	c(0.99)/t	-1.02	0.22	5.47E-06		0.13	eosinophil count, allergic rhinitis
5q14.1	rs115719402	AP3B1	g(0.96)/a	-0.60	0.13	7.20E-06		0.06	Vital capacity, BMI
	rs9602218		c(0.06)/a	0.58	0.12	1.74E-06	+-+	0.05	
	rs61960366	RNU6-67P, SLITRK1	g(0.97)/a	-0.79	0.15	7.09E-08	-+-	0.12	
13q31.1	rs74589927		g(0.02)/a	0.73	0.16	3.78E-06	+-+	0.02	RNU6-67P/ rs976078: food allergy
	rs2210726	VENTXP2,	c(0.91)/t	-0.47	0.10	1.33E-06		0.02	
	rs4390476	UBE2D3P4, MTND4P1	c(0.08)/a	0.46	0.10	8.81E-06	+++	0.12	
14q24.2	rs117443464	ZFYVE1	g(0.95)/a	-0.57	0.12	4.68E-06	+	0.19	LDL cholesterol and systolic blood pressure
20p12.3- p12.2	rs6077514	PLCB4	c(0.88)/t	-0.39	0.09	4.03E-06		0.43	neutrophil count

1p32.2         rs2051039         PPAP2B, PRKAA2         c(0.08)/t         0.47         0.10         6.06E-06         ++         0.08         PRKAA2: lymphocyte count and asthma susceptibility           1p31.1         rs72673642         HMGB1P18         g(0.77)/a         -0.31         0.07         6.25E-06          0.01         smoking, BMI           2q13         rs140983998         ACOXL, BUB1         c(0.98)/t         -0.88         0.19         4.71E-06          0.40         ACOXL: later onset asthma and obesity           2q14.3         rs148008098         AMMECR1L         c(0.96)/t         -0.69         0.15         3.41E-06          0.01         body height, blood protein; growth, bone, and heart alterations           3p24.2         rs4072729         RARB         c(0.03)/t         0.61         0.13         4.20E-06         ++         0.23         FEV1/FVC, adult lung function           3q13.2         rs145629570         NAA50, SIDT1, CO200         c(0.02)/t         0.92         0.18         6.83E-07         +++         0.10         SIDT1: FEV1/FVC; CD200: adult-onset non-allergic asthma           3q23         rs113643470         TFDP2, XRN1         c(0.98)/t         -0.91         0.19         1.68E-06          0.03         XRN1: eosi										
1p32.2 rs2051039	Locus	SNP	•		Beta	SE	P-value	direction	<del>_</del>	Previous relevant associations
1B32.2         rs2051039         PRKAA2         c(0.08)/t         0.47         0.10         6.06E-06         ++         0.08         susceptibility           1p31.1         rs72673642         HMGB1P18         g(0.77)/a         -0.31         0.07         6.25E-06          0.01         smoking, BMI           2q13         rs140983998         ACOXL, BUB1         c(0.98)/t         -0.88         0.19         4.71E-06          0.40         ACOXL: later onset asthma and obesity           2q14.3         rs148008098         AMMECRIL         c(0.96)/t         -0.69         0.15         3.41E-06          0.01         body height, blood protein; growth, bone, and heart alterations           3p24.2         rs4072729         RARB         c(0.03)/t         0.61         0.13         4.20E-06          0.23         FEV1/FVC, adult lung function           3q13.2         rs145629570         KIAA2018, NAA50, SIDT1, CD200         c(0.02)/t         0.92         0.18         6.83E-07         +++         0.10         SIDT1: FEV1/FVC; CD200: adult-onset non-allergic asthma           3q23         rs113643470         TFDP2, XRN1         c(0.98)/t         -0.91         0.19         1.68E-06          0.03         XRN1: eosinophil count; 3q23: allergic dise	1p36.13	rs9439669	KLHDC7A	t(0.82)/a	-0.34	0.07	5.15E-06		0.31	1p36.13: metabolic syndrome
2q13 rs140983998 ACOXL, BUB1 c(0.98)/t -0.88 0.19 4.71E-06 0.40 ACOXL: later onset asthma and obesity 2q14.3 rs148008098 AMMECR1L c(0.96)/t -0.69 0.15 3.41E-06 0.01 body height, blood protein; growth, bone, and heart alterations 3p24.2 rs4072729 RARB c(0.03)/t 0.61 0.13 4.20E-06 +-+ 0.23 FEV1/FVC, adult lung function 3q13.2 rs145629570 KIAA2018, NAA50, SIDT1, CD200 0.18 6.83E-07 +++ 0.10 SIDT1: FEV1/FVC; CD200: adult-onset non-allergic asthma 3q23 rs113643470 TFDP2, XRN1 c(0.98)/t -0.91 0.19 1.68E-06 0.03 XRN1: eosinophil count; 3q23: allergic disease and atopic sensitisation 4p11 rs17472015 SLAIN2, SLC10A4, FRYL c(0.01)/t 1.00 0.23 9.49E-06 +++ 0.46 FRYL: body height, age at menopause 7q36.1 rs118027705 GIMAP family, r(0.97)/t -0.77 0.17 6.48E-06 -+- 0.18 systolic blood pressure	1p32.2	rs2051039	,	c(0.08)/t	0.47	0.10	6.06E-06	+-+	0.08	, , ,
2q14.3 rs148008098 AMMECR1L c(0.96)/t -0.69 0.15 3.41E-06 0.01 body height, blood protein; growth, bone, and heart alterations  3p24.2 rs4072729 RARB c(0.03)/t 0.61 0.13 4.20E-06 ++ 0.23 FEV1/FVC, adult lung function  3q13.2 rs145629570 KIAA2018, NAA50, SIDT1, CD200 0.18 6.83E-07 +++ 0.10 SIDT1: FEV1/FVC; CD200: adult-onset non-allergic asthma  3q23 rs113643470 TFDP2, XRN1 c(0.98)/t -0.91 0.19 1.68E-06 0.03 XRN1: eosinophil count; 3q23: allergic disease and atopic sensitisation  4p11 rs17472015 SLAIN2, SLC10A4, FRYL c(0.01)/t 1.00 0.23 9.49E-06 +++ 0.46 FRYL: body height, age at menopause  7q36.1 rs118027705 GIMAP family, c(0.97)/t -0.77 0.17 6.48E-06 -+- 0.01 AOC1: CV disease, smoking; GIMAP family:	1p31.1	rs72673642	HMGB1P18	g(0.77)/a	-0.31	0.07	6.25E-06		0.01	smoking, BMI
3p24.2         rs48008098         AMMECRIL         c(0.96)/t         -0.69         0.15         3.41E-06          0.01         and heart alterations           3p24.2         rs4072729         RARB         c(0.03)/t         0.61         0.13         4.20E-06         +-+         0.23         FEV1/FVC, adult lung function           3q13.2         rs145629570         KIAA2018, NAA50, SIDT1, CD200         c(0.02)/t         0.92         0.18         6.83E-07         +++         0.10         SIDT1: FEV1/FVC; CD200: adult-onset non-allergic asthma           3q23         rs113643470         TFDP2, XRN1         c(0.98)/t         -0.91         0.19         1.68E-06          0.03         XRN1: eosinophil count; 3q23: allergic disease and atopic sensitisation           4p11         rs17472015         SLAIN2, SLC10A4, FRYL         c(0.01)/t         1.00         0.23         9.49E-06         +++         0.46         FRYL: body height, age at menopause           rs117660982         KRBA1, ZNF467         g(0.97)/a         -0.74         0.16         7.63E-06         -+-         0.18         systolic blood pressure           7q36.1         rs118027705         GIMAP family, c(0.97)/t         -0.77         0.17         6.48E-06         -+-         0.01         AOC1: CV disease, smoking; GIMAP fam	2q13	rs140983998	ACOXL, BUB1	c(0.98)/t	-0.88	0.19	4.71E-06		0.40	ACOXL: later onset asthma and obesity
3q13.2 rs145629570	2q14.3	rs148008098	AMMECR1L	c(0.96)/t	-0.69	0.15	3.41E-06		0.01	
3q13.2       rs145629570       NAA50, SIDT1, CD200       c(0.02)/t       0.92       0.18       6.83E-07       +++       0.10       SIDT1: FEVT/FVC; CD200: adult-onset non-allergic asthma         3q23       rs113643470       TFDP2, XRN1       c(0.98)/t       -0.91       0.19       1.68E-06        0.03       XRN1: eosinophil count; 3q23: allergic disease and atopic sensitisation         4p11       rs17472015       SLAIN2, SLC10A4, FRYL       c(0.01)/t       1.00       0.23       9.49E-06       +++       0.46       FRYL: body height, age at menopause         rs117660982       KRBA1, ZNF467       g(0.97)/a       -0.74       0.16       7.63E-06       -+-       0.18       systolic blood pressure         7q36.1       rs118027705       GIMAP family, C(0.97)/t       -0.77       0.17       6.48E-06       -+-       0.01       AOC1: CV disease, smoking; GIMAP family:	3p24.2	rs4072729	RARB	c(0.03)/t	0.61	0.13	4.20E-06	+-+	0.23	FEV1/FVC, adult lung function
7q36.1  rs113643470  rs1146470  rs1146470  rs1146470  rs1146470  rs1146470  rs11464	3q13.2	rs145629570	NAA50, SIDT1,	c(0.02)/t	0.92	0.18	6.83E-07	+++	0.10	• •
rs17472015 SLC10A4, FRYL C(0.01)/t 1.00 0.23 9.49E-06 +++ 0.46 FRYL: body height, age at menopause rs117660982 KRBA1, ZNF467 g(0.97)/a -0.74 0.16 7.63E-06 -+- 0.18 systolic blood pressure rs118027705 GIMAP family, c(0.97)/t -0.77 0.17 6.48E-06 -+- 0.01 AOC1: CV disease, smoking; GIMAP family:	3q23	rs113643470	TFDP2, XRN1	c(0.98)/t	-0.91	0.19	1.68E-06		0.03	
7q36.1  rs118027705 GIMAP family, c(0.97)/t -0.77 0.17 6.48F-06 -+- 0.01 AOC1: CV disease, smoking; GIMAP family:	4p11	rs17472015	· · · · · · · · · · · · · · · · · · ·	c(0.01)/t	1.00	0.23	9.49E-06	+++	0.46	FRYL: body height, age at menopause
rs118027705 GIMAP family, c(0.97)/t -0.77 0.17 6.48F-06 -+- 0.01 AOC1: CV disease, smoking; GIMAP family:		rs117660982	KRBA1, ZNF467	g(0.97)/a	-0.74	0.16	7.63E-06	-+-	0.18	systolic blood pressure
	7q36.1	rs118027705		c(0.97)/t	-0.77	0.17	6.48E-06	-+-	0.01	•


	rs139489493	LOC105375566	c(0.98)/t	-0.95	0.20	2.28E-06		0.03	
7q36.3	rs144271668	PTPRN2	c(0.01)/a	0.88	0.19	2.91E-06	+++	0.28	eczema
8q21.3	rs990182	LOC105375631	t(0.42)/a	0.28	0.06	2.57E-06	+++	0.46	8q21.3: type 1 diabetes
9p22.3	rs79110962	NFIB, ZDHHC21	c(0.08)/t	0.51	0.10	3.98E-07	+++	0.05	9p22.3: asthma (mean age<16 years)
10q23.31	rs7896106	SLC16A12, IFIT family, PANK1	g(0.35)/t	0.30	0.06	1.35E-06	+++	0.05	SLC16A12: Body height; PANK1: insulin
11q23.3	rs141958628	CBL, CCDC84, MCAM	c(0.98)/t	-0.98	0.20	1.33E-06	-+-	0.27	CCDC84: asthma, allergy
15q15.3- q21.1	rs139134265	SPG11, CTDSPL2	g(0.02)/c	0.87	0.20	9.11E-06	+-+	0.13	CTDSPL2: alcohol drinking
15q25.2	rs143862030	ADAMTSL3, GOLGA6L4, UBE2Q2P8	c(0.04)/t	0.64	0.13	1.65E-06	+++	0.08	ADAMTSL3: FEV1/FVC; lean mass
16p13.3	rs113390367	SSTR5-AS1, CACNA1H	g(0.86)/a	-0.40	0.08	1.04E-06		0.16	CACNA1H: eosinophil count
16p12.1	rs4788025	GSG1L	g(0.46)/a	-0.30	0.06	7.99E-07		0.19	16p12.1: current asthma and rhinoconjunctivitis at 10-15 years
22q13.32	rs133498	FAM19A5 or TAFA5	g(0.94)/a	-0.48	0.11	5.35E-06		0.84	Obesity and Metabolic Dysfunction


<sup>\*</sup>minimum p-value across associations with the other three wheezing phenotypes, using the never/infrequent wheeze as the baseline phenotype

eQTL: identified in expression analyses of whole blood and/or lung tissues using Genotype-Tissue Expression database (https://gtexportal.org) using the European reference panel.


<sup>428 †</sup> List of references or sources (GeneCards, GWAS Catalog, PhenoScanner) available in **Appendix 5-Tables 1-4.** 


**Figure 1.** Circular Manhattan plot showing an overview of the GWAS results by wheeze phenotype (from outside to inside: early-onset persistent, early-onset pre-school remitting, early-onset mid-childhood remitting and late-onset wheeze). The red line indicates the genome-wide significance threshold ( $P < 5 \times 10^{-8}$ ), while the blue line indicates the threshold for genetic variants that showed a suggestive significant association ( $P < 10^{-5}$ ).





**Figure 2.** Scatter plots illustrating the heterogeneity in the genetic profile of the wheezing phenotypes. Top plots compare phenotype-specific beta effects for persistent and early-onset mid-childhood remitting wheezing. Shared nominal beta effects only found when relaxing p< $10^{-4}$  for early-onset mid-childhood remitting wheezing. Bottom plots compare phenotype-specific beta effects for persistent and early-onset pre-school remitting. No shared beta effects (same or opposite direction) were found at p< $10^{-5}$  for any of the comparisons. Abbreviations used: pw=persistent, er= early-onset pre-school remitting and pe= early-onset mid-childhood remitting.






445

446 447

448

# Figure 3. Chromatin interactions between rs75260654 and the ANXA1 promoter in CD4+ cells in MAAS



rs75260654 physically interacts with ANXA1 promoter in CD4+ T-cells and overlaps a region of active (POLR2AphosphoS2 ChIP-seq) open (ATAC-seq) chromatin in A549 cell line (lung epithelial carcinoma). The region is also predicted to be an active enhancer (ChromHMM 18-state model) in the A549 cell type. Only ChromHMM enhancer chromatin are displayed. Yellow shaded area indicates the PCHi-C fragment overlapping rs75260654 (red bar) and interacting with the ANXA1 promoter.

**Figure 4.** eQTL *ANXA1* and rs75260654 across different tissue types. Point size is proportional to -log10 p-value.

## Tissue specfic effect of rs75260654 on ANXA1 Expression





Figure 5. Annexin A1 is induced following HDM challenge and mice deficient in *ANXA1* have exacerbated airway hyperreactivity. (A) Schematic of house dust mite allergen dosing protocol, N=4-6 per group, data representative of two animal experiments. (B, C) Immunofluorescent staining of paraffin embedded lung tissue sections incubated with anti-Annexin A1, counterstained with DAPI (N=4 per group). (D) mRNA expression of annexin A1 in lung tissue following HDM exposure, expression normalized to housekeeping gene hprt. Mice receiving HDM were analysed for changes in airway hyper-reactivity following methacholine (MCh) challenge in tracheotomized restrained mice. (E) Airway resistance at top MCh dose 100mg/ml (F) Eosinophils quantified in BAL, (F) T1/ST2+ lymphocytes quantified in the BAL. (H) IL-4 (I) IL-5 and (J) IL-13 quantified in lung tissue by ELISA. \* p<0.05 and \*\* p<0.01 relative to PBS control group by Mann Whitney test. + p<0.05 and ++ p<0.01 comparing HDM AnnexinA1 KO mice relative to HDM WT group by Mann Whitney test.



#### 474 **APPENDIX 1**

502

503

504

## 475 **Description of cohorts** 476 The Study Team for Early Life Asthma Research (STELAR) consortium(20) brings together five 477 UK population-based birth cohorts as described below. Informed consent was obtained from 478 parents, and study subjects gave their assent/consent when applicable. Data were 479 harmonised and imported into Asthma eLab web-based knowledge management platform to 480 facilitate joint analyses (www.asthmaelab.org)(20). 481 The Avon Longitudinal Study of Parents and Children (ALSPAC) 482 ALSPAC is a birth cohort study established in 1991 in Avon, UK(62, 63). Pregnant women with expected dates of delivery 1st April 1991 to 31st December 1992 were invited to take part in 483 484 the study. The initial number of pregnancies enrolled is 14,541. Of these initial pregnancies, 485 there was a total of 14,676 foetuses, resulting in 14,062 live births and 13,988 children who 486 were alive at 1 year of age. 487 When the oldest children were approximately 7 years of age, an attempt was made to bolster 488 the study with eligible cases who had failed to join originally. As a result, when considering 489 variables collected from the age of seven onwards (and potentially abstracted from obstetric 490 notes) there are data available for more than the 14,541 pregnancies mentioned above. The 491 number of new pregnancies not in the initial sample (known as Phase I enrolment) that are 492 currently represented on the built files and reflecting enrolment status at the age of 24 is 913 493 (456, 262 and 195 recruited during Phases II, III and IV respectively), resulting in an additional 494 913 children being enrolled. The phases of enrolment are described in more detail in the 495 cohort profile paper and its update. The total sample size for analyses using any data 496 collected after the age of seven is therefore 15,454 pregnancies, resulting in 15,589 foetuses. 497 Of these 14,901 were alive at 1 year of age. 498 Ethical approval: Ethical approval for the study was obtained from the ALSPAC Ethics and Law 499 Committee and the Local Research Ethics Committees. All self-completion questionnaire 500 content is approved by the ALSPAC Ethics and Law Committee. Ethics protocols' numbers: 501 Initial approval Bristol and Weston Health Authority: E1808 Children of the Nineties: Avon

Longitudinal Study of Pregnancy and Childhood (ALSPAC). (28th November 1989). Southmead

Health Authority: 49/89 Children of the Nineties -"ALSPAC". (5th April 1990). Frenchay Health

Authority: 90/8 Children of the Nineties. (28th June 1990).

- 505 Informed consent for the use of data collected via questionnaires and clinics was obtained
- 506 from participants following the recommendations of the ALSPAC Ethics and Law Committee at
- 507 the time. Data dictionary: The study website contains details of available data through a fully
- 508 searchable data dictionary: http://www.bristol.ac.uk/alspac/researchers/our-data/
- 509 We are extremely grateful to all the families who took part in this study, the midwives for
- 510 their help in recruiting them, and the whole ALSPAC team, which includes interviewers,
- 511 computer and laboratory technicians, clerical workers, research scientists, volunteers,
- 512 managers, receptionists and nurses.
- 513 The Manchester Asthma and Allergy Study (MAAS)
- 514 MAAS is an unselected birth cohort study established in 1995 in Manchester, UK(24). It
- consists of a mixed urban-rural population within 50 square miles of South Manchester and
- 516 Cheshire, United Kingdom located within the maternity catchment area of Wythenshawe and
- 517 Stepping Hill Hospitals. All pregnant women were screened for eligibility at antenatal visits (8-
- 518 10<sup>th</sup> week of pregnancy). Of the 1499 couples who met the inclusion criteria (≤10 weeks of
- 519 pregnancy, maternal age ≥18 years, and questionnaire and skin prick data test available for
- both parents), 288 declined to take part in the study and 27 were lost to follow-up between
- 521 recruitment and the birth of a child. A total of 1184 children were born into the study
- between February 1996 and April 1998. They were followed prospectively for 19 years to date
- and attended follow-up clinics for assessments, which included lung function measurements,
- 524 skin prick testing, biological samples (serum, plasma and urine), and questionnaire data
- 525 collection. The study was approved by the North West Greater Manchester East Research
- 526 Ethics Committee. Ethics protocols' numbers: ERP/94/032 Up to 5 yrs. Allergen avoidance,
- 527 Primary Prevention, genetics, sRaw age 3 and 5; SOU/00/259 5 year; ERP/95/137 Exposure to
- 528 pet allergens, atopy, genetics; ERP/97/023 IFWIN, genetics; 03/SM/400 8 year; 06/Q1403/142
- 529 10-12 years; 11/NW/0228 13-15 years; 14/NW/1309 18+ years.
- 530 The Study of Eczema and Asthma to Observe the influence of Nutrition (SEATON)
- 531 SEATON is an unselected birth cohort study established in 1997 in Aberdeen, UK, which was
- 532 designed to explore the relationship between antenatal dietary exposures and asthma
- outcomes in childhood(64). 2000 healthy pregnant women attending an antenatal clinic, at
- median 12 weeks gestation, were recruited. An interviewer administered a questionnaire to
- 535 the women and atopic status was ascertained by skin prick test (SPT). The cohort included

1924 children born between April 1998 and December 1999. Participants were recruited prenatally and followed up by self–completion questionnaire to 15 years of age using postal questionnaires to record the presence of asthma and allergic diseases. Lung function measurements and SPT to common allergens was performed at 5, 10 and 15 years. The study was approved by the North of Scotland Research Ethics Committee. Ethics protocol REC reference: 13/NS/0108; Protocol number: 2/048/13; Amendment number: AM03.

542 ASHFORD

The Ashford study is an unselected birth cohort study established in 1991 in Ashford, UK(65). It included 642 children born between 1992 and 1993. Participants were recruited prenatally and followed to age 14 years. Detailed standardised questionnaires were administered at each follow-up to collect information on the natural history of asthma and other allergic diseases. Lung function measurements and SPT was carried out at 5, 8 and 14 years of age. In 2015, the study children aged 20 were sent a self-completion questionnaire, which was returned by 60% of the participants. The Asthma in Ashford study was reviewed by the Imperial College Research Ethics Committee on 11/11/2014. On 08/01/2015 the Joint Research Compliance Office granted full approval of the study on the basis described in the revised documents. ICREC reference: 14|C2288.

553 The Isle of Wight (IOW) cohort

IOW is an unselected birth cohort study established in 1989 on the Isle of Wight, UK(23, 66, 67). After the exclusion of adoptions, perinatal deaths, and refusal for follow-up, written informed consent was obtained from parents to enrol 1,456 newborns born between 1<sup>st</sup> January 1989 and 28<sup>th</sup> February 1990. Follow-up-up assessments were conducted to 26 years of age to prospectively study the development of asthma and allergic diseases. At each follow-up, validated questionnaires were completed by the parents. Additionally, the Skin Prick Test (SPT) was performed on 980, 1036 and 853 participants at 4, 10 and 18 years of age to check allergic reactions to common allergens. At 10, 18, and 26 years, spirometry and methacholine challenge tests were performed to diagnose lung problems. Ethics protocols' numbers: Ethics approval for the IoW cohort was originally given by the Isle of Wight Local Research Ethics Committee (now named the National Research Ethics Service, NRES Committee South Central – Southampton B) in 1989 and at each subsequent follow up (1,2 and 4 years) (this is pre "numbers"); Age 10 follow up (including DNA and genotyping): Isle of

- 567 Wight Health Authority Local Research Ethics Committee 18/98; Age 18 Follow up(including
- 568 DNA and genotyping): Isle of Wight, Portsmouth & South East Hampshire Research Ethics
- 569 Committee 06/Q1701/34.
- 570 **Definition of variables**
- A list of all variables used in the current study, per cohort, is shown in **Appendix 1-Table 1**.
- 572 Demographic, exposures and outcomes
- 573 Postal questionnaires were used in ALSPAC and SEATON, while interviewer-administered
- 574 questionnaires were employed in other cohorts.
- 575 Parental history of asthma, eczema and hay fever was defined based on the responses given
- 576 to the question "have you (and/or your partner) ever had asthma/eczema/hay fever".
- 577 Maternal and paternal smoking were defined based on the response given to the question
- 578 "do you (or does your partner) smoke", administered during pregnancy. Low birth weight was
- defined as birth weight less than 2500 g based on NHS birth records.
- Asthma in MAAS was defined as a case if positive for two of the following criteria: doctor
- diagnosis of asthma in the past 12 months, current wheeze in the last 12 months, doctor
- 582 prescription for asthma. Asthma in ALSPAC was defined as a mothers' report of doctor ever
- 583 diagnosis of asthma.
- Current wheeze in MAAS was defined as a questionnaire report to the question "have you
- 585 wheezed in the last 12 months" upon attendance at a follow up clinic. Current wheeze in
- 586 ALSPAC was defined as a mothers' report to the question "has your child had any wheezing or
- 587 whistling in the last 12 months?".
- Asthma medication in ALSPAC was defined as a mothers' report to the question "has your
- 589 child taken any asthma medication in the last 12 months?". Lower respiratory hospital
- 590 admissions: Data on hospital admissions in MAAS were obtained by manually inspecting the
- 591 General Practice (GP) records for each individual.
- 592 Early-life risk factors were divided into four groups according to timing of exposure; maternal
- and child characteristics (gender, maternal smoking during pregnancy and maternal history of
- 594 asthma), perinatal (low birth weight adjusted for gestational age), environmental (pet
- ownership, smoke exposure after birth) and allergic sensitization (defined based on positive
- 596 skin prick test to cat, house dust mite or grass) variables.

- 597 Primary outcome: Joint wheeze phenotypes
- 598 We used latent class analysis (LCA) to identify longitudinal trajectories of wheeze(19) based
- on pooled analysis among 15,941 children with at least two observations on wheezing at five
- time periods that were approximately shared across all cohorts: infancy (1/2-1 year); early
- 601 childhood (2-3 years); pre-school/early school age (4-5 years); middle childhood (8-10 years);
- and adolescence (14-18 years). Cohort-specific definitions other variables derived from the
- questionnaires are provided in **Appendix 1-Table 2**.
- To control for cohort-specific variation, Cohort ID was included in the LCA model as an
- 605 additional predictor by transforming the 5-category variable into a set of four dummy
- of variables and including them as covariates. The largest cohort, ALSPAC, was treated as the
- 607 non-coded category to which all other cohorts were compared. The expectation
- 608 maximization algorithm was used to estimate relevant parameters, with 100,000 iterations
- and 500 replications.
- To assess model fit, we used (1) the Bayesian information criterion (BIC), (2) the Akaike
- 611 information criterion (AIC), (3) Lo-Mendell-Rubin likelihood ratio test (LMR), (4)
- Bootstrapped likelihood ratio and, (4) quality of classification certainty (model entropy). The
- 613 BIC is an index used in Bayesian statistics to choose among a set of competing models; the
- model with the lowest BIC is preferred. Using the lowest BIC as a selection criterion, the best
- 615 fitting model was chosen as the five-class solution with a nominal covariate (BIC:31340).
- 616 Analyses were carried out using Mplus 8, R (http://www.r-project.org/) and Stata 14
- 617 (StataCorp, College Station, Tex).
- 618 Based on the statistical fit, a five-class solution was selected as the optimal model(19), and
- the classes (wheeze phenotypes) were labeled as: (1) Never/Infrequent wheeze (52.4%); (2)
- 620 Early-onset pre-school remitting wheeze (18.6%), with high prevalence of wheeze during
- 621 infancy, decreasing to 20% around early-childhood and to less than 10% afterwards; (3) Early-
- 622 onset middle-childhood remitting wheeze (9.8%), with early-onset wheeze and peak
- 623 prevalence in early-childhood (~70%), and diminishing by middle-childhood (<5%); (4) Early-
- 624 onset persistent wheeze (10.4%) with 58% wheeze prevalence during infancy, and prevalence
- between 70- 80% thereafter; (5) Late-onset wheeze (8.8%) with very low prevalence until
- 626 middle childhood, increasing rapidly to 55% in adolescence. These latent classes were used in
- 627 the subsequent GWAS.

- 628 Minimising Bias and missing data effects
- 629 Extracted from reference Oskel et al.(19): "One of the advantages of our multicohort
- 630 approach is that individual studies that might not provide conclusive evidence to make
- 631 inference about the general population because of cohort specific effects and biases can
- 632 contribute to revealing a more accurate picture when integrated together. The integration of
- 633 five cohorts and their pooled analysis enhanced the credibility and generalizability of the
- 634 phenotyping results to the U.K. population. A further advantage is to minimize the study-
- 635 specific biases (including cohort specific effects, attrition effects, different recruitment
- 636 strategies, and geographic factors) affecting the certainty of allocation of individuals to each
- 637 latent class, while maximizing the benefits of individual cohort studies (e.g., potentially
- 638 important risk factors and outcomes are captured in some, but not all cohorts)."
- 639 "Another strength of pooling cohort data is that a multicohort design allowed us to analyze a
- 640 large sample with complete data on wheeze from birth to adolescence, thus increasing
- 641 statistical power to detect less prevalent phenotypes." However, "The optimal solution in the
- 642 model using 15,941 children (allowing for missing data) remained five classes (see Table E3,
- 643 Figure E1), and was very similar to that derived from a complete data set." We used results
- from the larger sample, that is individuals with at least 2 observations of wheezing, to assign
- 645 individuals to their most likely wheezing phenotype and used this as our primary outcome in
- 646 this study.
- 647 Included vs. excluded participants
- Related and non-European individuals were excluded as well as those individuals with missing
- 649 genetic data.
- In ALSPAC, 11,176 individuals had data on wheezing phenotypes, of these 6,833 were white
- 651 unrelated and had genetic data. We found more children from mothers who smoked during
- pregnancy in the excluded sample compared to the included sample; no difference in gender,
- 653 maternal history of asthma, current wheezing at 8 or 15 years, and small evidence for more
- asthma ever and current medication at 8 years in the excluded sample (Appendix 1-Table 4).
- 655 In MAAS, 1150 individuals had data on wheezing phenotypes, of these 887 were white
- 656 unrelated and had genetic data. We found no difference in children from mothers who
- 657 smoked during pregnancy in the excluded sample compared to the included sample; no
- difference in gender, maternal history of asthma or current wheezing at both 8 and 16 years.

659 There was small evidence for more asthma ever and current medication at 8 years in the 660 excluded sample (Appendix 1-Table 4). 661 In SEATON, 1535 individuals had data on joint wheezing phenotypes, of these 548 were white 662 unrelated and had genetic data. We found evidence for more children from mothers who 663 smoked during pregnancy in the excluded sample compared to the included sample; and 664 more males in the excluded sample. There was no difference in maternal history of asthma or 665 current wheezing, asthma ever or current medication at both 10 and 15 years in the excluded 666 sample compared to the included sample (Appendix 1-Table 4). 667 In ASHFORD, 620 individuals had data on joint wheezing phenotypes, of these 348 were white 668 unrelated and had genetic data. We found evidence for more children from mothers who 669 smoked during pregnancy in the excluded sample compared to the included sample; no 670 difference in gender, maternal history of asthma or asthma ever. There was small evidence 671 for less current wheezing at 8 years, or current medication at 8 years in the excluded sample 672 compared to the included sample (Appendix 1-Table 4). 673 In IOW, 1460 individuals had data on joint wheezing phenotypes, of these 952 were white 674 unrelated and had genetic data. We found evidence for more children from mothers who 675 smoked during pregnancy in the excluded sample compared to the included sample; no 676 difference in gender, maternal history of asthma, asthma ever at 10 and 18 years in the 677 excluded sample compared to the included sample. There was small evidence for more 678 children with current wheeze and medication at 8 years in the included sample compared to 679 the included sample (Appendix 1-Table 4).

# Appendix 1 Table 1. Definition of variables in each of the five STELAR birth cohorts

Variable	Definition
Cohort: ALSPAC	
Mother-asthma	Have you ever had asthma? (recruitment)
Mother smoking	Mother smoked when expecting (recruitment)
Doctor-diagnosed asthma ever	Has a doctor ever said that your child has asthma? (year 8 and 14)
Current wheezing	Two questions combined: Occurrence of wheezing and/or wheezing with whistling on the chest in the last 12 months (year $\frac{1}{2}$ , $4^{\frac{3}{4}}$ , $8^{\frac{1}{2}}$ and 14)
Current asthma medication	Asthma medication in the last 12 months (year 8 <sup>1/2</sup> and 14)
Current rhinitis	Child had problem with sneezing/runny nose without cold/flu in last 12 months (year 7 and $16^{1/2}$ )
Current hay-fever	Child had hay-fever in last 12 months (year 10 <sup>1/2</sup> and 14)
Cohort: MAAS	
Mother-asthma	Has a doctor ever told you that you had asthma? (recruitment)
Mother smoking	Do you smoke- mother (recruitment)
Doctor-diagnosed asthma ever	Has your doctor ever told you that your child has or had asthma? (year 8 & 16)
Asthma ever	Has your child ever suffered from asthma (year 8 and 16)
Current wheezing	Has your child had wheezing or whistling in the chest in the last 6/12 months (year 1, 3, 5, 8 and 16)
Current asthma medication	Asthma medication in the last 12 months (year 8 and year 16)
Current rhinitis	Has your child ever had a problem with sneezing, or a runny nose, or a blocked nose when he /she did not have a cold or the flu? (year 8 and year 16)
Current hay-fever	Does your child have hay-fever now? (year 8 and year 16)
Cohort: SEATON	

Mother-asthma	Do you suffer from asthma? (recruitment)
Mother smoking	Which of the following best describes your smoking status? (recruitment)
Doctor-diagnosed asthma ever	Has your child ever suffered from asthma? If yes, has this been confirmed by a doctor? (year 10 & 15)
Asthma ever	Has your child ever suffered from asthma? (year 10); Have you ever suffered from asthma? (year 15)
Current wheezing	Has your child had wheezing in the chest in the last 12 months (year 1, 2, 5, 10 and 15)
Current asthma medication  Current hay-fever	Has your child been prescribed medicines/inhalers for asthma in the last 12 months? (year 10)  Have you been prescribed medicines/inhalers for asthma in the last 12 months? (year 15)  Has your child suffered from hay-fever last 12 months? (year 10 & 15)
Cohort: ASHFORD	<u> </u>
	De very house on house you give heart told you house oothers? / reconsition out.)
Mother-asthma  Mother smoking	Do you have or have you ever been told you have asthma? (recruitment)  Do you smoke cigarettes? (recruitment)
_	
Doctor-diagnosed asthma ever	Has your doctor ever told you that your child has or had asthma? (year 8 & 14)
Asthma ever	In the past 12 months has your daughter suffered from asthma? (year 8); Has she/he ever suffered from asthma? (year 14)
Current wheezing	Which one best describes your child's wheeze in past 12 months? 'Yes' (B:1-6, C:7+), 'No' (A:0) (year 1, 2, 5, 8 and 14)
Current asthma medication	Over the last 12 months has your daughter taken any of the following treatments (preventer, reliever, nebulizer, steroids) for asthma? (year 8 and year 14)
Current rhinitis	In the last twelve months has your child had a problem with sneezing or a runny or blocked nose? (year 8 & 14)
Current hay-fever	In your opinion does your child have hay fever now? year 8
	Has your child ever had hay fever? year 14
Cohort: IOW	·
Mother-asthma	Do you or have you suffered from asthma or wheezing (recruitment)
Mother smoking	Do you smoke in the house? (recruitment)
Doctor-diagnosed	Asthma cared for by hospital specialist/ GP or nurse (year 10, 18 & 26)

#### asthma ever

Asthma ever Child ever had asthma (year 10 and 18)

**Current wheezing** Presence of wheeze since previous review (year 1, 2, 4, 10 and 18)

Asthma medication Child ever had asthma treatment (year 18 year) combined with asthma treatment questions

ever being asked at year 1, 2, 4, 10 and 18

Current rhinitis In the past 12 months have you had a problem with sneezing, or a runny or blocked nose when

you did not have a cold or the flu? (year 10, 18 & 26)

# 685 wheeze phenotypes

Birth Cohort:	IOW	MAAS	SEATON	ASHFORD	ALSPAC
Year of birth	1989	1995	1997	1992	1991
Questionnaire	Interviewer -administered	Interviewer -administered	Postal	Interviewer –administered	Postal
Data collection age (years)	1, 2, 4, 10, 18	1, 3, 5, 8, 16	1, 2, 5, 10, 15	1, 2, 5, 8, 14	1/2, 2 <sup>1/2</sup> , 4 <sup>3/4</sup> , 8 <sup>1/2</sup> , 14
No. of children with ≥2 observations on wheezing at five selected time points	1460	1150	1535	620	11176

Appendix 1 Table 3. Characteristics of the participants in STELAR cohorts included in this analysis (restricted to individuals with genetic data).

Numbers are N (%) except for age, where we report Mean (SD).

688

686

	ALSPAC		MAAS		SEATON		ASHFORD		IOW	
	N=6833		N=887		N=548		N=348		N=952	
	(71.4%)		(9.3%)		(5.7%)		(3.6%)		(9.9%)	
Males	3492 (51.1)		475 (53.6)		260 (47.5)		179 (51.4)		466 (49.0)	
Maternal history of asthma	748 (11.5)		120 (13.5)		77 (14.1)		49 (14.1)		106 (11.2)	
Maternal smoking	1423 (22.1)		122 (13.8)		107 (19.5)		52 (14.9)		217 (23.1)	
Wheeze Phenotypes										
Never/Infrequent	4331 (63.4)		506 (57.1)		332 (60.6)		145 (41.7)		573 (60.2)	
Early-onset persistent	656 (9.6)		133 (15.0)		36 (6.6)		41 (11.8)		77 (8.1)	
Early-onset pre-school remitting	1076 (15.8)		145 (16.4)		117 (21.4)		145 (41.7)		0	
Early-onset mid-childhood remitting	474 (6.9)		48 (5.4)		13 (2.4)		13 (3.7)		55 (5.8)	
Late-onset	296 (4.3)		55 (6.2)		50 (9.1)		4 (1.2)		247 (26.0)	
	7-8 years	14-15 years	8 years	16 years	10 years	15 years	8 years	14 years	10 years	18 years
Age Mean (SD) in years	8.7 (0.3)	15.4 (0.3)	7.98 (0.16)	16.09 (0.62)	10.15 (0.18)	15.09 (0.28)	7.97 (NA)	13.95 (NA)	9.98 (0.27)	17.87 (0.59)
Doctor-Diagnosed Asthma ever*	1060 (19.7)	796 (23.2)	198 (23.9)	198 (30.0)	86 (16.0)	80 (19.5)	75 (21.6)	83 (23.9)	350 (40.9)	255 (28.6)
Asthma ever	NA	NA	193 (22.8)	192 (29.5)	87 (16.2)	66 (21.9)	54 (15.6)	65 (18.7)	194 (20.9)	264 (29.3)
Current wheeze	683 (12.5)	306 (9.0)	150 (17.6)	112 (16.9)	67 (12.4)	63 (15.5)	54 (15.6)	54 (15.5)	190 (20.4)	227 (25.1)
Current asthma medication	695 (12.9)	361 (10.6)	141 (16.5)	114 (17.1)	68 (12.6)	58 (14.0)	50 (14.41)	49 (14.1)	41 (11.81)	38 (10.9)

<sup>\*</sup> DDA ever not available in IOW, we used Asthma cared for by hospital specialist/ GP or nurse as proxy


Appendix 1 Table 4. Comparison of included vs. excluded participants in the five cohorts at different ages

ALSPAC	N	Included	N	Excluded	P-value						
Males (%)	6833	3492 (51.1)	4343	2269 (52.2)	0.24						
Maternal history asthma (%)	6497	748 (11.5)	4038	453 (11.2)	0.64						
Maternal smoking-pregnancy (%)	6438	1423 (22.1)	4019	1167 (29.0)	<0.001						
	At 7.5	-8.5 years				At 14	-15 years				
ALSPAC	N	Included	N	Excluded	P-value	N	Included	N	N	Excluded	P-value
Age Mean (SD) years	5139	8.7 (0.3)	1872	8.7 (0.3)	<0.001	3885	15.4 (0.3)	1	1237	15.5 (0.4)	<0.001
Current wheeze (%)	5453	683 (12.5)	2579	344 (13.3)	0.308	3419	306 (9.0)	1	1078	105 (9.7)	0.432
Asthma ever (%)	5377	1060 (19.7)	2605	562 (21.6)	0.053	3425	796 (23.2	) 1	1079	279 (25.9)	0.079
Current asthma medication (%)	5379	695 (12.9)	2529	368 (14.6)	0.047	3400	361 (10.6	) 1	1077	134 (12.4)	0.096
MAAS	N	Included	N	Excluded	P-value						
Males (%)	887	475 (53.6)	263	149 (56.7)	0.38						
Maternal history asthma (%)	886	120 (13.5)	259	45 (17.4)	0.12						
Maternal smoking* (%)	884	122 (13.8)	260	47 (18.1)	0.09						
	At 8 ye	ears				At 16 ye	ears				
MAAS	N	Included	N	Excluded	P-value	N	Included	N	Exclud	led I	P-value
Age Mean (SD) years	827	7.98 (0.16)	149	8.00 (0.21)	0.31	605	16.09 (0.62)	59	15.98	(0.60)	0.20
Current wheeze (%)	853	150 (17.6)	172	35 (20.4)	0.39	664	112 (16.9)	82	15 (18	.3)	0.11
Asthma ever (%)	845	193 (22.8)	173	52 (30.1)	0.043 47	651	192 (29.5)	79	28 (35	.4)	0.28

Current asthma medication (%)	855	141 (16.5)	173	43 (24.9)	0.009	666	114 (17.1)	83	14 (16.9)	0.96
<u>SEATON</u>	N	Included	N	Excluded	P-value					
Males (%)	548	260 (47.5)	987	525 (53.2)	0.031					
Maternal history asthma (%)	548	77 (14.1)	985	161 (16.4)	0.24					
Maternal smoking* (%)	548	107 (19.5)	987	276 (28.0)	<0.001					
	At 10	years				At 1	L5 years			
SEATON	N	Included	N	Excluded	P-value	N	Included	N	Excluded	P-value
Age Mean (SD) years	548	10.15 (0.18)	987	10.23 (0.16)	<0.001	545	15.09 (0.28)	916	15.11 (0.26)	0.20
Current wheeze (%)	541	67 (12.4)	376	42 (11.2)	0.58	407	63 (15.5)	310	48 (15.5)	0.99
Asthma ever (%)	537	87 (16.2)	374	53 (14.2)	0.40	409	66 (21.9)	302	85 (20.8)	0.73
Current asthma medication (%)	542	68 (12.6)	378	39 (10.3)	0.30	414	58 (14.0)	309	34 (11.0)	0.23
<u>ASHFORD</u>	N	Included	N	Excluded	P-value					
Males (%)	348	179 (51.4)	272	153 (56.3)	0.23					
Maternal history asthma (%)	348	49 (14.1)	272	38 (14.0)	0.97					
Maternal smoking* (%)	348	52 (14.9)	270	61 (22.6)	0.015					
	At 8 y	ears				At 14	years			
ASHFORD	N	Included	N	Excluded	P-value	N	Included	N	Excluded	P-value
Age Mean (SD) years	348	NA	272	NA	NA	348	NA	272	NA	NA
Current wheeze (%)	347	54 (15.6)	246	25 (10.2)	0.06	348	54 (15.5)	150	18 (12.00)	0.31
Asthma ever (%)	347	54 (15.6)	246	38 (15.5)	0.97	348	65 (18.7)	150	25 (16.7)	0.59
Current asthma medication (%)	347	50 (14.41)	246	22 (8.9)	0.05	348	49 (14.1)	150	16 (10.7)	0.30
Isle Of Wight	N	Included	N	Excluded	P-value					

Males (%)	952	466 (49.0)	508	275 (54.1)	0.06					
Maternal history asthma (%)	946	106 (11.2)	505	52 (10.3)	0.60					
Maternal smoking* (%)	941	217 (23.1)	502	147 (29.3)	0.01					
	At 10	At 10 years								
IOW	N	Included	N	Excluded	P-value	N	Included	N	Excluded	P-value
Age Mean (SD) years	932	9.98 (0.27)	426	10.04 (0.31)	<0.001	914	17.87 (0.59)	389	18.14 (0.67)	<0.001
Current wheeze (%)	932	190 (20.4)	426	69 (16.2)	0.07	903	227 (25.1)	377	58 (15.4)	<0.002
Asthma ever (%)	930	194 (20.9)	425	80 (18.8)	0.39	900	264 (29.3)	385	108 (28.1)	0.64
Current asthma medication (%)	347	41 (11.81)	246	15 (6.10)	0.02	348	38 (10.9)	150	13(8.7)	0.45





#### APPENDIX 2

#### Genotyping and imputation

695 ALSPAC

693

- 696 Participants were genotyped using the Illumina HumanHap550 quad genome-wide SNP
- 697 genotyping platform (Illumina Inc., San Diego, CA, USA) by the Wellcome Trust Sanger Institute
- 698 (WTSI; Cambridge, UK) and the Laboratory Corporation of America (LCA, Burlington, NC, USA),
- 699 using support from 23andMe. Haplotypes were estimated using ShapeIT (v2.r644) which uses
- relationship information to improve phasing accuracy. The phased haplotypes were then imputed
- 701 to the Haplotype Reference Consortium (HRCr1.1, 2016) panel(68) of approximately 31,000
- 702 phased whole genomes. The HRC panel was phased using Shapelt v2, and the imputation was
- 703 performed using the Michigan imputation server.
- 704 *MAAS*
- 705 In MAAS, we used the Illumina 610 quad genome-wide SNP genotyping platform (Illumina Inc., San
- 706 Diego, CA, USA). Prior to imputation samples were excluded on the basis of gender mismatches;
- 707 minimal or excessive heterozygosity, genotyping call rates of <97%. SNPS were excluded if they
- 708 had call rates of < 95%, minor allele frequencies of < 0.5% and HWE p<3x10-8. Prior to imputation
- 709 each chromosome was pre-phased using EAGLE2 (v2.0.5)(68) as recommended by the Sanger
- 710 imputation server (69). We then imputed with PBWT (70) with the Haplotype Reference
- 711 Consortium (release 1.1) of 32,470 reference genomes (69) using the Sanger Imputation Server.
- 712 IOW, SEATON and ASHFORD
- 713 IOW, SEATON and ASHFORD were genotyped using the illumina Infinium Omni2.5-8 v1.3 BeadChip
- 714 genotyping platform (Illumina Inc., San Diego, CA, USA). Genotype QC and imputation was carried
- 715 out as described for MAAS.
- 716 Exclusions
- 717 Individuals were excluded on the basis of gender mismatches; minimal or excessive
- 718 heterozygosity; disproportionate levels of individual missingness (>3%), insufficient sample
- 719 replication (IBD < 0.8) or evidence of cryptic relatedness (IBD > 0.1). Following imputation, single
- 720 nucleotide polymorphisms (SNPs) with a minor allele frequency of <1%, a call rate of <95%,
- 721 evidence for violations of Hardy-Weinberg equilibrium (P<5E-7) or imputation quality measure

722 (MaCH-Rsq or IMPUTE-info score) <0.40 were excluded. All individuals with non-European

723 ancestry and siblings were removed.

### **GWAS Meta-analysis**

724

727

728

729

730

732

733

734

735

736

737

738

739

740

742

744

745

749

752

725 GWAS of the joint wheezing phenotypes were performed independently in ALSPAC, MAAS and the 726

combined IOW-SEATON-ASHFORD (combined as they were genotyped on the same platform, at

the same time, and quality-controlled and imputed together). All genetic data were imputed to a

new Haplotype Reference Consortium panel. This comprises around 31,000 sequenced individuals

(mostly European), so the coverage of European haplotypes is much greater than in other panels.

As a consequence, we expect to improve imputation accuracy, particularly at lower frequencies.

731 We used SNPTEST v2.5.2(26) with a frequentist additive multinomial regression model (-method

newml, never/infrequent wheeze as the reference) to investigate the association between SNPs

and wheezing phenotypes. No covariates were included in the model and only individuals of

European descent were included in this analysis. A meta-analysis of the three GWASs including

5,887 controls and 943 cases for early-onset persistent, 1482 cases for early-onset remitting, 603

cases for mid-childhood-onset remitting and 652 cases for late-onset wheeze, was performed

using METAL(27) with a total of 8,057,852 SNPs present. We used the option SCHEME STDERR in

METAL to implement an effect-size based method weighted by each study-specific standard error

in a fixed-effects model. We performed clumping to keep only one representative SNP per Linkage

disequilibrium (LD) block and used locus zoom plots to short-list independent SNPs for further

741 annotation.

#### LD clumping, pre-Selection and Gene Annotation

LD clumping was performed for all SNPs with p-value<10<sup>-5</sup> for at least one wheezing phenotype. In 743

order to avoid redundancy between SNPs and to ensure associations are independent, we used

significance thresholds of 0.05 for index and clumped SNPs (--clump-p1 0.05,--clump-p2 0.05), LD

746 threshold of 0.80 (--clump-r2 0.80) and physical distance threshold of 250kb (--clump-kb 250).

747 European 1000 Genome data were used to infer LD structure.

748 Locus Zoom plots (http://locuszoom.org/)(71) were used for close inspection of all independent

signals. Loci showing a peak with different colour dots (possibly indicating more than one causal

variant) were short-listed for further annotation. SNPnexus database (https://www.snp-750

751 nexus.org/v4/)(72) was used to annotate the overlapping, upstream and downstream genes; the

GWAS catalogue (by SNP and then gene) (https://www.ebi.ac.uk/gwas/search), GeneCards

753 (https://www.genecards.org/)(73) database and phenoscanner

- 754 (http://www.phenoscanner.medschl.cam.ac.uk/) were used to further explore previously
- 755 associated relevant phenotypes and gene function. Lead SNPs were looked in
- 756 https://www.regulomedb.org/ to assess potential functionality.

#### 757 **Genetic Control**

- 758 The genomic inflation factor ( $\lambda$ ) was calculated using the scipy stats chi2 module in Python. The
- 759 chi-squared test statistics from the meta-analysis p-values were first obtained. Then the observed
- 760 median chi-squared statistic from the calculated chi-squared test statistics were calculated. Finally,
- 761 the genomic inflation factor (λ) was derived by dividing the observed median chi-squared statistic
- by the expected median chi-squared statistic.

### 763 **Heterogeneity Scatter Plots**

- Heterogeneity scatter plots were based on filtering signals for each pair of wheezing phenotypes.
- 765 For example, for group1=persistent and group 2= early-onset mid-childhood remitting wheezing:
- 766 If group1 has a p-value<10<sup>-5</sup> and group2 has a p-value>0.05, and group1 has a negative effect size
- 767 (beta) while the lower bound of group2's effect size (beta CI) is greater than group1's effect size,
- 768 then we classified the result as group1 specific. If group1 has a p-value<10<sup>-5</sup> and group2 has a p-
- value>0.05, and group1 has a positive effect size (beta) while the upper bound of group2's effect
- size (beta + CI) is less than group1's effect size, then we classified the result as group1 specific.
- 771 If group2 has a p-value< 10<sup>-5</sup> and group1 has a p-value>0.05, and group2 has a negative effect size
- 772 (beta) while the lower bound of group1's effect size (beta CI) is greater than group2's effect size,
- then we classified the result as group2 specific. If group2 has a p-value<10<sup>-5</sup> and group1 has a p-
- value>0.05, and group2 has a positive effect size (beta) while the upper bound of group1's effect
- size (beta + CI) is less than group2's effect size, then we classified the result as group2 specific.
- 776 If both group1 and group2 have p-values< 10<sup>-5</sup>, and their effect sizes (betas) have the same sign
- 777 (i.e., both positive or both negative), then we classified the result as "Same direction".
- 778 If both group1 and group2 have p-values<10<sup>-5</sup>, and their effect sizes (betas) have opposite signs
- 779 (i.e., one positive and one negative), then we classified the result as "Opposite direction".

#### Gene expression in whole blood and lung tissues

- 781 The top independent SNPs associated with each of the wheeze phenotypes were assessed for their
- 782 association with cis- and trans-acting gene expression (mRNA) in whole blood and lung tissues. We
- 783 identified potential eQTL signals using Genotype-Tissue Expression database
- 784 (https://gtexportal.org) using the European reference panel.

**Appendix 2 Table 1.** List of 134 independent SNPs identified after clumping and associated with at least one wheezing phenotype ( $p<10x^{-5}$ )

CHR	SNP	ВР	short-listed after inspection of locus zoom plot
Persist	ent Wheezing		piot
1	rs4620530	240063821	yes
2	rs13398488	7142199	yes
2	rs77062323	53049017	no
2	rs6543291	106011626	yes
3	rs77655717	128737320	yes
4	rs77822621	1008212	yes
4	rs7680608	1050437	yes
4	rs115228498	142969757	yes
4	rs145937716	143192224	no
5	rs116494115	7736317	yes
5	rs78701483	95680422	no
6	rs138099941	7654240	no
6	rs9346404	71606613	no
6	rs143979498	151040328	no
7	rs76871421	105676144	yes
8	rs59670576	128555771	no
9	rs116933120	27458652	no
9	rs75260654	75788108	yes
9	rs116849664	75820902	yes
9	rs143481506	139515723	no
10	rs7088157	100038964	yes
11	rs112474574	5885773	yes
11	rs116861530	116962661	yes
13	rs7982350	73106322	no
13	rs17461573	106711373	no
14	rs1105683	56213787	yes
15	rs2202714	49811991	yes
15	rs117540214	84338642	yes
17	rs17676191	37949924	yes
17	rs79026872	37965932	yes
17	rs4795400	38067020	yes
17	rs1031460	38072247	yes
17	rs56199421	38090808	yes
17	rs4795406	38100134	yes
17	rs72832972	38110575	yes
17	rs4794821	38124203	yes
17	rs59843584	38124892	yes
18	rs111812993	30353181	no
19	rs4804311	8615589	yes
19	rs2013694	8616392	yes
			,


19	rs73501545	8620823	yes
19	rs111644945	8625081	yes
22	rs5994170	17615213	yes
22	rs34902370	17632194	yes
Early-o	nset Remitting W	heezing	
1	rs12730098	212427488	yes
1	rs75639566	233019116	no
2	rs2880066	17107219	yes
2	rs10180268	17126699	yes
3	rs115031796	86691640	no
3	rs3861377	173317378	yes
3	rs10513743	176022304	yes
5	rs10075253	75548246	yes
5	rs12520884	84406634	no
6	rs117477297	92565052	no
6	rs2453395	166286532	yes
7	rs56027869	50072919	no
7	rs4730561	78531705	yes
, 7	rs73144976	78586112	yes
7	rs67259321	78686582	-
7	rs146771277	154438861	yes no
9	rs10758259	34392908	
9 11			yes
	rs7128994	71242209	no
11	rs72994149	106837223	yes
12	rs117367256	93508478	no
13	rs2872948	57442480	yes
13	rs73527654	57447994	yes
13	rs2151504	82291577	no
15	rs116966886	47043587	yes
15	rs117565527	47342882	yes
	nildhood onset Re		ng
1	rs35725789	159207367	yes
1	rs146141555	159227423	yes
1	rs146575092	159374228	yes
1	rs140877050	220848829	no
1	rs72745905	223451086	no
2	rs7595553	36127878	yes
2	rs145007503	50688324	no
2	rs6546068	64583398	no
2	rs17387431	206651315	no
2	rs144791928	236963432	no
3	rs34315999	8969653	yes
3	rs115245770	99209128	no
3	rs146961758	194285978	yes
4	rs138794367	103859545	yes
5	rs115719402	77538102	yes
6	rs76026399	47531792	no
-		331,32	0


7	rs73172838	154842348	no
8	rs112631708	134500083	no
9	rs72752356	98094970	no
13	rs113195384	46333770	no
13	rs9602218	84139813	yes
13	rs61960366	84144202	yes
13	rs74589927	84208697	yes
13	rs2210726	84492936	yes
13	rs4390476	84598570	yes
14	rs117443464	73460284	yes
16	rs72820814	81916262	no
17	rs190526697	12274299	no
18	rs75286534	26206826	no
18	rs138888086	63591085	no
18	rs76551535	71879807	no
19	rs77496444	19192132	no
20	rs6077514	9302948	yes
Late-on	set Wheezing		
1	rs9439669	18859049	yes
1	rs2051039	57067560	yes
1	rs72673642	80727443	yes
2	rs147557117	19778063	no
2	rs140983998	111402871	yes
2	rs117617447	123387601	no
2	rs13025116	127505482	no
2	rs148008098	128633620	yes
3	rs4072729	24780393	yes
3	rs143960666	31227943	no
3	rs4677102	72193991	no
3	rs145629570	113422516	yes
3	rs113643470	141728174	yes
4	rs17472015	48467594	yes
7	rs117660982	149438923	yes
7	rs118027705	150456728	yes
7	rs139489493	150481499	yes
7	rs144271668	157934780	yes
8	rs990182	89976447	yes
9	rs79110962	14432953	yes
10	rs9325460	82492323	no
10	rs7896106	91196402	yes
10	rs115465993	109372900	no
11	rs16935643	41395746	no
11	rs141958628	119083284	yes
14	rs113363660	69410278	no
15	rs139134265	44923960	yes
15	rs143862030	84922146	yes
16	rs113390367	1118849	yes


16	rs4788025	28003221	yes	
18	rs72918264	51009510	no	
22	rs133498	48913809	yes	


# Appendix 2 Figure 1. Zoom locus plots for short-listed independent top hits for Persistent

# 789 Wheezing













# school Remitting Wheezing









# 814 childhood Remitting Wheezing








# Appendix 2 Figure 4. Zoom locus plots for short-listed independent top hits for Late-onset


# 823 Wheezing











#### **APPENDIX 3**

836

846

847

848

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863

864

865

866

### 837 **Post-GWAS: rs75260654 (ANXA1)**

- 838 Annotation & distribution
- 839 Information including chromosome, strand, clinical significance was retrieved from ENSEMBL using
- the R package biomaRt(74, 75). The effects of rs75260654 on genomic features were predicted by
- querying the using Ensembl Variant Effect Predictor (VEP) (76) web tool.
- rs75260654 distribution in the GRCh38.p13 build of the human genome across African, Asian and
- 843 European populations of the 1000 Genomes Project Phase 3 were accessed by guerying the
- 844 Ensembl (www.ensembl.org ) web browser on 24 May 2021.
- 845 Promoter Capture
  - The Hi-C libraries were prepared from CD4+ T-cells isolated from 7 healthy individuals (2 libraries per individual) from the MAAS cohort using the Arima-HiC kit (Arima Genomics). Promoter Capture Hi-C (PCHi-C) libraries were generated by capturing the restriction fragments (RF) overlapping the TSS of 18775 protein coding genes using the Agilent SureSelectXT HS Target Enrichment System according to the manufacturers' protocols. The final design included 305419 probes covering 13.476Mb and 18630 protein-coding genes. The restriction fragments (RF) overlapping the TSS (+/-1 RF, 3 RF per promoter) were captured with custom-designed biotinylated RNA baits. Libraries were sequenced to ~300M 2x150bp reads each (~600M reads/individual). The 3'-end of the reads was quality trimmed with Sickle. The sequencing data were processed with the HiCUP pipeline to map the sequencing reads and eliminate experimental artefacts and PCR duplicates(77). The BAM files from technical replicates were merged. Promoter interactions were called using the CHiCAGO pipeline(78), which calls statistically significant interactions in PCHi-C data while accounting for noise and PCHi-C specific bias. A CHiCAGO score > 5 (soft-thresholded -log weighted p-value) was considered significant. To gain information from all the available data, the BAM files from all 7 individuals were supplied as biological replicates in the analysis with CHiCAGO. Moreover, to increase power, restriction fragments were binned as follows: 10 consecutive RF that were not covered by the baits were binned together; the 3 baited fragments for each promoter were binned with 1 RF upstream and 1 downstream, totaling 5 fragments per promoter. If the bins for two consecutive promoters overlapped, these were binned together into a single larger bin. Publicly available ENCODE ATAC-seq (A549 cell line) and ChIP-seq (A549 cell line ENCFF900GVO) and POLR2A ChIP-seq (A549 cell line, ENCFF737ZKN) data and 18-

867 state ChromHMM from the EpiMap Project (BSS00007) (79) for A549 cell line were downloaded. The PCHi-C interactions of interest and their overlap with ATAC-seq and ChIP-seq peaks, and 868 869 putative enhancers from the 18-state ChromHMM model were visualised using the Sushi R 870 package. eQTL catalogue lookup 871 872 We queried the eQTL catalogue (https://www.ebi.ac.uk/eqtl/; accessed 6 May 2021) using tabix-873 0.2.6 to assess if rs75260654, rs116849664 or rs78320984 are eQTLs in studies that utilised the 874 following cell types: lung, T cells, blood, monocytes, neutrophils, NK cells, fibroblasts, B cells, CD4+ 875 T cells, CD8+ T cells, Th17 cells, Th1 cells, Th2 cells, Treg naive, Treg memory, CD16+ monocytes, 876 Cultured fibroblasts, EBV-transformed lymphocytes. We defined nominal significance as p<=0.05. 877 Variant effect 878 Variant effect on tissue-specific gene expression, which is based on GTEx eQTL, was retrieved on 879 May 24 from eQTL Ensembl database (https://www.ensembl.org/) and eQTLGene Consortium 880 (https://www.eqtlgen.org/cis-eqtls.html).Using downloaded correlation of variant on tissue 881 specific gene expression from Ensembl, the relative effect of T allele on the ANXA1 expression 882 across 86 tissue types was presented in scatter plot using R version 3.6.1.(80) To get information 883 on the functional role of ANXA1, the top 30 interacting proteins and enrichment were retrieved

from STRING database(81) into cystoscape for visualization.(82)

#### APPENDIX 4

885

886

#### Results in context of literature

- Previously relevant associated traits for each region/gene now associated with different wheezing
- phenotypes are presented in **Appendix 5-Tables 1-4.**
- 889 Persistent Wheeze
- 890 We identified two GWAS-significant loci: 17q21, p<5.5×10<sup>-9</sup>, and a novel locus on 9q21.13
- 891 (ANXA1), p<6.7×10<sup>-9</sup>. The remaining 17 loci  $(4.0\times10^{-7}\le p\text{-values}\le 9.8\times10^{-6})$  included regions
- 892 previously associated with childhood asthma (1q43, 4p16.3, 4q31.21, 5p15.31, 7q22.3, 17q12),
- 893 asthma and rhinitis (2p25.1), eosinophil count (3q21.3, 10q24.2, 11q23.3, 22q11.1), bronchial
- 894 hyper-responsiveness (2q12.2), lung function (1q43, 3q21.3, 5q13.3, 15q25.2, 19p13.2),
- 895 triglycerides measurement and/or glucose metabolism (11q23.3, 19p13.2 and 22q11.1), severe
- asthma (14g22.1) and severe asthma and insulin resistance (11p15.4). See Appendix 5-Table 1.
- 897 Early-onset Preschool Remitting Wheeze
- 898 Among the regions associated with early-onset preschool remitting wheeze, we identified loci
- 899 previously associated with smoking (3q26.31, 7q21.11 and 15q21.1), waist circumference and
- 900 obesity (1q32.3), asthma and/or BMI (5q13.3, 6q27, 7q21.11), allergic disease and atopy (7q21.11)
- and airway repair (2p24.2 and 9p13.3). See **Appendix 5-Table 2.**
- 902 Early-onset Mid-childhood Remitting Wheeze
- 903 Loci associated with this phenotype were previously associated with neutrophil counts (1q23.2,
- 904 3q29, 20p12.3-p12.2), eosinophil counts and allergic rhinitis (4q24), pollution and DNA
- 905 methylation (2p22.3), atopy (3p25.3), food allergy (13q31.1) and BMI (3q29, 5q14.1). See
- 906 Appendix 5-Table 3.
- 907 Late-onset Wheeze
- 908 Regions associated with late-onset wheeze were previously associated with adult-onset non-
- allergic asthma (3q13.2), asthma/allergic disease and allergy/atopic sensitization (3q23, 7q36.1),
- 910 asthma and/or allergy in adolescence (9p22.3, 16p12.1), late-onset asthma and obesity (2q13),
- 911 lung function or body height (2q14.3, 3p24.2, 3q13.2, 15q25.2), lymphocyte count and asthma
- 912 susceptibility (1p32.2), obesity and/or metabolic syndrome/dysfunction (1p36.13 and 22q13.32),

eczema (7q36.3), insulin resistance (10q23.31), type 1 diabetes (8q21.3), alcohol drinking (15q15.3-q21.1) and sex hormone-binding globulin levels (11q23.3). See **Appendix 5-Table 4.** 

Appendix 4 Table 1. References to previous relevant associated traits for each region/gene identified in Early Onset Persistent Wheezing

Early Onset Persistent Wh	neezing		
Gene(s)	Locus	Previous Associated Trait	Reference or Source
CHRM3	1q43	FEV <sub>1</sub> , FEV <sub>1</sub> /FVC, asthma- high priority drug	Patel, K.R. et al. Targeting acetylcholine receptor M3
		target	prevents the progression of airway hyperreactivity in a
			mouse model of childhood asthma. FASEB J 31, 4335-
			4346 (2017).
RNF144A	2p25.1	asthma, allergy, childhood onset asthma,	Schoettler, N. et al. Advances in asthma and allergic
		allergic rhinitis	disease genetics: Is bigger always better? J Allergy Clin
			Immunol <b>144</b> , 1495-1506 (2019).
FHL2	2q12.2	bronchial hyper-responsiveness, airway	Kurakula, K. et al. Deficiency of FHL2 attenuates airway
		inflammation, novel gene associated with	inflammation in mice and genetic variation associates
		asthma severity in human	with human bronchial hyper-responsiveness. Allergy 70,
			1531-44 (2015).
RAB7A	3q21.3	eosinophil count	GeneCards
RAB43	3q21.3	response to bronchodilator, FEV <sub>1</sub> /FEC ratio	GWAS Catalog
RNF212, IDUA, DGKQ,	4p16.3	asthma	Gautam, Y. et al. Comprehensive functional annotation
SLC26A1			of susceptibility variants associated with asthma. Hum
			Genet <b>139</b> , 1037-1053 (2020).
INPP4B	4q31.21	atopic asthma	Sharma, M. et al. A genetic variation in inositol
			polyphosphate 4 phosphatase a enhances susceptibility

			to asthma. Am J Respir Crit Care Med 177, 712-9 (2008).
ADCY2	5p15.31	asthma × air pollution, childhood asthma	Gref, A. et al. Genome-Wide Interaction Analysis of Air
			Pollution Exposure and Childhood Asthma with
			Functional Follow-up. Am J Respir Crit Care Med 195,
			1373-1383 (2017).
CDHR3	7q22.3	childhood <b>asthma</b>	Everman, J.L. et al. Functional genomics of CDHR3
			confirms its role in HRV-C infection and childhood
			asthma exacerbations. J Allergy Clin Immunol 144, 962-
			971 (2019).
ANXA1	9q21.13	FEV <sub>1</sub> /FVC, response to bronchodilators in	Lutz, S.M. et al. A genome-wide association study
		smokers	identifies risk loci for spirometric measures among
			smokers of European and African ancestry. BMC Genet
			<b>16</b> , 138 (2015).
ANXA1	9q21.13	anti-inflammatory properties, strongly	Vieira Braga FA et al. A cellular census of human lungs
		expressed in bronchial mast cells	identifies novel cell states in health and in asthma.
			(2019).
ANXA1	9q21.13	potentially involved in epithelial airway	Leoni, G. et al. Annexin A1-containing extracellular
		repair	vesicles and polymeric nanoparticles promote epithelial
			wound repair. <i>J Clin Invest</i> <b>125</b> , 1215-27 (2015).
R3HCC1L	10q24.2	atopic <b>eczema</b> , psoriasis	GWAS Catalog
R3HCC1L	10q24.2	eosinophil count, BMI	GeneCards

TRIM5, TRIM6, TRIM22	11p15.4	severe asthma and insulin resistance	Kimura, T. et al. Precision autophagy directed by
			receptor regulators - emerging examples within the
			TRIM family. <i>J Cell Sci</i> <b>129</b> , 881-91 (2016).
SIK3	11q23.3	triglycerides, glucose metabolism, eosinophil	Sun, Z. et al. The potent roles of salt-inducible kinases
		count	(SIKs) in metabolic homeostasis and tumorigenesis. Sig
			Transduct Target Ther <b>5</b> (2020).
KTN1	14q22.1	severe asthma	Bigler, J. et al. A Severe Asthma Disease Signature from
			Gene Expression Profiling of Peripheral Blood from U-
			BIOPRED Cohorts. Am J Respir Crit Care Med 195, 1311-
			1320 (2017).
FAM227B	5q13.3	rs35251997 and FEV <sub>1</sub> , FEV <sub>1</sub> /FVC	Shrine, N. et al. New genetic signals for lung function
			highlight pathways and chronic obstructive pulmonary
			disease associations across multiple ancestries. Nat
			Genet <b>51</b> , 481-493 (2019).
ADAMTSL3	15q25.2	FEV1/FVC	Sakornsakolpat, P. et al. Genetic landscape of chronic
			obstructive pulmonary disease identifies heterogeneous
			cell-type and phenotype associations. Nat Genet 51,
			494-505 (2019).
IKZF3, GSDMB, LRRC3C,	17q12	early-onset asthma, persistent wheezing	Granell R et al. Examination of the relationship between
GSDMA		(chr17q12-q21)	variation at 17q21 and childhood wheeze phenotypes. J
			Allergy Clin Immunol. 2013 Mar;131(3):685-94.

MARCH2,	HNRNPM,	19p13.2	triglycerides,	HDL-chole:	sterol,	metabolic	Sajuthi, S.P. et al. Genetic regulation of adipose tissue
MYO1F			syndrome				transcript expression is involved in modulating serum
							triglyceride and HDL-cholesterol. <i>Gene</i> <b>632</b> , 50-58
							(2017).
MYO1F		19p13.2	FEV <sub>1</sub> and FVC				GeneCards
CECDE.		22 44 4					
CECR5		22q11.1	triglycerides,	eosinophil	count	and body	Liu, D.J. et al. Exome-wide association study of plasma
			height				lipids in >300,000 individuals. <i>Nat Genet</i> <b>49</b> , 1758-1766
							(2017).

Appendix 4 Table 2. References to previous relevant associated traits for each region/gene identified in Early-onset Pre-school Remitting

## 918 Wheezing

Early-onset Pre-schoo	l Remitting Wi		
Gene(s)	Locus	Previous Associated Trait	Reference or source
PPP2R5A	1q32.3	waist circumference & obesity	Kim, H.J. et al. Combined linkage and association analyses identify a novel locus for obesity near PROX1 in Asians. Obesity (Silver Spring) 21, 2405-12 (2013).
FAM49A or CYRIA	2p24.2	airway repair in non-atopic asthma	Hoang, T.T. <i>et al.</i> Epigenome-wide association study of DNA methylation and adult asthma in the Agricultural Lung Health Study. <i>Eur Respir J</i>

			<b>56</b> (2020).
NLGN1	3q26.31	smoking	Drgon, T. et al. Genome-wide association for
			nicotine dependence and smoking cessation
			success in NIH research volunteers. Mol Med 15,
			21-7 (2009).
NAALADL2	3q26.31	suggestive association with severe asthma	Herrera-Luis E et al. Genome-wide association
		exacerbations	study reveals a novel locus for asthma with severe
			exacerbations in diverse populations. Pediatr
			Allergy Immunol. 2021;32(1):106-115.
SV2C	5q13.3	BMI, diastolic blood pressure	GeneCards
PDE10A	6q27	Birthweight, asthma and BMI	Melen, E. et al. Analyses of shared genetic factors
			between asthma and obesity in children. J Allergy
			Clin Immunol <b>126</b> , 631-7 e1-8 (2010).
MAGI2	7q21.11	allergic diseases & atopy	Freidin, M.B. et al. [Genome-wide association
			study of allergic diseases in Russians of Western
			Siberia]. <i>Mol Biol (Mosk)</i> <b>45</b> , 464-72 (2011).
MAGI2	7q21.11	smoking	Quach, B.C. et al. Expanding the genetic
			architecture of nicotine dependence and its
			shared genetics with multiple traits. Nat Commun
			<b>11</b> , 5562 (2020).
MAGI2	7q21.11	BMI	GeneCards

MAGI2	7q21.11	airway wall thickness	GWAS Catalog
C9orf24	9p13.3	airway repair	Yoshisue, H. et al. Characterization of ciliated
			bronchial epithelium 1, a ciliated cell-associated
			gene induced during mucociliary differentiation.
			Am J Respir Cell Mol Biol <b>31</b> , 491-500 (2004).
GUCY1A2	11q22.3	systolic/diastolic blood pressure	GeneCards
PRR20A/B/C/D/E	13q21.1	systolic <b>blood pressure</b>	GeneCards
SEMA6D	15q21.1	smoking	Minica, C.C. et al. Pathways to smoking
			behaviours: biological insights from the Tobacco
			and Genetics Consortium meta-analysis. Mol
			Psychiatry 22, 82-88 (2017).

920

Appendix 4 Table 3. References to previous relevant associated traits for each region/gene identified in Early-onset Mid-childhood Remitting

## 922 Wheezing

Early-onset Mid-	-childhood Remitt	ting Wheezing	
Gene(s)	Locus	Previous Associated Trait	Reference
CADM3, FCER1A, MPTX1, OR10J1	1q23.2	neutrophil count, CRP	Barreto, M. <i>et al.</i> Duffy phenotyping and FY*B-67T/C genotyping as screening test for benign constitutional neutropenia. <i>Hematol Transfus Cell Ther</i> (2020).
MRPL50P1	2р22.3	PM 2.5 exposure level and global DNA methylation level	Liu, J. <i>et al.</i> Genetic variants, PM2.5 exposure level and global DNA methylation level: A multi-center population-based study in Chinese. <i>Toxicol Lett</i> <b>269</b> , 77-82 (2017).
RAD18	3p25.3	atopy/SPT	Bouzigon, E. et al. Meta-analysis of 20 genome-wide linkage studies evidenced new regions linked to asthma and atopy. Eur J Hum Genet 18, 700-6 (2010).
MRPL50P1	3q29	ВМІ	Kettunen, J. et al. Multicenter dizygotic twin cohort study confirms two linkage susceptibility loci for body mass index at 3q29 and 7q36 and identifies three further potential novel loci. Int J Obes (Lond)

			<b>33</b> , 1235-42 (2009).
LSG1	3q29	BMI, eosinophil & neutrophil count	GeneCards
TMEM44-AS1,	3q29	diastolic blood pressure	GeneCards
TMEM44,			
ATP13A3			
SLC9B1	4q24	eosinophil count	Aschard, H. et al. Sex-specific effect of IL9
			polymorphisms on lung function and
			polysensitization. Genes Immun 10, 559-65 (2009).
SLC9B1	4q24	allergic rhinitis	Haagerup, A. et al. Allergic rhinitisa total genome-
			scan for susceptibility genes suggests a locus on
			chromosome 4q24-q27. Eur J Hum Genet <b>9</b> , 945-52
			(2001).
AP3B1	5q14.1	vital capacity, BMI	GeneCards, GWAS Catalog
RNU6-67P,	13q31.1	RNU6-67P/ rs976078: food allergy	Liu, X. et al. Genome-wide association study of
SLITRK1,			maternal genetic effects and parent-of-origin effects
VENTXP2,			on food allergy. Medicine (Baltimore) 97, e0043
UBE2D3P4,			(2018).
MTND4P1			
ZFYVE1	14q24.2	LDL cholesterol and systolic blood pressure	GWAS Catalog

PLCB4	20p12.3-p12.2	neutrophil count	Okada, Y. et al. Common variations in PSMD3-CSF3
			and PLCB4 are associated with neutrophil count.
			Hum Mol Genet <b>19</b> , 2079-85 (2010).

Late-onset Whe	ezing		
Gene(s)	Locus	Previous Associated Trait	Reference
KLHDC7A	1p36.13	1p36.13: metabolic syndrome	Hoffmann, K. et al. A German genome-wide
			linkage scan for type 2 diabetes supports the
			existence of a metabolic syndrome locus on
			chromosome 1p36.13 and a type 2 diabetes locus
			on chromosome 16p12.2. Diabetologia <b>50</b> , 1418-
			22 (2007).
PPAP2B,	1p32.2	PRKAA2: lymphocyte count and asthma	Cusanovich, D.A. et al. The combination of a
PRKAA2		susceptibility	genome-wide association study of lymphocyte
			count and analysis of gene expression data reveals
			novel asthma candidate genes. Hum Mol Genet
			<b>21</b> , 2111-23 (2012).
HMGB1P18	1p31.1	smoking, BMI	GeneCards
ACOXL, BUB1	2q13	ACOXL: later onset asthma and obesity	Zhu, Z. et al. Shared genetic and experimental
			links between obesity-related traits and asthma
			subtypes in UK Biobank. J Allergy Clin Immunol
			<b>145</b> , 537-549 (2020).

AMMECR1L	2q14.3	body height, blood protein, growth, bone, and	Moyses-Oliveira, M. et al. Inactivation of
		heart alterations	AMMECR1 is associated with growth, bone, and
			heart alterations. <i>Hum Mutat</i> <b>39</b> , 281-291 (2018).
RARB	3p24.2	FEV <sub>1</sub> /FVC, adult lung function	Collins, S.A. et al. HHIP, HDAC4, NCR3 and RARB
			polymorphisms affect fetal, childhood and adult
			lung function. Eur Respir J 41, 756-7 (2013).
KIAA2018,	3q13.2	SIDT1: FEV <sub>1</sub> /FVC, CD200: adult-onset non-allergic	Siroux, V. et al. Genetic heterogeneity of asthma
NAA50, SIDT1,		asthma	phenotypes identified by a clustering approach.
CD200			Eur Respir J <b>43</b> , 439-52 (2014).
TFDP2, XRN1	3q23	XRN1: eosinophil count, 3q23: allergic disease and	Freidin, M.B. et al. [Genome-wide association
		atopic sensitisation	study of allergic diseases in Russians of Western
			Siberia]. <i>Mol Biol (Mosk)</i> <b>45</b> , 464-72 (2011).
SLAIN2,	4p11	FRYL: body height, age at menopause	GeneCards
SLC10A4, FRYL			
KRBA1, ZNF467	7q36.1	systolic blood pressure	GWAS Catalog
GIMAP family,	7q36.1	AOC1: CV disease, smoking, GIMAP family:	Heinonen, M.T. et al. GIMAP GTPase family genes:
AOC1,		autoimmune diabetes, asthma and allergy	potential modifiers in autoimmune diabetes,
LOC105375566			asthma, and allergy. J Immunol 194, 5885-94
			(2015).
PTPRN2	7q36.3	eczema	Bogari, N.M. et al. Whole exome sequencing
			detects novel variants in Saudi children diagnosed

			with eczema. <i>J Infect Public Health</i> <b>13</b> , 27-33 (2020).
LOC105375631	8q21.3	8q21.3: type 1 diabetes	Mukhopadhyay, N., Noble, J.A., Govil, M.,
			Marazita, M.L. & Greenberg, D.A. Identifying
			genetic risk loci for diabetic complications and
			showing evidence for heterogeneity of type 1
			diabetes based on complications risk. PLoS One
			<b>13</b> , e0192696 (2018).
NFIB, ZDHHC21	9p22.3	9p22.3: asthma (mean age<16 years)	Denham, S. et al. Meta-analysis of genome-wide
			linkage studies of asthma and related traits. Respir
			Res <b>9</b> , 38 (2008).
SLC16A12, IFIT	10q23.31	SLC16A12: Body height, PANK1: insulin resistance	Yang, L. et al. P53/PANK1/miR-107 signalling
family, PANK1			pathway spans the gap between metabolic
			reprogramming and insulin resistance induced by
			high-fat diet. <i>J Cell Mol Med</i> <b>24</b> , 3611-3624 (2020).
CBL, CCDC84,	11q23.3	CBL: Sex hormone-binding globulin levels; MCAM:	GWAS Catalog
MCAM		Blood protein levels	
SPG11, CTDSPL2	15q15.3-q21.1	CTDSPL2: alcohol drinking	GWAS Catalog
ADAMTSL3,	15q25.2	ADAMTSL3: FEV <sub>1</sub> /FVC, lean mass	Karasik, D. et al. Disentangling the genetics of lean
GOLGA6L4,			mass. Am J Clin Nutr <b>109</b> , 276-287 (2019).
UBE2Q2P8			

SSTR5-AS1,	16p13.3	CACNA1H: eosinophil count	GWAS Catalog
CACNA1H			
GSG1L	16p12.1	16p12.1: current asthma and rhino-conjunctivitis at	Sottile, G. et al. An association analysis to identify
		10-15 years	genetic variants linked to asthma and rhino-
			conjunctivitis in a cohort of Sicilian children. <i>Ital J</i>
			Pediatr <b>45</b> , 16 (2019).
FAM19A5 or	22q13.32	Obesity and Metabolic Dysfunction	Recinella L. et al. Adipokines: New Potential
TAFA5			Therapeutic Target for Obesity and Metabolic,
			Rheumatic, and Cardiovascular Diseases. Front
			Physiol. 2020 Oct 30;11:578966

927 APPENDIX 5
 928 Appendix 5 Table 1. SNPs near ANXA1 associated with persistent wheeze

Chr	Rsid	position		A 2	freqA	beta	se	P value	Direction (3 GWAS)
9	rs75260654	75788108	t	c	0.02	0.90	0.16	6.66e-09	
9	rs116849664	75820902	t	c	0.02	0.89	0.16	1.99e-08	
9	rs78320984	75844302	t	g	0.02	0.81	0.15	6.41e-08	

929 A1 is the effect allele, A2 is the reference allele.

## 930 **Appendix 5 Table 2.** Allele Frequencies of *rs75260654* across different wheeze phenotypes

Phenotype	CC	CT	TT
Never/infrequent	5641 (97.2)	161 (2.8)	0 (0)
Early-onset pre-school remitting	1409 (97.1)	42 (2.9)	0 (0)
Early-onset mid-childhood remitting	572 (96.1)	23 (3.9)	0 (0)
Late-onset	613 (95.2)	31 (4.8)	0 (0)
Early-onset persistent	867 (94.2)	51 (5.5)	2 (0.2)

## 931 **Appendix 5 Table 3.** Selected immune eQTLs of rs75260654

Rsid	P value	beta	se	an	symbol	Study
rs75260654	0.014	-0.65	0.26	382	ANXA1	Quach_2016_monocyte_R848
rs75260654	0.015	-1.02	0.41	396	ANXA1	Quach_2016_monocyte_IAV

## 932 Appendix 5 Table 4. Lung eQTLs of rs75260654

Rsid	P value	beta	se	an	symbol	Study
rs116849664	0.0489	0.22	0.11	620	ANXA1	GTEx exon lung

rs78320984 0.0489 0.22 0.11 620 *ANXA1* GTEx\_exon\_lung

**Appendix 5 Table 5.** Functional enrichment for *ANXA1*: Top 10 GO terms

Term name	Description	FDR value
GO.0007186	G protein-coupled receptor signaling pathway	3.57×10 <sup>-18</sup>
GO.0006954	inflammatory response	1.13×10 <sup>-16</sup>
GO.0006874	cellular calcium ion homeostasis	5.55×10 <sup>-15</sup>
GO.0007204	positive regulation of cytosolic calcium ion concentration	1.65×10 <sup>-14</sup>
GO.0060326	cell chemotaxis	1.95×10 <sup>-14</sup>
GO.0006955	immune response	4.23×10 <sup>-14</sup>
GO.0006935	Chemotaxis	4.93×10 <sup>-14</sup>
GO.0006952	defense response	1.68×10 <sup>-13</sup>
GO.0050801	ion homeostasis	2.23×10 <sup>-13</sup>
GO.0002376	immune system process	2.87×10 <sup>-13</sup>

#### Replication of ANXA1 top hits in PIAMA cohort

### PIAMA cohort description

PIAMA (Prevention and Incidence of Asthma and Mite Allergy) is an ongoing birth cohort study. Details of the study design have been published previously(83, 84). In brief, pregnant women were recruited from the general population through antenatal clinics in the north, west and centre of the Netherlands in 1996-1997. The baseline study population consisted of 3963 new-borns. Questionnaires were completed by the parents during pregnancy when the child was 3 months old, and then annually from 1 up to 8 years; at ages 11, 14 and 17 years, questionnaires were completed by the parents as well as the participants themselves.

#### LCA wheezing phenotypes

A 6 class LCA model was identified including 3,832 individuals with at least 2 observations of wheeze between 1 and 11-12 years of age. The identified classes were labelled: never/Infrequent (2909,75.91%), pre-school onset remitting (571, 14.90%), mid-childhood school remitting (108, 2.82%), intermediate onset remitting (106, 2.77%), school-age onset persisting (74, 1.93%) and continuous wheeze (64, 1.67%).

#### Replication analyses

We analyzed associations between SNPs downstream of *ANXA1* (Appendix 5-Table 1, Appendix 5-Figure 1) and continuous wheezing in PIAMA, using the never/infrequent wheezing as the baseline category. Analyses were carried out in SPSS using a logistic regression model.


# **Appendix 5 Table 6** Replication of associations between SNPs downstream of *ANXA1* and early-onset persistent wheezing in PIAMA

				CW (40) vs NI (1557)		
Rsid	chr:position	R2	A2/freqA2	beta	se	p-value
rs75260654	9:75788108	0.60	c/0.02	-0.287	0.91	0.75
rs116849664	9:75820902	0.61	c/0.02	0.119	1.08	0.91


rs78320984	9:75844302	0.59	g/0.02	0.125	1.04	0.90

A2 is the reference allele. CW=continuous wheezing, IR=intermediate wheezing derived from LCA 1 to 12 years in PIAMA.

**Appendix 5 Figure 1.** Linkage disequilibrium between SNPs downstream of *ANXA1* that were associated with persistent wheeze.



# 965 *IL-13* and *NR3C1*



#### **APPENDIX 6**

#### **Functional mouse experiments**

969 *Mice* 

967

- 970 In accordance with the Animals (scientific procedures) act 1986, all animal experiments were 971 conducted under the approved UK Home Office Project License No: PPL 70/7643, reviewed by 972 Imperial College's Animal Welfare and Ethical Review body. Female WT BALB/c and annexin 973 A1 knockout (KO) mice were purchased from Charles River (Bicester, UK). Animals aged 6-8 974 weeks of age received 25ug intranasal instillation of either HDM (Greer Laboratories, Lenoir, 975 NC, USA; Cat: XPB70D3A25), or PBS 3x a week for 3 weeks. Mice were sacrificed 4 hours post-976 final HDM challenge. Mice were housed under specific pathogen-free conditions and a 12:12 977 light:dark cycle. Food and water were supplied ad libitum. All animal experiments were 978 completed twice, with N=4-6 per group.
- 979 Airway hyperresponsiveness
- Airway hyperreactivity was measured using Flexivent™. Lung resistance was measured in response to increasing doses of methacholine (3-100mg/ml, Sigma, Poole, UK, Cat: A2251) in
- 982 tracheotomised anaesthetised mice using an EMMS system (Electro-Medical Measurement
- 983 Systems, UK).
- 984 Flow Cytometry Analysis
- 985 Bronchoalveolar lavage (BAL) was collected. BAL cells were restimulated with ionomycin and
- 986 phorbol 12-myristate 13-acetate in the presence of brefeldin (Sigma), as previously
- 987 described(85). Specific antibodies for T1/ST2 staining were purchased from Morwell
- 988 Diagnostics (Zurich, Switzerland). Cells were also stained for lineage negative cocktail, Ly6G,
- 989 CD45, CD11b, CD11c, SiglecF. Labelled cells were acquired on a BD Fortessa (BD Biosciences,
- 990 Oxford, UK) and analysed using FlowJo software (Treestar, Ashland, Oregan, USA). Details of
- antibodies used can be found in the **Appendix 6-table 1**.

992 Analysis of cytokines and chemokines

993 Murine lung tissue homogenate supernatants were processed as previously described(85).

Cytokine levels were analysed by ELISA: IL-4, IL-5 (PharMingen, Oxford, UK), IL-13 Ready-Set-

995 Go kits (eBioscience).

Real time-PCR

994

996

997

998

999

1000

1001

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

Total RNA was extracted from murine lung tissue using an RNeasy Mini Kit (Qiagen). Total RNA (1µg) was reverse transcribed into cDNA using a High-Capacity cDNA Reverse Transcription Kit (Life Technologies, UK). Real-time PCR reactions were performed using TagMan Gene Expression Master Mix and TagMan Gene Expression probes, annexin A1, and

HPRT (Applied Biosystems). Values were normalised to HPRT and gene expression was

analysed using the change-in-threshold 2-ΔCT method.

#### Annexin A1 Immunohistochemistry

Paraffin-embedded mouse lung sections were stained with annexin A1 (R&D Systems, MAB3770). Annexin A1 primary antibody was followed by a secondary detection antibody (donkey anti-goat 488, Thermofisher, A11055). Annexin A1+ cells were quantified by manual counting under microscope and numbers averaged over four fields, from five biological replicates per group.

### Statistical analysis

Data are expressed as median ± IQR. Statistical differences between groups were calculated using Mann Whitney U test, unless otherwise specified. p -values are indicated in figures.

## **Appendix 6 Table 1** Antibodies used in the Flow Cytometry Analysis

			Conjugated	
Molecule	Manufacturer	Isotype	dye	Clone
T1 /ST2	Morwell Diagnostics GMBH,			
T1/ST2	Switzerland	Rat IgG1	FITC	DJ8
CD45	e-Bioscience Ltd, Hatfield, UK	Rat IgG2b	PerCP-CY5.5	30-F11
CD11b	BD Biosciences, Oxford, UK	Rat IgG2b	e450	M1/70

CD11c	e-Bioscience Ltd, Hatfield, UK	Hamster IgG1	PerCP-CY5.5	N418
Siglec F	BD Biosciences, Oxford, UK	Rat IgG2a	PE	E50-2440

#### 1014 **REFERENCES**

1021

1022

1023

10241025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

- 1. Pavord ID, Beasley R, Agusti A, Anderson GP, Bel E, Brusselle G, Cullinan P, Custovic A, Ducharme
   FM, Fahy JV, Frey U, Gibson P, Heaney LG, Holt PG, Humbert M, Lloyd CM, Marks G, Martinez
   FD, Sly PD, von Mutius E, Wenzel S, Zar HJ, Bush A. After asthma: redefining airways diseases.
   Lancet 2018; 391: 350-400.
- 2. Duffy DL, Martin NG, Battistutta D, Hopper JL, Mathews JD. Genetics of asthma and hay fever in
   Australian twins. *Am Rev Respir Dis* 1990; 142: 1351-1358.
  - 3. Simpson A, Custovic A, Tepper R, Graves P, Stern DA, Jones M, Hankinson J, Curtin JA, Wu J, Blekic M, Bukvic BK, Aberle N, Marinho S, Belgrave D, Morgan WJ, Martinez FD. Genetic variation in vascular endothelial growth factor-a and lung function. *Am J Respir Crit Care Med* 2012; 185: 1197-1204
  - 4. Daniels SE, Bhattacharrya S, James A, Leaves NI, Young A, Hill MR, Faux JA, Ryan GF, le Souef PN, Lathrop GM, Musk AW, Cookson WO. A genome-wide search for quantitative trait loci underlying asthma. *Nature* 1996; 383: 247-250.
  - 5. Moffatt MF, Kabesch M, Liang L, Dixon AL, Strachan D, Heath S, Depner M, von Berg A, Bufe A, Rietschel E, Heinzmann A, Simma B, Frischer T, Willis-Owen SA, Wong KC, Illig T, Vogelberg C, Weiland SK, von Mutius E, Abecasis GR, Farrall M, Gut IG, Lathrop GM, Cookson WO. Genetic variants regulating ORMDL3 expression contribute to the risk of childhood asthma. *Nature* 2007; 448: 470-473.
  - 6. Moffatt MF, Gut IG, Demenais F, Strachan DP, Bouzigon E, Heath S, von Mutius E, Farrall M, Lathrop M, Cookson W, Consortium G. A large-scale, consortium-based genomewide association study of asthma. *N Engl J Med* 2010; 363: 1211-1221.
- 1036 7. Demenais F, Margaritte-Jeannin P, Barnes KC, Cookson WOC, Altmuller J, Ang W, Barr RG, Beaty TH, 1037 Becker AB, Beilby J, Bisgaard H, Bjornsdottir US, Bleecker E, Bonnelykke K, Boomsma DI, 1038 Bouzigon E, Brightling CE, Brossard M, Brusselle GG, Burchard E, Burkart KM, Bush A, Chan-1039 Yeung M, Chung KF, Couto Alves A, Curtin JA, Custovic A, Daley D, de Jongste JC, Del-Rio-1040 Navarro BE, Donohue KM, Duijts L, Eng C, Eriksson JG, Farrall M, Fedorova Y, Feenstra B, 1041 Ferreira MA, Australian Asthma Genetics Consortium c, Freidin MB, Gajdos Z, Gauderman J, 1042 Gehring U, Geller F, Genuneit J, Gharib SA, Gilliland F, Granell R, Graves PE, Gudbjartsson DF, 1043 Haahtela T, Heckbert SR, Heederik D, Heinrich J, Heliovaara M, Henderson J, Himes BE, Hirose 1044 H, Hirschhorn JN, Hofman A, Holt P, Hottenga J, Hudson TJ, Hui J, Imboden M, Ivanov V, 1045 Jaddoe VWV, James A, Janson C, Jarvelin MR, Jarvis D, Jones G, Jonsdottir I, Jousilahti P, 1046 Kabesch M, Kahonen M, Kantor DB, Karunas AS, Khusnutdinova E, Koppelman GH, Kozyrskyj 1047 AL, Kreiner E, Kubo M, Kumar R, Kumar A, Kuokkanen M, Lahousse L, Laitinen T, Laprise C, 1048 Lathrop M, Lau S, Lee YA, Lehtimaki T, Letort S, Levin AM, Li G, Liang L, Loehr LR, London SJ, 1049 Loth DW, Manichaikul A, Marenholz I, Martinez FJ, Matheson MC, Mathias RA, Matsumoto K, 1050 Mbarek H, McArdle WL, Melbye M, Melen E, Meyers D, Michel S, Mohamdi H, Musk AW, 1051 Myers RA, Nieuwenhuis MAE, Noguchi E, O'Connor GT, Ogorodova LM, Palmer CD, Palotie A, 1052 Park JE, Pennell CE, Pershagen G, Polonikov A, Postma DS, Probst-Hensch N, Puzyrev VP, Raby 1053 BA, Raitakari OT, Ramasamy A, Rich SS, Robertson CF, Romieu I, Salam MT, Salomaa V, 1054 Schlunssen V, Scott R, Selivanova PA, Sigsgaard T, Simpson A, Siroux V, Smith LJ, Solodilova M, 1055 Standl M, Stefansson K, Strachan DP, Stricker BH, Takahashi A, Thompson PJ, Thorleifsson G, 1056 Thorsteinsdottir U, Tiesler CMT, Torgerson DG, Tsunoda T, Uitterlinden AG, van der Valk RJP, 1057 Vaysse A, Vedantam S, von Berg A, von Mutius E, Vonk JM, Waage J, Wareham NJ, Weiss ST, 1058 White WB, Wickman M, Widen E, Willemsen G, Williams LK, Wouters IM, Yang JJ, Zhao JH, 1059 Moffatt MF, Ober C, Nicolae DL. Multiancestry association study identifies new asthma risk 1060 loci that colocalize with immune-cell enhancer marks. Nat Genet 2018; 50: 42-53. 1061
  - 8. El-Husseini ZW, Gosens R, Dekker F, Koppelman GH. The genetics of asthma and the promise of genomics-guided drug target discovery. *Lancet Respir Med* 2020; 8: 1045-1056.
  - 9. Kim KW, Ober C. Lessons Learned From GWAS of Asthma. Allergy Asthma Immun 2019; 11: 170-187.

1062

- 10. Ober C, Yao TC. The genetics of asthma and allergic disease: a 21st century perspective. *Immunol Rev* 2011; 242: 10-30.
- 1066 11. Custovic A, Marinho S, Simpson A. Gene-environment interactions in the development of asthma and atopy. *Expert Rev Respir Med* 2012; 6: 301-308.
- 1068 12. Custovic A. "Asthma" or "Asthma Spectrum Disorder"? *J Allergy Clin Immunol Pract* 2020; 8: 2628-1069 2629.
- 13. Haider S, Granell R, Curtin J, Fontanella S, Cucco A, Turner S, Simpson A, Roberts G, Murray CS,
   Holloway JW, Devereux G, Cullinan P, Arshad SH, Custovic A. Modeling Wheezing Spells
   Identifies Phenotypes with Different Outcomes and Genetic Associates. *Am J Respir Crit Care* Med 2022; 205: 883-893.
- 1074 14. Custovic A, Henderson J, Simpson A. Does understanding endotypes translate to better asthma management options for all? *J Allergy Clin Immunol* 2019; 144: 25-33.
  - 15. Aaron SD, Vandemheen KL, FitzGerald JM, Ainslie M, Gupta S, Lemiere C, Field SK, McIvor RA, Hernandez P, Mayers I, Mulpuru S, Alvarez GG, Pakhale S, Mallick R, Boulet LP, Canadian Respiratory Research N. Reevaluation of Diagnosis in Adults With Physician-Diagnosed Asthma. *JAMA* 2017; 317: 269-279.
  - 16. Looijmans-van den Akker I, van Luijn K, Verheij T. Overdiagnosis of asthma in children in primary care: a retrospective analysis. *Br J Gen Pract* 2016; 66: e152-157.
  - 17. Robinson PFM, Fontanella S, Ananth S, Martin Alonso A, Cook J, Kaya-de Vries D, Polo Silveira L, Gregory L, Lloyd C, Fleming L, Bush A, Custovic A, Saglani S. Recurrent Severe Preschool Wheeze: From Prespecified Diagnostic Labels to Underlying Endotypes. *Am J Respir Crit Care Med* 2021; 204: 523-535.
  - 18. Howard R, Rattray M, Prosperi M, Custovic A. Distinguishing asthma phenotypes using machine learning approaches. *Current allergy and asthma reports* 2015; 15: 1-10.
  - 19. Oksel C, Granell R, Haider S, Fontanella S, Simpson A, Turner S, Devereux G, Arshad SH, Murray CS, Roberts G, Holloway JW, Cullinan P, Henderson J, Custovic A, Stelar investigators bTi.

    Distinguishing Wheezing Phenotypes from Infancy to Adolescence. A Pooled Analysis of Five Birth Cohorts. *Ann Am Thorac Soc* 2019; 16: 868-876.
  - 20. Custovic A, Ainsworth J, Arshad H, Bishop C, Buchan I, Cullinan P, Devereux G, Henderson J, Holloway J, Roberts G, Turner S, Woodcock A, Simpson A. The Study Team for Early Life Asthma Research (STELAR) consortium 'Asthma e-lab': team science bringing data, methods and investigators together. *Thorax* 2015; 70: 799-801.
  - 21. Golding J, Pembrey M, Jones R. ALSPAC-the Avon longitudinal study of parents and children. I. Study methodology. *Paediatric and perinatal epidemiology* 2001; 15: 74-87.
    - 22. Cullinan P, MacNeill SJ, Harris JM, Moffat S, White C, Mills P, Newman Taylor AJ. Early allergen exposure, skin prick responses, and atopic wheeze at age 5 in English children: a cohort study. *Thorax* 2004; 59: 855-861.
- 1101 23. Arshad SH, Holloway JW, Karmaus W, Zhang H, Ewart S, Mansfield L, Matthews S, Hodgekiss C,
   1102 Roberts G, Kurukulaaratchy R. Cohort Profile: The Isle Of Wight Whole Population Birth Cohort
   1103 (IOWBC). Int J Epidemiol 2018; 47: 1043-1044i.
- 24. Custovic A, Simpson BM, Murray CS, Lowe L, Woodcock A, Asthma NACM, Allergy Study G. The
   National Asthma Campaign Manchester Asthma and Allergy Study. *Pediatr Allergy Immunol* 2002; 13: 32-37.
- 1107 25. Martindale S, McNeill G, Devereux G, Campbell D, Russell G, Seaton A. Antioxidant intake in
   1108 pregnancy in relation to wheeze and eczema in the first two years of life. American journal of
   1109 respiratory and critical care medicine 2005; 171: 121-128.
- 26. Marchini J, Howie B. Genotype imputation for genome-wide association studies. *Nat Rev Genet* 2010; 11: 499-511.
- 27. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 2010; 26: 2190-2191.

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

- 28. Melen E, Himes BE, Brehm JM, Boutaoui N, Klanderman BJ, Sylvia JS, Lasky-Su J. Analyses of shared
   genetic factors between asthma and obesity in children. *J Allergy Clin Immunol* 2010; 126:
   631-637 e631-638.
- 29. Herrera-Luis E, Espuela-Ortiz A, Lorenzo-Diaz F, Keys KL, Mak ACY, Eng C, Huntsman S, Villar J,
   Rodriguez-Santana JR, Burchard EG, Pino-Yanes M. Genome-wide association study reveals a
   novel locus for asthma with severe exacerbations in diverse populations. *Pediatr Allergy Immunol* 2021; 32: 106-115.
- 30. Zhu Z, Guo Y, Shi H, Liu CL, Panganiban RA, Chung W, O'Connor LJ, Himes BE, Gazal S, Hasegawa K,
   1122 Camargo CA, Jr., Qi L, Moffatt MF, Hu FB, Lu Q, Cookson WOC, Liang L. Shared genetic and
   1123 experimental links between obesity-related traits and asthma subtypes in UK Biobank. J
   1124 Allergy Clin Immunol 2020; 145: 537-549.
- 31. Cusanovich DA, Billstrand C, Zhou X, Chavarria C, De Leon S, Michelini K, Pai AA, Ober C, Gilad Y.
   The combination of a genome-wide association study of lymphocyte count and analysis of
   gene expression data reveals novel asthma candidate genes. *Hum Mol Genet* 2012; 21: 2111 2123.
  - 32. Siroux V, Gonzalez JR, Bouzigon E, Curjuric I, Boudier A, Imboden M, Anto JM, Gut I, Jarvis D, Lathrop M, Omenaas ER, Pin I, Wjst M, Demenais F, Probst-Hensch N, Kogevinas M, Kauffmann F. Genetic heterogeneity of asthma phenotypes identified by a clustering approach. *Eur Respir J* 2014; 43: 439-452.
  - 33. Heinonen MT, Laine AP, Soderhall C, Gruzieva O, Rautio S, Melen E, Pershagen G, Lahdesmaki HJ, Knip M, Ilonen J, Henttinen TA, Kere J, Lahesmaa R, Finnish Pediatric Diabetes R. GIMAP GTPase family genes: potential modifiers in autoimmune diabetes, asthma, and allergy. *J Immunol* 2015; 194: 5885-5894.
  - 34. Denham S, Koppelman GH, Blakey J, Wjst M, Ferreira MA, Hall IP, Sayers I. Meta-analysis of genome-wide linkage studies of asthma and related traits. *Respir Res* 2008; 9: 38.
  - 35. Sottile G, Ferrante G, Torregrossa M, Cibella F, Cilluffo G, Fasola S, Alessandro R, Seidita G, Viegi G, La Grutta S. An association analysis to identify genetic variants linked to asthma and rhinoconjunctivitis in a cohort of Sicilian children. *Ital J Pediatr* 2019; 45: 16.
  - 36. Lutz SM, Cho MH, Young K, Hersh CP, Castaldi PJ, McDonald ML, Regan E, Mattheisen M, DeMeo DL, Parker M, Foreman M, Make BJ, Jensen RL, Casaburi R, Lomas DA, Bhatt SP, Bakke P, Gulsvik A, Crapo JD, Beaty TH, Laird NM, Lange C, Hokanson JE, Silverman EK, Investigators E, Investigators CO. A genome-wide association study identifies risk loci for spirometric measures among smokers of European and African ancestry. *BMC Genet* 2015; 16: 138.
  - 37. Vieira Braga FA, Kar G, Berg M, Carpaij OA, Polanski K, Simon LM, Brouwer S, Gomes T, Hesse L, Jiang J, Fasouli ES, Efremova M, Vento-Tormo R, Talavera-Lopez C, Jonker MR, Affleck K, Palit S, Strzelecka PM, Firth HV, Mahbubani KT, Cvejic A, Meyer KB, Saeb-Parsy K, Luinge M, Brandsma CA, Timens W, Angelidis I, Strunz M, Koppelman GH, van Oosterhout AJ, Schiller HB, Theis FJ, van den Berge M, Nawijn MC, Teichmann SA. A cellular census of human lungs identifies novel cell states in health and in asthma. *Nat Med* 2019; 25: 1153-1163.
- 38. Leoni G, Neumann PA, Kamaly N, Quiros M, Nishio H, Jones HR, Sumagin R, Hilgarth RS, Alam A,
   Fredman G, Argyris I, Rijcken E, Kusters D, Reutelingsperger C, Perretti M, Parkos CA,
   Farokhzad OC, Neish AS, Nusrat A. Annexin A1-containing extracellular vesicles and polymeric
   nanoparticles promote epithelial wound repair. J Clin Invest 2015; 125: 1215-1227.
- 39. Quach H, Rotival M, Pothlichet J, Loh YE, Dannemann M, Zidane N, Laval G, Patin E, Harmant C,
   Lopez M, Deschamps M, Naffakh N, Duffy D, Coen A, Leroux-Roels G, Clement F, Boland A,
   Deleuze JF, Kelso J, Albert ML, Quintana-Murci L. Genetic Adaptation and Neandertal
   Admixture Shaped the Immune System of Human Populations. *Cell* 2016; 167: 643-656 e617.
- 40. Võsa U, Claringbould A, Westra H-J, Bonder MJ, Deelen P, Zeng B, Kirsten H, Saha A, Kreuzhuber R,
   Kasela S, Pervjakova N, Alvaes I, Fave M-J, Agbessi M, Christiansen M, Jansen R, Seppälä I,
   Tong L, Teumer A, Schramm K, Hemani G, Verlouw J, Yaghootkar H, Sönmez R, Brown A,
   Kukushkina V, Kalnapenkis A, Rüeger S, Porcu E, Kronberg-Guzman J, Kettunen J, Powell J, Lee

- 1165 B, Zhang F, Arindrarto W, Beutner F, Brugge H, Dmitreva J, Elansary M, Fairfax BP, Georges M, 1166 Heijmans BT, Kähönen M, Kim Y, Knight JC, Kovacs P, Krohn K, Li S, Loeffler M, Marigorta UM, 1167 Mei H, Momozawa Y, Müller-Nurasyid M, Nauck M, Nivard M, Penninx B, Pritchard J, Raitakari 1168 O, Rotzchke O, Slagboom EP, Stehouwer CDA, Stumvoll M, Sullivan P, Hoen PACt, Thiery J, 1169 Tönjes A, van Dongen J, van Iterson M, Veldink J, Völker U, Wijmenga C, Swertz M, Andiappan 1170 A, Montgomery GW, Ripatti S, Perola M, Kutalik Z, Dermitzakis E, Bergmann S, Frayling T, van 1171 Meurs J, Prokisch H, Ahsan H, Pierce B, Lehtimäki T, Boomsma D, Psaty BM, Gharib SA, 1172 Awadalla P, Milani L, Ouwehand W, Downes K, Stegle O, Battle A, Yang J, Visscher PM, Scholz 1173 M, Gibson G, Esko T, Franke L. Unraveling the polygenic architecture of complex traits using 1174 blood eQTL metaanalysis. 2018: 447367.
- 1175 41. Gregory LG, Causton B, Murdoch JR, Mathie SA, O'Donnell V, Thomas CP, Priest FM, Quint DJ, Lloyd 1176 CM. Inhaled house dust mite induces pulmonary T helper 2 cytokine production. Clin Exp 1177 Allergy 2009; 39: 1597-1610.
  - 42. Perretti M, Chiang N, La M, Fierro IM, Marullo S, Getting SJ, Solito E, Serhan CN. Endogenous lipidand peptide-derived anti-inflammatory pathways generated with glucocorticoid and aspirin treatment activate the lipoxin A4 receptor. Nat Med 2002; 8: 1296-1302.
    - 43. D'Acquisto F, Paschalidis N, Raza K, Buckley CD, Flower RJ, Perretti M. Glucocorticoid treatment inhibits annexin-1 expression in rheumatoid arthritis CD4+ T cells. Rheumatology (Oxford) 2008; 47: 636-639.
  - 44. Vong L, Ferraz JG, Dufton N, Panaccione R, Beck PL, Sherman PM, Perretti M, Wallace JL. Upregulation of Annexin-A1 and lipoxin A(4) in individuals with ulcerative colitis may promote mucosal homeostasis. PLoS One 2012; 7: e39244.
    - 45. McArthur S, Juban G, Gobbetti T, Desgeorges T, Theret M, Gondin J, Toller-Kawahisa JE, Reutelingsperger CP, Chazaud B, Perretti M, Mounier R. Annexin A1 drives macrophage skewing to accelerate muscle regeneration through AMPK activation. J Clin Invest 2020; 130: 1156-1167.
- 1191 46. Rhen T, Cidlowski JA. Antiinflammatory action of glucocorticoids--new mechanisms for old drugs. N 1192 Engl J Med 2005; 353: 1711-1723.
- 1193 47. Lee SH, Lee PH, Kim BG, Seo HJ, Baek AR, Park JS, Lee JH, Park SW, Kim DJ, Park CS, Jang AS. 1194 Annexin A1 in plasma from patients with bronchial asthma: its association with lung function. 1195 BMC Pulm Med 2018; 18: 1.
- 1196 48. Eke Gungor H, Tahan F, Gokahmetoglu S, Saraymen B. Decreased levels of lipoxin A4 and annexin A1 in wheezy infants. Int Arch Allergy Immunol 2014; 163: 193-197.
  - 49. Ng FS, Wong KY, Guan SP, Mustafa FB, Kajiji TS, Bist P, Biswas SK, Wong WS, Lim LH. Annexin-1deficient mice exhibit spontaneous airway hyperresponsiveness and exacerbated allergenspecific antibody responses in a mouse model of asthma. Clin Exp Allergy 2011; 41: 1793-1803.
- 1201 50. Montoro DT, Haber AL, Biton M, Vinarsky V, Lin B, Birket SE, Yuan F, Chen S, Leung HM, Villoria J, 1202 Rogel N, Burgin G, Tsankov AM, Waghray A, Slyper M, Waldman J, Nguyen L, Dionne D, 1203 Rozenblatt-Rosen O, Tata PR, Mou H, Shivaraju M, Bihler H, Mense M, Tearney GJ, Rowe SM, 1204 Engelhardt JF, Regev A, Rajagopal J. A revised airway epithelial hierarchy includes CFTR-1205 expressing ionocytes. Nature 2018; 560: 319-324.
- 1206 51. Bonnelykke K, Sleiman P, Nielsen K, Kreiner-Moller E, Mercader JM, Belgrave D, den Dekker HT, 1207 Husby A, Sevelsted A, Faura-Tellez G, Mortensen LJ, Paternoster L, Flaaten R, Molgaard A, 1208 Smart DE, Thomsen PF, Rasmussen MA, Bonas-Guarch S, Holst C, Nohr EA, Yadav R, March ME, 1209 Blicher T, Lackie PM, Jaddoe VW, Simpson A, Holloway JW, Duijts L, Custovic A, Davies DE, 1210 Torrents D, Gupta R, Hollegaard MV, Hougaard DM, Hakonarson H, Bisgaard H. A genome-1211 wide association study identifies CDHR3 as a susceptibility locus for early childhood asthma 1212 with severe exacerbations. Nat Genet 2014; 46: 51-55.
- 1213 52. Schoettler N, Rodriguez E, Weidinger S, Ober C. Advances in asthma and allergic disease genetics: 1214 Is bigger always better? J Allergy Clin Immunol 2019; 144: 1495-1506.

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1197

1198

1199

- 53. Granell R, Henderson AJ, Timpson N, St Pourcain B, Kemp JP, Ring SM, Ho K, Montgomery SB,
   Dermitzakis ET, Evans DM, Sterne JA. Examination of the relationship between variation at
   17q21 and childhood wheeze phenotypes. J Allergy Clin Immunol 2013; 131: 685-694.
- 54. Spycher B, Henderson J, Granell R, Evans DM, Smith GD, Timpson NJ, Sterne JAC. Genome-wide
   prediction of childhood asthma and related phenotypes in a longitudinal birth cohort. *Eur Respir J* 2012; 40.
- 55. Savenije OE, Mahachie John JM, Granell R, Kerkhof M, Dijk FN, de Jongste JC, Smit HA, Brunekreef
   B, Postma DS, Van Steen K, Henderson J, Koppelman GH. Association of IL33-IL-1 receptor-like
   1 (IL1RL1) pathway polymorphisms with wheezing phenotypes and asthma in childhood. J
   Allergy Clin Immunol 2014; 134: 170-177.
  - 56. Crawford F CG, Chappell FM, et al. A systematic review and individual patient data meta-analysis of prognostic factors for foot ulceration in people with diabetes: the international research collaboration for the prediction of diabetic foot ulcerations (PODUS). Southampton (UK): NIHR Journals Library; 2015 Jul. (Health Technology Assessment, No. 19.57.) Chapter 5, Validation of the model. Available from: <a href="https://www.ncbi.nlm.nih.gov/books/NBK305598/">https://www.ncbi.nlm.nih.gov/books/NBK305598/</a>.
  - 57. Oksel C, Haider S, Fontanella S, Frainay C, Custovic A. Classification of Pediatric Asthma: From Phenotype Discovery to Clinical Practice. *Front Pediatr* 2018; 6: 258.
  - 58. Oksel C, Granell R, Mahmoud O, Custovic A, Henderson AJ, Stelar, Breathing Together i. Causes of variability in latent phenotypes of childhood wheeze. *J Allergy Clin Immunol* 2019; 143: 1783-1790 e1711.
  - 59. Savenije OE, Granell R, Caudri D, Koppelman GH, Smit HA, Wijga A, de Jongste JC, Brunekreef B, Sterne JA, Postma DS, Henderson J, Kerkhof M. Comparison of childhood wheezing phenotypes in 2 birth cohorts: ALSPAC and PIAMA. *J Allergy Clin Immunol* 2011; 127: 1505-1512 e1514.
  - 60. Torgerson DG, Ampleford EJ, Chiu GY, Gauderman WJ, Gignoux CR, Graves PE, Himes BE, Levin AM, Mathias RA, Hancock DB, Baurley JW, Eng C, Stern DA, Celedon JC, Rafaels N, Capurso D, Conti DV, Roth LA, Soto-Quiros M, Togias A, Li X, Myers RA, Romieu I, Van Den Berg DJ, Hu D, Hansel NN, Hernandez RD, Israel E, Salam MT, Galanter J, Avila PC, Avila L, Rodriquez-Santana JR, Chapela R, Rodriguez-Cintron W, Diette GB, Adkinson NF, Abel RA, Ross KD, Shi M, Faruque MU, Dunston GM, Watson HR, Mantese VJ, Ezurum SC, Liang L, Ruczinski I, Ford JG, Huntsman S, Chung KF, Vora H, Li X, Calhoun WJ, Castro M, Sienra-Monge JJ, del Rio-Navarro B, Deichmann KA, Heinzmann A, Wenzel SE, Busse WW, Gern JE, Lemanske RF, Jr., Beaty TH, Bleecker ER, Raby BA, Meyers DA, London SJ, Mexico City Childhood Asthma S, Gilliland FD, Children's Health S, study H, Burchard EG, Genetics of Asthma in Latino Americans Study SoG-E, Admixture in Latino A, Study of African Americans AG, Environments, Martinez FD, Childhood Asthma R, Education N, Weiss ST, Childhood Asthma Management P, Williams LK, Study of Asthma P, Pharmacogenomic Interactions by R-E, Barnes KC, Genetic Research on Asthma in African Diaspora S, Ober C, Nicolae DL. Meta-analysis of genome-wide association studies of asthma in ethnically diverse North American populations. *Nat Genet* 2011; 43: 887-
- 61. Wojcik GL, Graff M, Nishimura KK, Tao R, Haessler J, Gignoux CR, Highland HM, Patel YM, Sorokin EP, Avery CL, Belbin GM, Bien SA, Cheng I, Cullina S, Hodonsky CJ, Hu Y, Huckins LM, Jeff J, Justice AE, Kocarnik JM, Lim U, Lin BM, Lu Y, Nelson SC, Park SL, Poisner H, Preuss MH, Richard MA, Schurmann C, Setiawan VW, Sockell A, Vahi K, Verbanck M, Vishnu A, Walker RW, Young KL, Zubair N, Acuna-Alonso V, Ambite JL, Barnes KC, Boerwinkle E, Bottinger EP, Bustamante CD, Caberto C, Canizales-Quinteros S, Conomos MP, Deelman E, Do R, Doheny K, Fernandez-Rhodes L, Fornage M, Hailu B, Heiss G, Henn BM, Hindorff LA, Jackson RD, Laurie CA, Laurie CC, Li Y, Lin DY, Moreno-Estrada A, Nadkarni G, Norman PJ, Pooler LC, Reiner AP, Romm J, Sabatti C, Sandoval K, Sheng X, Stahl EA, Stram DO, Thornton TA, Wassel CL, Wilkens LR, Winkler CA, Yoneyama S, Buyske S, Haiman CA, Kooperberg C, Le Marchand L, Loos RJF, Matise TC, North

- 1265 KE, Peters U, Kenny EE, Carlson CS. Genetic analyses of diverse populations improves discovery for complex traits. *Nature* 2019; 570: 514-518.
- 1267 62. Boyd A, Golding J, Macleod J, Lawlor DA, Fraser A, Henderson J, Molloy L, Ness A, Ring S, Davey
  1268 Smith G. Cohort Profile: The 'Children of the 90s'--the index offspring of the Avon Longitudinal
  1269 Study of Parents and Children. *Int J Epidemiol* 2013; 42: 111-127.
- 1270 63. Fraser A, Macdonald-Wallis C, Tilling K, Boyd A, Golding J, Davey Smith G, Henderson J, Macleod J,
   1271 Molloy L, Ness A, Ring S, Nelson SM, Lawlor DA. Cohort Profile: the Avon Longitudinal Study of
   1272 Parents and Children: ALSPAC mothers cohort. *Int J Epidemiol* 2013; 42: 97-110.
  - 64. Martindale S, McNeill G, Devereux G, Campbell D, Russell G, Seaton A. Antioxidant intake in pregnancy in relation to wheeze and eczema in the first two years of life. *Am J Respir Crit Care Med* 2005; 171: 121-128.
  - 65. Atkinson W, Harris J, Mills P, Moffat S, White C, Lynch O, Jones M, Cullinan P, Newman Taylor AJ. Domestic aeroallergen exposures among infants in an English town. *Eur Respir J* 1999; 13: 583-589.
  - 66. Kurukulaaratchy RJ, Fenn M, Twiselton R, Matthews S, Arshad SH. The prevalence of asthma and wheezing illnesses amongst 10-year-old schoolchildren. *Respir Med* 2002; 96: 163-169.
- 1281 67. Kurukulaaratchy RJ, Fenn MH, Waterhouse LM, Matthews SM, Holgate ST, Arshad SH.
   1282 Characterization of wheezing phenotypes in the first 10 years of life. *Clin Exp Allergy* 2003; 33:
   1283 573-578.
  - 68. Loh PR, Danecek P, Palamara PF, Fuchsberger C, Y AR, H KF, Schoenherr S, Forer L, McCarthy S, Abecasis GR, Durbin R, A LP. Reference-based phasing using the Haplotype Reference Consortium panel. *Nat Genet* 2016; 48: 1443-1448.
- 1287 69. McCarthy S, Das S, Kretzschmar W, Delaneau O, Wood AR, Teumer A, Kang HM, Fuchsberger C, Danecek P, Sharp K, Luo Y, Sidore C, Kwong A, Timpson N, Koskinen S, Vrieze S, Scott LJ, Zhang 1288 1289 H, Mahajan A, Veldink J, Peters U, Pato C, van Duijn CM, Gillies CE, Gandin I, Mezzavilla M, 1290 Gilly A, Cocca M, Traglia M, Angius A, Barrett JC, Boomsma D, Branham K, Breen G, Brummett 1291 CM, Busonero F, Campbell H, Chan A, Chen S, Chew E, Collins FS, Corbin LJ, Smith GD, 1292 Dedoussis G, Dorr M, Farmaki AE, Ferrucci L, Forer L, Fraser RM, Gabriel S, Levy S, Groop L, 1293 Harrison T, Hattersley A, Holmen OL, Hveem K, Kretzler M, Lee JC, McGue M, Meitinger T, 1294 Melzer D, Min JL, Mohlke KL, Vincent JB, Nauck M, Nickerson D, Palotie A, Pato M, Pirastu N, 1295 McInnis M, Richards JB, Sala C, Salomaa V, Schlessinger D, Schoenherr S, Slagboom PE, Small K, 1296 Spector T, Stambolian D, Tuke M, Tuomilehto J, Van den Berg LH, Van Rheenen W, Volker U, 1297 Wijmenga C, Toniolo D, Zeggini E, Gasparini P, Sampson MG, Wilson JF, Frayling T, de Bakker 1298 PI, Swertz MA, McCarroll S, Kooperberg C, Dekker A, Altshuler D, Willer C, Iacono W, Ripatti S, 1299 Soranzo N, Walter K, Swaroop A, Cucca F, Anderson CA, Myers RM, Boehnke M, McCarthy MI, 1300 Durbin R, Haplotype Reference C. A reference panel of 64,976 haplotypes for genotype 1301 imputation. Nat Genet 2016; 48: 1279-1283.
  - 70. Durbin R. Efficient haplotype matching and storage using the positional Burrows-Wheeler transform (PBWT). *Bioinformatics* 2014; 30: 1266-1272.
  - 71. Pruim RJ, Welch RP, Sanna S, Teslovich TM, Chines PS, Gliedt TP, Boehnke M, Abecasis GR, Willer CJ. LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics* 2010; 26: 2336-2337.
- 1307 72. Dayem Ullah AZ, Oscanoa J, Wang J, Nagano A, Lemoine NR, Chelala C. SNPnexus: assessing the
   1308 functional relevance of genetic variation to facilitate the promise of precision medicine.
   1309 Nucleic Acids Res 2018; 46: W109-W113.
- 73. Stelzer G, Rosen N, Plaschkes I, Zimmerman S, Twik M, Fishilevich S, Stein TI, Nudel R, Lieder I,
   Mazor Y, Kaplan S, Dahary D, Warshawsky D, Guan-Golan Y, Kohn A, Rappaport N, Safran M,
   Lancet D. The GeneCards Suite: From Gene Data Mining to Disease Genome Sequence
   Analyses. Curr Protoc Bioinformatics 2016; 54: 1 30 31-31 30 33.

1274

1275

1276

1277

1278

1279

1280

1284

1285

1286


1302


1303


1304


1305


- 74. Durinck S, Moreau Y, Kasprzyk A, Davis S, De Moor B, Brazma A, Huber W. BioMart and
   Bioconductor: a powerful link between biological databases and microarray data analysis.
   Bioinformatics 2005; 21: 3439-3440.
- 75. Durinck S, Spellman PT, Birney E, Huber W. Mapping identifiers for the integration of genomic datasets with the R/Bioconductor package biomaRt. *Nat Protoc* 2009; 4: 1184-1191.
- 76. McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GR, Thormann A, Flicek P, Cunningham F. The Ensembl Variant Effect Predictor. *Genome Biol* 2016; 17: 122.
- 1321 77. Wingett S, Ewels P, Furlan-Magaril M, Nagano T, Schoenfelder S, Fraser P, Andrews S. HiCUP: pipeline for mapping and processing Hi-C data. *F1000Res* 2015; 4: 1310.
  - 78. Cairns J, Freire-Pritchett P, Wingett SW, Varnai C, Dimond A, Plagnol V, Zerbino D, Schoenfelder S, Javierre BM, Osborne C, Fraser P, Spivakov M. CHiCAGO: robust detection of DNA looping interactions in Capture Hi-C data. *Genome Biol* 2016; 17: 127.
  - 79. Adsera CB, Park YP, Meuleman W, Kellis M. Integrative analysis of 10,000 epigenomic maps across 800 samples for regulatory genomics and disease dissection. 2019: 810291.
  - 80. Ihaka R, Gentleman R. R: A Language for Data Analysis and Graphics. *Journal of Computational and Graphical Statistics* 1996; 5: 299-314.
  - 81. Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, Simonovic M, Doncheva NT, Morris JH, Bork P, Jensen LJ, Mering CV. STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res* 2019; 47: D607-D613.
  - 82. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* 2003; 13: 2498-2504.
    - 83. Brunekreef B, Smit J, de Jongste J, Neijens H, Gerritsen J, Postma D, Aalberse R, Koopman L, Kerkhof M, Wilga A, van Strien R. The prevention and incidence of asthma and mite allergy (PIAMA) birth cohort study: design and first results. *Pediatr Allergy Immunol* 2002; 13: 55-60.
    - 84. Wijga AH, Kerkhof M, Gehring U, de Jongste JC, Postma DS, Aalberse RC, Wolse AP, Koppelman GH, van Rossem L, Oldenwening M, Brunekreef B, Smit HA. Cohort profile: the prevention and incidence of asthma and mite allergy (PIAMA) birth cohort. *Int J Epidemiol* 2014; 43: 527-535.
    - 85. Branchett WJ, Walker SA, Lloyd CM. Experimental Mouse Models of Asthma and Analysis of CD4 T Cells. *Methods Mol Biol* 2021; 2285: 329-348.
















## Tissue specfic effect of rs75260654 on ANXA1 Expression









































FAM19A5→ L MIR4535→

←LOC284933 48.8 49 Position on chr22 (Mb)

MIR3201→

LINC01310→

49.2



