

# Longitudinal Associations Between Physical Health Conditions in Childhood and ADHD Symptoms at Age 17

**Running title:** ADHD AND PHYSICAL HEALTH ASSOCIATIONS

**Author names and affiliations:**

Mrs Claire Reed (MSc)<sup>1</sup>, Professor Samuele Cortese (MD, PhD)<sup>1,2,3,4,5</sup>, Professor Henrik Larsson (PhD)<sup>6,7</sup>, Professor Cédric Galéra (MD, PhD)<sup>8,9,10,11</sup>, Dr Joanne Cotton (PhD)<sup>12</sup>, Dr Valerie Brandt (PhD)<sup>1</sup>

<sup>1</sup>University of Southampton, Southampton, UK

<sup>2</sup>Solent NHS Trust, Southampton, UK

<sup>3</sup>University of Nottingham, Nottingham, UK

<sup>4</sup>Hassenfeld Children's Hospital at NYU Langone, NY, USA

<sup>5</sup>University Child Study Center, New York, NY, USA.

<sup>6</sup>Karolinska Institute, Stockholm, Sweden

<sup>7</sup>Örebro University, Örebro, Sweden

<sup>8</sup>University of Bordeaux, Bordeaux, France

<sup>9</sup>INSERM, Bordeaux Population Health Center, UMR1219, Bordeaux, France

<sup>10</sup>Centre Hospitalier Perrens, Bordeaux, France

<sup>11</sup>Research Unit on Children's Psychosocial Maladjustment, Montreal, QC, Canada

<sup>12</sup>University of Cambridge, Cambridge, UK

**Corresponding author:** Claire Reed, c.l.reed@soton.ac.uk; University of Southampton, Southampton, UK.

**Funding for this study:** This study received no funding.

**Acknowledgements:** We are grateful to the Centre for Longitudinal Studies (CLS), UCL Social Research Institute, for the use of these data and to the UK Data Service for making them available. However, neither CLS nor the UK Data Service bear any responsibility for the analysis or interpretation of these data.

**Financial disclosures:** Henrik Larsson acknowledges financial support from the European Union's Horizon 2020 research and innovation programme under grant agreement number 965381, the Swedish Research Council (2018-02599; 2022-01119) and the Swedish Brain Foundation (FO2021-0115). Valerie Brandt acknowledges financial support from the Academy of Medical Sciences.

**Conflict of interest:** Henrik Larsson reports receiving grants from Shire Pharmaceuticals; personal fees from and serving as a speaker for Medice, Shire/Takeda Pharmaceuticals and Evolan Pharma AB; and sponsorship for a conference on attention-deficit/hyperactivity disorder from Shire/Takeda Pharmaceuticals and Evolan Pharma AB, all outside the submitted work. Henrik Larsson is editor-in-chief of JCPP Advances. Prof. Cortese has

declared honoraria and reimbursement for travel and accommodation expenses for lectures from the following non-profit associations: Association for Child and Adolescent Central Health (ACAMH), Canadian ADHD Alliance Resource (CADDRA), British Association of Pharmacology (BAP), and from Healthcare Convention for educational activity on ADHD. Joanne Cotton was funded by an ESRC/UKRI postdoctoral fellowship, Grant number: G108309 (October 2020 - August 2022).CR, CG and VB have no COIs to declare.

**Keywords:** ADHD, physical health, environmental factors, ADHD medication.

## Abstract

**Objective:** While evidence suggests significant cross-sectional relationships between Attention-Deficit/Hyperactivity Disorder (ADHD) and several physical health conditions, less is known about their longitudinal associations. We investigated the cumulative effect of childhood physical health conditions on ADHD symptoms at age 17, controlling for environmental factors, ADHD medication status and ADHD symptoms at age three.

**Method:** Using Millennium Cohort Study data (Weighted  $n=8,059$ ), we assessed whether four physical health clusters (sensory, neurological, atopic, and cardio-metabolic) were associated with scores on the ADHD subscale from the Strengths and Difficulties Questionnaire at age 17. Environmental factors were grouped into five cumulative risk indices: prenatal, perinatal, postnatal environment, postnatal maternal wellbeing, and sociodemographic factors. Regression analyses determined whether each physical health cluster was associated with ADHD score while controlling for environmental factors, ADHD medication and earlier symptoms.

**Results:** Sensory, neurological, and cardio-metabolic clusters were all significantly associated with ADHD symptoms ( $\beta$  range = 0.04-0.09,  $p < .001$ ). The overall model explained 2% of the variance. This rose to 21% ( $\Delta R^2 = .06$ ) after adjusting for confounders. The sensory ( $\beta = 0.06$ ) and neurological ( $\beta = 0.06$ ) clusters remained significant ( $R^2 = .21$ ,  $\Delta R^2 = .06$ ) but the cardio-metabolic cluster was no longer a significant predictor.

**Conclusion:** Sensory or neurological conditions in childhood were associated with higher ADHD symptoms aged 17 after adjustment of confounders. This was not the case for atopic or cardio-metabolic conditions. These findings have implications for the care of children with sensory/neurological conditions and future research examining ADHD etiopathophysiology.

## Introduction

Attention-Deficit/Hyperactivity Disorder (ADHD) is characterised by developmentally atypical, pervasive, and impairing symptoms of inattention and/or hyperactivity/impulsivity.<sup>1</sup> ADHD is a complex condition, with many comorbidities often complicating the diagnostic process, symptom management and the quality of life of individuals with ADHD and their families.<sup>2</sup>

There is increasing evidence that ADHD is associated not only with mental health but also with physical conditions. A recent umbrella review<sup>3</sup> encompassing ten meta-analyses on ADHD found significant cross-sectional associations of ADHD with asthma and obesity when considering the most rigorous estimates (adjusted effect sizes). When considering unadjusted effect sizes, there was also convincing or highly suggestive evidence for an association of ADHD with rhinitis and dermatitis.

Another recent birth cohort study<sup>4</sup> (n = 2057) assessed cross-sectional associations between ADHD symptoms and a large number of physical conditions at different time points during development, from five months to 17 years. Some of the associations between ADHD symptoms and several physical conditions were consistently found in early childhood, middle childhood, and adolescence (e.g., asthma, sleep problems), while others were present only at one time point, or were confounded by socioeconomic status or psychiatric comorbidities (e.g., body mass index, dental caries). It is possible that either ADHD contributes to the increased risk for physical conditions, or that, vice versa, physical conditions increase the risk of ADHD or that common underlying factors contribute to both.<sup>5</sup>

However, while the bulk of the studies on the links between ADHD and physical conditions have addressed cross-sectional associations, longitudinal associations shedding light on the direction of the relationship have been less explored and have focused on single physical conditions, e.g., obesity/increased body mass index.<sup>6</sup>

To address these limitations, the present study used a large-scale population-representative sample to examine the hypothesis that a cumulative number of physical health conditions across child development are associated with ADHD symptoms in adolescence.

When assessing this relationship, it is important to control for two important confounders: 1) environmental factors associated with an increased risk of ADHD. In this regard, there is meta-analytic evidence that ADHD is more common in children whose mothers smoked during pregnancy or suffered pre-eclampsia.<sup>7</sup> Other risk factors include low birth weight and prematurity<sup>8</sup>, and several indicators of socioeconomic disadvantage such as low family income, single parent households and low maternal education.<sup>9</sup> 2) medications for ADHD that may impact on a number of physiologic systems,- for instance, stimulants can negatively affect (at least at the beginning of the treatment) weight <sup>10</sup> and sleep.<sup>11</sup>

To gain insight into the longitudinal relationship between physical conditions and ADHD symptom development, we explored whether a cumulative number of physical disorders in childhood predicted ADHD symptoms at age 17, controlling for cumulative environmental risk, ADHD medication use and ADHD symptoms reported at age three.

## Methods

### Sample

We used data from the Millennium Cohort Study (MCS), a longitudinal study that collected information from over 19,000 UK families with children born between 2000 and 2002. To date, seven waves of data have been collected, at ages 9 months and 3, 5, 7, 11, 14, and 17 years, respectively. In this study, as in previous research<sup>12</sup>, only the first sibling of each family was included (4.9% excluded) to prevent confounding effects of family membership. Participants were also excluded if data relating to their biological mother were unavailable (5.1%) or all physical health predictor variables were missing (1.1%). As in previous longitudinal analyses<sup>13</sup>, only participants with non-missing outcome data at age 17 were included in the analysis.

The number of participating families varied between waves, and households sampled from disadvantaged areas were more likely to withdraw.<sup>14</sup> As such, standardised weights were applied to ensure the sample was representative of the UK population and to account for attrition rates. Weights were calculated by multiplying the wave one sample design weights by the non-response weights of each wave. As data in this study were taken from all waves, weights from the most recent data sweep were applied. Descriptive statistics for all weighted variables in the current sample for the study (n=8059) are reported in Tables 1 and 2. This study was approved by the Ethics committee of University of [anonymised], in addition to the informed consent procedures obtained in the MCS.<sup>15</sup> Parents gave written informed consent to the open use of the anonymised dataset.

### Measures

#### *Outcome measure – parent-reported ADHD symptoms*

The parent-reported hyperactivity/inattention subscale of the Strengths and Difficulties Questionnaire (SDQ<sup>16</sup>), measured at age 17, was used as a continuous dependent variable. The

subscale hyperactivity/inattention is composed of five items ("restless, overactive, cannot stay still for long", "constantly fidgeting or squirming", "easily distracted, concentration wanders", "thinks things out before acting", "good attention span, sees tasks through to the end"). The SDQ items are rated from 0 to 2 points (0= never, 1= sometime true, 2= certainly true- or reverse coded for positive statements). Scores on the hyperactivity/inattention subscale range between 0 and 10, with a higher score indicating more ADHD symptoms. The SDQ has been found to be a valid measure with the hyperactivity/inattention subscale predicting ADHD diagnosis<sup>17</sup> with a Cronbach's alpha of 0.88, indicating good reliability. In the current sample, SDQ scores are significantly correlated ( $p < .001$ ) across all six waves (ages 3-17 years) in which they are reported.

ADHD was diagnosed in a subset of children, with  $n = 174$  children having an ADHD diagnosis by age 17. The main analysis was replicated with diagnosed ADHD as the outcome variable to test robustness of the results. We found that adolescents with an ADHD diagnosis at age 17 had a significantly higher hyperactivity / inattention SDQ score ( $M = 5.09$ ) than adolescents without an ADHD diagnosis ( $M = 2.38$ ;  $t(8386) = 16.31$ ,  $p < .001$ ,  $d = 1.25$ ).

### ***Predictors - Physical Health Conditions***

At each wave of data collection (age 9 months to 17 years), parents were asked about their child's physical health. This information was gathered in several ways, including asking whether they had ever seen a health care professional about any problems and, if yes, asking to give details. Parents were also asked about specific diagnoses (e.g., 'has -child's name- ever had asthma?'). Hospital admissions were also recorded along with the reason for admission, such as convulsions, fits or epilepsy. We created variables to identify whether each child had ever experienced the condition. Physical conditions where a diagnosis could be given repeatedly and on a temporary basis, such as infection or diarrhoea, were excluded.

We grouped the physical conditions into four clusters, according to similar physiological mechanisms, reflecting their grouping in the ICD-11, as well as existing empirical evidence.<sup>18-23</sup> a) Sensory cluster (eyesight problems and hearing problems), b) Atopic cluster (eczema, asthma, and hay fever), c) Neurological cluster (fits and epilepsy, sleeping problems, stutter, movement problems and other neurological disorders), d) Cardio-metabolic cluster (obesity [at wave 7 only to exclude the natural weight fluctuations of young children], diabetes, and any reported heart or circulation condition including congenital heart disease). Cohort members' cumulative physical risk was therefore indicated by the number of conditions they had ever had in each cluster.

***Control Variables - Cumulative Risk Indices (CRIs) and age three SDQ score.***

Informed by previous evidence (Table 1), 20 environmental risk factors for ADHD were identified and controlled for within the available MCS data.<sup>24</sup> Data were collected during interviews with parents at the first data sweep (age 9 months). Individual risk factors were classified as either 'high risk' or 'low risk' (e.g., caesarean section yes/ no). Continuous variables were assigned a cut-off with reference to previous literature (Table 1). Risk factors were grouped and totalled to create five cumulative risk indices (CRIs) as follows: 1) *Prenatal risk index*, including maternal pre-pregnancy BMI, blood pressure problems in pregnancy, maternal diabetes and smoking during pregnancy, 2) *Perinatal risk index*, including birth weight, gestation, delivery method, and pregnancy lie/presentation, 3) *Postnatal risk (environment) index*, including maternal age, number of parents in the household, maternal education, and breastfeeding, 4) *Postnatal risk (maternal wellbeing) index*, including maternal distress, maternal depression/anxiety, and maternal attachment. Maternal distress was assessed using a 9-item short form version<sup>25</sup> of the Rutter Malaise Inventory.<sup>26</sup> The highest loading items in the first principal factor were used to create the 9-item scale measuring psychological



distress. Mothers were asked a series of questions requiring a yes/no response such as “Do you often feel miserable or depressed?” High risk maternal distress was indicated by a score of  $\geq 4$ <sup>27</sup> and psychometric properties are reported to be good.<sup>28</sup> A diagnosis of depression and/or anxiety was also included in the maternal wellbeing index, along with maternal attachment measured using a 6-item modified version of the Condon Maternal Attachment Questionnaire.<sup>29</sup> In this scale, mothers are asked about their feelings in different situations such as when they are caring for or having to leave their baby. Responses were recoded to ensure all items followed the same 1-5 scale which was then summed to give a total score out of 30. In line with previous literature,<sup>30</sup> high risk was categorised as the 25% of scores with the lowest attachment. 5) *Socioeconomic status and demographic risk index* including sex, ethnicity, household income, housing tenure and overcrowding.

To account for ADHD symptoms experienced at a younger age, SDQ score reported by parents at age three was added as an additional predictor.

### ***ADHD medication***

Only at wave six, when cohort members were 14, parents were asked if their child was taking any prescribed medication for ADHD. A list of possible medications was given, including both stimulants and non-stimulants and medication, and the variable was coded as yes if parents indicated their child was taking any medication on the list.

### **Statistical Analysis**

Stepwise, multiple regression analysis was conducted using IBM SPSS Statistics for Windows (Version 28.0, Chicago, Illinois) to analyse the relationship between the physical health clusters and ADHD symptoms at age 17 before and after controlling for cumulative environmental risk, ADHD medication and age three SDQ score. SDQ scores on the hyperactivity/inattention

subscale at age 17 were entered as the dependent variable in all four regression models. The predictors entered into each model were as follows: Model 1- physical health clusters; model 2- physical health clusters and cumulative risk indices; model 3- physical health clusters, cumulative risk indices and ADHD medication; model 4- physical health clusters, cumulative risk indices, ADHD medication and SDQ score at age three. Variance inflation factor values did not indicate multicollinearity for any of the variables. To ensure correct temporal ordering of the regression model, physical health diagnoses from all timepoints were used to predict ADHD symptoms and diagnosis at age 17. However, to confirm robustness of the findings when accounting for earlier ADHD symptoms, we included in the main analysis a fourth regression model with SDQ score at age three as an additional predictor. To show further robustness of the results, model 3 was repeated as a binary logistic regression with ADHD diagnosis (yes/no) as the outcome.

## Results

Weighted frequencies of physical health clusters and cumulative risk indices are reported in Table 2. As clusters were formed based on physiological similarity, they consisted of differing numbers of conditions and proportional frequencies. For example, the cardio-metabolic cluster consisted of three conditions with obesity markedly more prevalent than the other two disorders, whereas the sensory cluster consisted of two conditions with similar frequencies. In total, 91 children (1%) were taking ADHD medication and the average SDQ score at age 17 was 2.64 ( $SD = 2.34$ ). Skewness and kurtosis were found to be within the normal range.

Model 1 (physical health clusters only) was overall significant ( $F(4,5411) = 29.37, p < .001$ ) and explained 2% of the variance. As individual predictors, the sensory ( $\beta = 0.08, p < .001$ ), neurological ( $\beta = 0.10, p < .001$ ) and cardio-metabolic ( $\beta = 0.04, p < .001$ ) clusters all significantly predicted ADHD symptoms at age 17. Model 2 (physical health clusters + CRIs) was also statistically significant ( $F(9,5406) = 61.59, p < .001$ ) with 10% of the variance explained ( $R^2_{adj} = .09, \Delta R^2 = .07$ ). Again, the sensory ( $\beta = 0.08, p < .001$ ) and neurological ( $\beta = 0.09, p < .001$ ) clusters were significantly associated with ADHD symptoms but the cardio-metabolic cluster was no longer a significant predictor. This remained unchanged in model 3 ( $F(10,5405) = 96.37, p < .001$ ) with only the sensory ( $\beta = 0.07, p < .001$ ) and neurological ( $\beta = 0.09, p < .001$ ) clusters significantly predicting ADHD symptoms at age 17. Overall, model 3 explained 15% of the variance ( $R^2_{adj} = .15, \Delta R^2 = .06$ ). Model 4 was again overall significant ( $F(11,5404) = 129.27, p < .001$ ). The model explained 21% of the variance ( $R^2_{adj} = .21, \Delta R^2 = .06$ ) with the sensory ( $\beta = 0.06, p < .001$ ) and neurological cluster ( $\beta = 0.06, p < .001$ ) significantly predicting age 17 ADHD symptoms. The atopic cluster was not a significant predictor in any of the four regression models. The CRIs all significantly predicted ADHD symptoms in both model 2 and model 3 ( $\beta$  range = 0.03-0.18). In model 4, all CRIs significantly predicted ADHD symptoms except perinatal factors. A summary of the regression statistics

can be found in Table 3. The mean SDQ score for each cumulative number of conditions in each cluster are shown in Figure 1.

A binary logistic regression with ADHD diagnosis as the outcome variable showed that the sensory cluster significantly predicted ADHD diagnosis, ( $b = 0.27, p = .020, OR = 1.31$  (95% CI: 1.04, 1.65)).

The neurological cluster also significantly predicted ADHD diagnosis, ( $b = 0.66, p < .001, OR = 1.94$  (95% CI: 1.48, 2.53)). Each additional neurological condition almost doubled the odds of an ADHD diagnosis.

## Discussion

This is the first study that analyzed the longitudinal association between a broad range of physical conditions in childhood and ADHD symptoms at age 17, controlling for cumulative environmental risk factors for ADHD, ADHD medication status and earlier ADHD symptoms. The sensory and neurological clusters were both significant predictors of hyperactivity/inattention symptoms at age 17 and remained so after controlling for confounders. Moreover, the odds of having an ADHD diagnosis at age 17 was almost twice as likely when adolescents had predated neurological issues. The atopic cluster was not a significant predictor in any of the regression models and, after controlling for environmental risk and ADHD medication, the cardio-metabolic cluster was also not a significant predictor.

Children with more sensory conditions had higher hyperactivity and inattention symptoms at age 17. This is in line with previous studies showing that both eyesight and hearing problems increase the risk of ADHD<sup>31-33</sup>. Our findings build on this known association between ADHD and eyesight/hearing problems by showing that the likelihood of hyperactivity and inattention symptoms increases as the number of sensory conditions increases. DeCarlo and colleagues<sup>33</sup> proposed that sensory disorders place a large demand on executive functioning. Sensory problems may deplete executive functioning skills, leading to a reduced ability to successfully perform tasks known to be impaired in ADHD, such as those requiring organisation and concentration. It follows, therefore, that an increased number of sensory conditions may exacerbate inattention or hyperactivity symptoms in people who may already struggle with executive functioning.

This raises the question, which should be explored in future research, of whether hearing and eyesight problems magnify existing symptoms, mimic ADHD symptoms that may otherwise not be present, or whether ADHD and sensory problems may share neurobiological underpinnings.<sup>34</sup> Our findings suggest that, to maintain quality of life, people with sensory

disorders may need support with symptoms beyond those primarily associated with their health condition, because people with high levels of ADHD symptoms often experience a poorer quality of life, even if they do not meet the criteria for a diagnosis of ADHD.<sup>35</sup>

A higher number of neurological conditions during development also significantly predicted ADHD symptoms at age 17 when controlling for risk factors, medication and earlier ADHD symptoms. This is in line with previous studies reporting associations between ADHD and individual neurological conditions, such as stutter<sup>36</sup> and epilepsy.<sup>37</sup> Sleep problems, stutter and epilepsy are all known to individually predict ADHD, as well as each other.<sup>21</sup> Our results expand this knowledge by showing that a cumulation of neurological conditions further increases the likelihood of experiencing ADHD symptoms and an ADHD diagnosis later on in adolescence.

Previous literature has suggested this relationship may be bi-directional<sup>38</sup> as ADHD and conditions such as epilepsy may have shared risk factors or a shared neurobiological origin, with each predicting an increased risk of the other. Previous studies have found a shared genetic association between ADHD and neurological conditions, such as epilepsy<sup>39</sup> and sleep problems.<sup>40</sup> It is possible that some genetic variants affect several neurological, as well as mental conditions, increasing polygenic risk.<sup>41</sup>

Neurological conditions such as epilepsy are also known to impact executive functioning networks<sup>42</sup> which may modulate ADHD symptom expression. Executive functions involve the frontal cortex<sup>43</sup> which is associated with both epilepsy and ADHD.<sup>39</sup> Alterations in executive functioning networks which are associated with ADHD and neurological conditions may offer another explanation of the comorbidity between the conditions.

Contrary to previous research,<sup>44</sup> atopic disorders were not significantly associated with higher ADHD symptoms. This result was surprising considering the wealth of evidence to support an association between atopic conditions and ADHD. Both eczema and hay fever had frequencies

higher than expected and notably higher than other conditions (42.3% and 30.8% respectively). It may be that these conditions were over-reported in the sample, which influenced the results. Future research utilising clinical diagnoses rather than parent-reported physical conditions may reduce measurement error and shed light on the association between cumulative atopic conditions and ADHD symptoms. Another possible explanation for our findings is that individual conditions predict ADHD symptoms but that an accumulation of conditions does not increase the likelihood further.

This may also be the case for cardio-metabolic disorders as the cardio-metabolic cluster also produced different results to the sensory and neurological clusters. The cumulative effect of cardio-metabolic conditions significantly predicted ADHD symptoms in the first regression model but not when controlling for environmental risk or ADHD medication. Previous research has reported an increased likelihood of developing diabetes in young people with ADHD, even after controlling for medication and demographic factors<sup>45</sup>. Our findings suggest that this relationship may not be bidirectional as cardiometabolic conditions did not predict an increase in hyperactivity/inattention symptoms. Rather than an accumulation of cardiometabolic conditions being associated with an increase in hyperactivity/inattention symptoms, it may be that environmental factors, or even other ADHD-related features such as impulsive decision making, are driving this relationship. These findings add weight to the complexity of the association between physical health and ADHD, demonstrating the influence of environmental risk on this relationship. For cardio-metabolic conditions at least, an individual's environment may play a larger part in determining the nature of the relationship.

All cumulative risk indices were themselves significantly associated with ADHD symptoms at age 17. The postnatal mother environment and SES/demographics indices had the largest effect size, and both consisted of environmental factors which are considered to reflect mainly psychosocial rather than biological influences compared to the other indices. For example,

gestation and birth weight are irreversible factors which affect a child's biology at a fixed point in time, potentially with long term consequences. Factors affecting the whole household, such as income or housing tenure, have a greater effect on a child's physical environment but are also subject to greater fluctuation and their influence may vary depending on changes over time. This suggests further investigation is warranted to explore how different environmental factors affect the relationship between cumulative physical conditions and ADHD.

This study has several strengths. It used a large sample, weighted to represent the general population. Data were obtained longitudinally, allowing analysis of environmental risk before the age of three as a predictor of ADHD symptoms in later adolescence. It is also the first study to control for both environmental risk and medication use while examining the cumulative effect of several physical health conditions. The results further our insight into the pathophysiology of ADHD and point to associations that may eventually inform clinical practice in terms of implementation of additional support programmes. A key area for future research would be to explore the effects of early treatment and support for physical conditions on later development of ADHD symptoms. However, when considering their possible clinical implications, our findings should be interpreted with caution. Although the SDQ differentiated well between children with and without an ADHD diagnosis, it is not a screening instrument for ADHD. The sensitivity analyses that we conducted with ADHD diagnosis are limited by only a subset of children having been evaluated for ADHD, therefore, our figures are likely underestimated. However, the consistency of the results in our sensitivity analysis indicates that our findings may extend to cases with a categorical diagnosis. There are some additional limitations to consider. First, the analyses were restricted by the amount and variety of physical health and environmental data available. Second, this study only looked at physical health as a predictor of ADHD symptoms and did not explore the reverse relationship nor the possibility of a bi-directional association. Third, due to a lack of data, it was not possible to analyse the



effect of parental history of ADHD or physical health. ADHD is known to be highly heritable<sup>46</sup> as are many physical health conditions. Therefore, it is likely that a parent's health and neurodiversity may impact both ADHD symptoms and physical health of their child as well as the home environment.

In summary, this study found a significant relationship between a number of cumulative physical disorders diagnosed during childhood and ADHD symptoms in adolescence. The results indicate that sensory disorders and neurological disorders predicted ADHD symptoms even when relevant environmental risk is controlled for, suggesting possible biological commonalities including genetic association.

## References

1. American Psychiatric Association. *Diagnostic and statistical manual of mental disorders, (DSM-5), 5th edn American Psychiatric Publishing Arlington, VA.* vol 890425558. ISBN-10. 2013. <https://doi.org/10.1176/appi.books.9780890425596>
2. Reale L, Bartoli B, Cartabia M, et al. Comorbidity prevalence and treatment outcome in children and adolescents with ADHD. *European child & adolescent psychiatry.* 2017;26(12):1443-1457. doi: 10.1007/s00787-017-1005-z.
3. Arrondo G, Solmi M, Dragioti E, et al. Associations between mental and physical conditions in children and adolescents: An umbrella review. *Neuroscience & Biobehavioral Reviews.* 2022:104662. <https://doi.org/10.1016/j.neubiorev.2022.104662>
4. Galéra C, Cortese S, Orri M, et al. Medical conditions and Attention-Deficit/Hyperactivity Disorder symptoms from early childhood to adolescence. *Molecular Psychiatry.* 2021:1-9. DOI: [10.1038/s41380-021-01357-x](https://doi.org/10.1038/s41380-021-01357-x)
5. Hané T, Cortese S. Attention deficit/hyperactivity-disorder and obesity: a review and model of current hypotheses explaining their comorbidity. *Neuroscience & Biobehavioral Reviews.* 2018;92:16-28. DOI: [10.1016/j.neubiorev.2018.05.017](https://doi.org/10.1016/j.neubiorev.2018.05.017)
6. Kase BE, Rommelse N, Chen Q, et al. Longitudinal associations between symptoms of ADHD and BMI from late childhood to early adulthood. *Pediatrics.* 2021;147(6) DOI: [10.1542/peds.2020-036657](https://doi.org/10.1542/peds.2020-036657)
7. Arango C, Dragioti E, Solmi M, et al. Risk and protective factors for mental disorders beyond genetics: an evidence-based atlas. *World Psychiatry.* 2021;20(3):417-436. DOI: [10.1002/wps.20894](https://doi.org/10.1002/wps.20894)
8. Serati M, Barkin JL, Orsenigo G, Altamura AC, Buoli M. Research Review: The role of obstetric and neonatal complications in childhood attention deficit and hyperactivity

disorder—a systematic review. *Journal of Child Psychology and Psychiatry*.

2017;58(12):1290-1300. DOI: [10.1111/jcpp.12779](https://doi.org/10.1111/jcpp.12779)

9. Russell AE, Ford T, Williams R, Russell G. The association between socioeconomic disadvantage and attention deficit/hyperactivity disorder (ADHD): a systematic review. *Child Psychiatry & Human Development*. 2016;47(3):440-458. DOI: [10.1007/s10578-015-0578-3](https://doi.org/10.1007/s10578-015-0578-3)

10. Carucci S, Balia C, Gagliano A, et al. Long term methylphenidate exposure and growth in children and adolescents with ADHD. A systematic review and meta-analysis. *Neuroscience & Biobehavioral Reviews*. 2021;120:509-525.

DOI: [10.1016/j.neubiorev.2020.09.031](https://doi.org/10.1016/j.neubiorev.2020.09.031)

11. Faraone SV, Po MD, Komolova M, Cortese S. Sleep-associated adverse events during methylphenidate treatment of attention-deficit/hyperactivity disorder: A meta-analysis. *The Journal of clinical psychiatry*. 2019;80(3):19675. DOI: [10.4088/JCP.18r12210](https://doi.org/10.4088/JCP.18r12210)

12. Flouri E, Midouhas E, Ruddy A, Moulton V. The role of socio-economic disadvantage in the development of comorbid emotional and conduct problems in children with ADHD. *European child & adolescent psychiatry*. 2017;26(6):723-732.

DOI: [10.1007/s00787-017-0940-z](https://doi.org/10.1007/s00787-017-0940-z)

13. Brandt V, Hall C, Eisenbarth H, Hall J. Mutual relationships between head injury and conduct problems in children aged 9 months to 14 years in the UK Millennium Cohort Study. *medRxiv*. 2022; <https://doi.org/10.1101/2022.01.09.22268970>

14. Ploubidis, GB; Mostafa, T; (2017) Millennium Cohort: Study Sixth Survey 2015-2016. Technical report on response (Age 14). (Centre for Longitudinal Studies ). UCL Institute of Education: London, UK

15. Shepherd P, Gilbert E. British Cohort Study ethical review and consent. *London: Centre for Longitudinal Studies*. 2019;

16. Goodman R. The Strengths and Difficulties Questionnaire: a research note. *Journal of child psychology and psychiatry*. 1997;38(5):581-586. <https://doi.org/10.1111/j.1469-7610.1997.tb01545.x>
17. Hall CL, Guo B, Valentine AZ, et al. The validity of the Strengths and Difficulties Questionnaire (SDQ) for children with ADHD symptoms. *PloS one*. 2019;14(6):e0218518. DOI: [10.1371/journal.pone.0218518](https://doi.org/10.1371/journal.pone.0218518)
18. Organization WH. International classification of diseases—Eleventh revision. *Geneva, Switzerland: World Health Organization*. 2019;
19. Chen M-H, Su T-P, Chen Y-S, et al. Is atopy in early childhood a risk factor for ADHD and ASD? A longitudinal study. *Journal of psychosomatic research*. 2014;77(4):316-321. DOI: [10.1016/j.jpsychores.2014.06.006](https://doi.org/10.1016/j.jpsychores.2014.06.006)
20. Eklund H, Findon J, Cadman T, et al. Needs of adolescents and young adults with neurodevelopmental disorders: comparisons of young people and parent perspectives. *Journal of Autism and Developmental Disorders*. 2018;48:83-91. doi: [10.1007/s10803-017-3295-x](https://doi.org/10.1007/s10803-017-3295-x)
21. Merlo S, Briley PM. Sleep problems in children who stutter: evidence from population data. *Journal of Communication Disorders*. 2019;82:105935. DOI: [10.1016/j.jcomdis.2019.105935](https://doi.org/10.1016/j.jcomdis.2019.105935)
22. Pruetz DG, Shaw DM, Chen H-H, et al. Identifying developmental stuttering and associated comorbidities in electronic health records and creating a phenome risk classifier. *Journal of fluency disorders*. 2021;68:105847. DOI: [10.1016/j.jfludis.2021.105847](https://doi.org/10.1016/j.jfludis.2021.105847)

23. Garcia-Argibay M, du Rietz E, Lu Y, et al. The role of ADHD genetic risk in mid-to-late life somatic health conditions. *Translational Psychiatry*. 2022;12(1):152.  
DOI: [10.1038/s41398-022-01919-9](https://doi.org/10.1038/s41398-022-01919-9)
24. University of London IoE, Centre for Longitudinal Studies. Data from: Millennium Cohort Study: First Survey, 2001-2003. 2022;14th Edition. doi:10.5255/UKDA-SN-4683-6
25. Dex S, Joshi H. *Millennium Cohort Study First Survey: A User's Guide to Initial Findings*. Centre for Longitudinal Studies, Institute of Education, University of London; 2004.
26. Rutter M, Tizard J, Whitmore K. Health, education, and behaviour. *London, ENG: Longmans Green*. 1970;
27. Noonan K, Burns R, Violato M. Family income, maternal psychological distress and child socio-emotional behaviour: Longitudinal findings from the UK Millennium Cohort Study. *SSM-population health*. 2018;4:280-290. DOI: [10.1016/j.ssmph.2018.03.002](https://doi.org/10.1016/j.ssmph.2018.03.002)
28. McGee R, Williams S, Silva PA. An evaluation of the Malaise Inventory. *Journal of Psychosomatic Research*. 1986;30(2):147-152. DOI: [10.1016/0022-3999\(86\)90044-9](https://doi.org/10.1016/0022-3999(86)90044-9)
29. Condon JT, Corkindale CJ. The assessment of parent-to-infant attachment: development of a self-report questionnaire instrument. *Journal of Reproductive and Infant Psychology*. 1998;16(1):57-76. DOI: [10.1016/0022-3999\(86\)90044-9](https://doi.org/10.1016/0022-3999(86)90044-9)
30. Curran EA, Cryan JF, Kenny LC, Dinan TG, Kearney PM, Khashan AS. Obstetrical mode of delivery and childhood behavior and psychological development in a British cohort. *Journal of autism and developmental disorders*. 2016;46(2):603-614. DOI: [10.1007/s10803-015-2616-1](https://doi.org/10.1007/s10803-015-2616-1)

31. Su CC, Tsai CY, Tsai TH, Tsai IJ. Incidence and risk of attention-deficit hyperactivity disorder in children with amblyopia: A nationwide cohort study. *Clinical & experimental ophthalmology*. 2019;47(2):259-264. DOI: [10.1111/ceo.13465](https://doi.org/10.1111/ceo.13465)
32. Hancock KJ, Brennan-Jones CG, Vithiatharan R, et al. Mental health problems among 4–17-year-olds with hearing problems: Results from a nationally representative study. *Hearing, Balance and Communication*. 2017;15(3):145-155.  
[10.1080/21695717.2017.1325094](https://doi.org/10.1080/21695717.2017.1325094)
33. DeCarlo DK, Swanson M, McGwin G, Visscher K, Owsley C. ADHD and vision problems in the National Survey of Children's Health. *Optometry and vision science: official publication of the American Academy of Optometry*. 2016;93(5):459.  
DOI: [10.1097/OPX.0000000000000823](https://doi.org/10.1097/OPX.0000000000000823)
34. Bellato A, Perna J, Ganapathy PS, et al. Association between ADHD and vision problems. A systematic review and meta-analysis. *Molecular Psychiatry*. 2023;28(1):410-422. <https://doi.org/10.1038/s41380-022-01699-0>
35. Franklin MS, Mrazek MD, Anderson CL, et al. Tracking distraction: The relationship between mind-wandering, meta-awareness, and ADHD symptomatology. *Journal of attention disorders*. 2017;21(6):475-486. <https://doi.org/10.1177/1087054714543494>
36. Druker K, Hennessey N, Mazzucchelli T, Beilby J. Elevated attention deficit hyperactivity disorder symptoms in children who stutter. *Journal of fluency disorders*. 2019;59:80-90. DOI: [10.1016/j.jfludis.2018.11.002](https://doi.org/10.1016/j.jfludis.2018.11.002)
37. Brikell I, Ghirardi L, D’Onofrio BM, et al. Familial liability to epilepsy and attention-deficit/hyperactivity disorder: a nationwide cohort study. *Biological psychiatry*. 2018;83(2):173-180. DOI: [10.1016/j.biopsych.2017.08.006](https://doi.org/10.1016/j.biopsych.2017.08.006)

38. Chou I-C, Chang Y-T, Chin Z-N, et al. Correlation between epilepsy and attention deficit hyperactivity disorder: a population-based cohort study. *PLoS One*. 2013;8(3):e57926. DOI: [10.1371/journal.pone.0057926](https://doi.org/10.1371/journal.pone.0057926)
39. Wu Y, Li Y, Zhu J, Long J. Shared genetics and causality underlying epilepsy and attention-deficit hyperactivity disorder. *Psychiatry Research*. 2022;316:114794. DOI: [10.1016/j.psychres.2022.114794](https://doi.org/10.1016/j.psychres.2022.114794)
40. Carpena MX, Bonilla C, Matijasevich A, et al. Sleep-related traits and attention-deficit/hyperactivity disorder comorbidity: Shared genetic risk factors, molecular mechanisms, and causal effects. *The World Journal of Biological Psychiatry*. 2021;22(10):778-791. DOI: [10.1080/15622975.2021.1907719](https://doi.org/10.1080/15622975.2021.1907719)
41. Brainstorm C, Anttila V, Bulik-Sullivan B, et al. Analysis of shared heritability in common disorders of the brain. *Science*. Jun 22 2018;360(6395)doi:10.1126/science.aap8757
42. Allone C, Buono VL, Corallo F, et al. Neuroimaging and cognitive functions in temporal lobe epilepsy: A review of the literature. *Journal of the neurological sciences*. 2017;381:7-15. DOI: [10.1016/j.jns.2017.08.007](https://doi.org/10.1016/j.jns.2017.08.007)
43. Friedman NP, Robbins TW. The role of prefrontal cortex in cognitive control and executive function. *Neuropsychopharmacology*. 2022;47(1):72-89. <https://doi.org/10.1038/s41386-021-01132-0>
44. Sun S, Kuja-Halkola R, Chang Z, Cortese S, Almqvist C, Larsson H. Familial liability to asthma and ADHD: A Swedish national register-based study. *JCPP Advances*. 2021;1(4):e12044. DOI: [10.1002/jcv2.12044](https://doi.org/10.1002/jcv2.12044)
45. Chen M-H, Pan T-L, Hsu J-W, et al. Risk of type 2 diabetes in adolescents and young adults with attention-deficit/hyperactivity disorder: a nationwide longitudinal study. *The Journal of clinical psychiatry*. 2018;79(3):0-0. DOI: [10.4088/JCP.17m11607](https://doi.org/10.4088/JCP.17m11607)

46. Faraone SV, Larsson H. Genetics of attention deficit hyperactivity disorder. *Molecular psychiatry*. 2019;24(4):562-575. DOI: [10.1038/s41380-018-0070-0](https://doi.org/10.1038/s41380-018-0070-0)
47. Pinares-Garcia P, Stratikopoulos M, Zagato A, Loke H, Lee J. Sex: a significant risk factor for neurodevelopmental and neurodegenerative disorders. *Brain sciences*. 2018;8(8):154. DOI: [10.3390/brainsci8080154](https://doi.org/10.3390/brainsci8080154)
48. Assari S, Caldwell CH. Family income at birth and risk of attention deficit hyperactivity disorder at age 15: racial differences. *Children*. 2019;6(1):10. DOI: [10.3390/children6010010](https://doi.org/10.3390/children6010010)
49. Russell G, Ford T, Rosenberg R, Kelly S. The association of attention deficit hyperactivity disorder with socioeconomic disadvantage: alternative explanations and evidence. *Journal of Child Psychology and Psychiatry*. 2014;55(5):436-445. DOI: [10.1111/jcpp.12170](https://doi.org/10.1111/jcpp.12170)
50. Russell AE, Ford T, Russell G. Socioeconomic associations with ADHD: findings from a mediation analysis. *PloS one*. 2015;10(6):e0128248. DOI: [10.1371/journal.pone.0128248](https://doi.org/10.1371/journal.pone.0128248)
51. Andersen CH, Thomsen PH, Nohr EA, Lemcke S. Maternal body mass index before pregnancy as a risk factor for ADHD and autism in children. *European child & adolescent psychiatry*. 2018;27(2):139-148. DOI: [10.1007/s00787-017-1027-6](https://doi.org/10.1007/s00787-017-1027-6)
52. He Y, Chen J, Zhu L-H, Hua L-L, Ke F-F. Maternal smoking during pregnancy and ADHD: results from a systematic review and meta-analysis of prospective cohort studies. *Journal of attention disorders*. 2020;24(12):1637-1647. DOI: [10.1177/1087054717696766](https://doi.org/10.1177/1087054717696766)
53. Rougeaux E, Hope S, Viner RM, Deighton J, Law C, Pearce A. Is mental health competence in childhood associated with health risk behaviors in adolescence? Findings from the UK Millennium Cohort Study. *Journal of Adolescent Health*. 2020;67(5):677-684. doi: [10.1016/j.jadohealth.2020.04.023](https://doi.org/10.1016/j.jadohealth.2020.04.023)



54. Murray AL, Hall HA, Speyer LG, et al. Developmental trajectories of ADHD symptoms in a large population-representative longitudinal study. *Psychological Medicine*. 2021;1-7. DOI: [10.1017/S0033291721000349](https://doi.org/10.1017/S0033291721000349)
55. Chang Z, Lichtenstein P, D’Onofrio BM, et al. Maternal age at childbirth and risk for ADHD in offspring: a population-based cohort study. *International journal of epidemiology*. 2014;43(6):1815-1824. DOI: [10.1093/ije/dyu204](https://doi.org/10.1093/ije/dyu204)
56. Torvik FA, Eilertsen EM, McAdams TA, et al. Mechanisms linking parental educational attainment with child ADHD, depression, and academic problems: a study of extended families in The Norwegian Mother, Father and Child Cohort Study. *Journal of Child Psychology and Psychiatry*. 2020;61(9):1009-1018. DOI: [10.1111/jcpp.13197](https://doi.org/10.1111/jcpp.13197)
57. Hall HA, Speyer LG, Murray AL, Auyeung B. Prenatal maternal infections and children’s neurodevelopment in the UK Millennium Cohort Study: A focus on ASD and ADHD. *Journal of Attention Disorders*. 2021:10870547211015422. DOI: [10.1177/10870547211015422](https://doi.org/10.1177/10870547211015422)
58. Zeng Y, Tang Y, Tang J, et al. Association between the different duration of breastfeeding and attention deficit/hyperactivity disorder in children: A systematic review and meta-analysis. *Nutritional Neuroscience*. 2020;23(10):811-823. DOI: [10.1080/1028415X.2018.1560905](https://doi.org/10.1080/1028415X.2018.1560905)
59. Bendiksen B, Aase H, Diep LM, Svensson E, Friis S, Zeiner P. The associations between pre-and postnatal maternal symptoms of distress and preschooler’s symptoms of ADHD, oppositional defiant disorder, conduct disorder, and anxiety. *Journal of attention disorders*. 2020;24(7):1057-1069. DOI: [10.1177/1087054715616185](https://doi.org/10.1177/1087054715616185)
60. Sagiv SK, Epstein JN, Bellinger DC, Korrick SA. Pre-and postnatal risk factors for ADHD in a nonclinical pediatric population. *Journal of attention disorders*. 2013;17(1):47-57. DOI: [10.1177/1087054711427563](https://doi.org/10.1177/1087054711427563)

61. Storebø OJ, Rasmussen PD, Simonsen E. Association between insecure attachment and ADHD: environmental mediating factors. *Journal of Attention Disorders*. 2016;20(2):187-196. DOI: [10.1177/1087054713501079](https://doi.org/10.1177/1087054713501079)

**Table 1.** Weighted Frequencies of Sample Demographics and Other Environmental Risk Factors. (N=8059)

	<b>High Risk Criteria</b>	<b>High Risk N (%)</b>
<b>SES<sup>a</sup> and demographics</b>		
Sex <sup>47</sup>	Male	4016 (49.8)
Ethnicity <sup>48</sup>	Any ethnicity other than white	968 (12)
Household income <sup>49</sup>	Below 60% median poverty indicator	2502 (31.1)
Household crowding <sup>12</sup>	Fewer rooms than people (excluding bathrooms)	743 (9.2)
Housing Tenure <sup>50</sup>	Social housing or renting from local authority	1825 (22.7)
<b>Prenatal factors</b>		
Maternal pre-pregnancy BMI <sup>b51</sup>	>24.9	2089 (25.9)
Antenatal blood pressure <sup>7</sup>	Mother diagnosed with Pre-eclampsia/ Hypertension	561 (7)
Maternal diabetes diagnosis <sup>7</sup>	Mother has Diabetes diagnosis	159 (2)
Smoking in pregnancy <sup>52, 53</sup>	Mother smoked 1 or more cigarettes during pregnancy	1716 (21.3)
<b>Perinatal factors</b>		
Birthweight <sup>8</sup>	<2.5kg	502 (6.2)
Gestation <sup>8, 54</sup>	<37 weeks gestation (premature)	663 (8.2)
Delivery <sup>7</sup>	Caesarean section	1584 (19.7)
Presentation/lie <sup>7</sup>	Breech presentation/ other abnormal lie	452 (5.6)
<b>Postnatal environmental factors</b>		
Maternal age <sup>55</sup>	≤19 years	671 (8.3)
Number of parents in household <sup>49</sup>	1 parent living in household	1270 (15.8)
Mother's education <sup>56, 57</sup>	Mother educated to <NVQ <sup>c</sup> level 3 (2 A-levels)	4847 (60.1)
Breastfeeding <sup>58</sup>	No breastfeeding	2480 (30.8)
<b>Postnatal maternal wellbeing factors</b>		
Maternal distress <sup>59, 27</sup>	<4 on Rutter Malaise Inventory	358 (4.4)
Maternal mental health diagnosis <sup>60</sup>	Depression/anxiety diagnosis	2008 (24.9)
Maternal attachment <sup>61, 30</sup>	>22 on Condon Maternal Attachment Scale	1870 (23.2)

<sup>a</sup> SES = socioeconomic status, <sup>b</sup> BMI = Body Mass Index, <sup>c</sup> NVQ = National Vocational Qualifications

**Table 2.** Weighted Frequencies of Individual and Clustered Physical Health Conditions, and Cumulative Risk Indices (N=8059)

	N (%)	Number of Conditions N(%)					
		0	1	2	3		
<b>Sensory cluster</b>		5051 (62.7)	2515 (31.2)	493 (6.1)			
Eyesight problems	1802 (22.4)						
Hearing problems	1700 (21.1)						
<b>Atopic cluster</b>		3271 (40.6)	2790 (34.6)	1477 (18.3)	508 (6.3)		
Asthma	1375 (17.1)						
Eczema	3411 (42.3)						
Hay fever	2482 (30.8)						
<b>Neurological cluster</b>		6851 (85)	1093 (13.6)	109 (1.4)	6 (0.1)		
Movement problems	452 (5.6)						
Neurological disorders	41 (0.5)						
Sleeping problems	82 (1)						
Stutter	261 (3.2)						
Fits or Epilepsy	491 (6.1)						
<b>Cardio-metabolic cluster</b>		5983 (74.2)	2052 (25.5)	24 (0.3)			
Diabetes	22 (0.3)						
Obesity age 17	2031 (25.2)						
Cardiovascular disorders	47 (0.6)						
		Number of Environmental Risk Factors N(%)					
<b>Cumulative Risk Indices</b>		0	1	2	3	4	5
Prenatal factors		4336 (53.8)	2983 (37)	682 (8.5)	53 (0.7)	5 (0.1)	
Perinatal factors		5747 (71.3)	1596 (19.8)	556 (6.9)	148 (1.8)	13 (0.2)	
Postnatal environmental factors		2707 (33.6)	2535 (31.5)	1917 (23.8)	701 (8.7)	199 (2.5)	
Postnatal maternal wellbeing factors		4625 (57.4)	2719 (33.7)	622 (7.7)	90 (1.1)		
Socioeconomic status and demographics		2229 (27.7)	3175 (39.4)	1431 (17.8)	901 (11.2)	298 (3.7)	24 (0.3)

**Table 3.** Regression Statistics for all Models with SDQ<sup>a</sup> Score at age 17 as the Outcome Variable.

	<b>R<sup>2</sup></b>	<b>R<sup>2</sup> change</b>	<b>B</b>	<b>β</b>	<b>95% Confidence Interval</b>	
					<b>LL</b>	<b>UL</b>
<b>Model 1</b>	.02	.02				
Constant			2.10**		2.00	2.20
Clusters						
Sensory			0.31**	0.09	0.22	0.41
Neurological			0.50**	0.09	0.36	0.64
Atopic			0.05	0.02	-0.02	0.11
Cardio-metabolic			0.17**	0.04	0.04	0.31
<b>Model 2</b>	.09	.07				
Constant			1.27**		1.15	1.39
Clusters						
Sensory			0.28**	0.08	0.19	0.37
Neurological			0.44**	0.08	0.38	0.64
Atopic			0.04	0.02	-0.03	0.09
Cardio-metabolic			0.04	0.01	-0.08	0.16
CRIs <sup>b</sup>						
Prenatal risk			0.22**	0.07	0.14	0.30
Perinatal risk			0.09*	0.03	0.01	0.16
Mother environment			0.26**	0.12	0.26	0.38
Mother wellbeing			0.18**	0.05	0.15	0.31
SES <sup>c</sup> & demographics			0.40**	0.18	0.31	0.42
<b>Model 3</b>	.15	.06				
Constant			1.34**		1.23	1.46
Clusters						
Sensory			0.28**	0.08	0.17	0.34
Neurological			0.40**	0.08	0.35	0.60
Atopic			0.02	0.01	-0.05	0.07

Cardio-metabolic			0.05	0.01	-0.06	0.17
<b>CRIs</b>						
Prenatal risk			0.19**	0.06	0.11	0.27
Perinatal risk			0.08*	0.03	0.01	0.15
Mother environment			0.23**	0.10	0.24	0.36
Mother wellbeing			0.14**	0.04	0.11	0.27
SES & demographics			0.36**	0.16	0.27	0.38
ADHD <sup>d</sup> medication			4.86**	0.24	4.35	5.27
<b>Model 4</b>	.21	.06				
Constant			0.70**		0.56	0.84
<b>Clusters</b>						
Sensory			0.22**	0.06	0.13	0.31
Neurological			0.32**	0.06	0.20	0.45
Atopic			0.01	0.00	-0.05	0.07
Cardio-metabolic			0.01	0.00	-0.11	0.13
<b>CRIs</b>						
Prenatal risk			0.16**	0.05	0.08	0.24
Perinatal risk			0.07	0.02	-0.01	0.14
Mother environment			0.16**	0.07	0.10	0.22
Mother wellbeing			0.08*	0.02	0.001	0.16
SES & demographics			0.27**	0.12	0.21	0.32
ADHD Medication			4.39**	0.22	3.91	4.87
SDQ score at age three			0.25**	0.25	0.22	0.27

\* p < .05, \*\* p < .001

<sup>a</sup> SDQ = Strengths and Difficulties Questionnaire, <sup>b</sup> CRIs = Cumulative Risk Indices, <sup>c</sup> SES = Socioeconomic Status, <sup>d</sup> ADHD = Attention Deficit/Hyperactivity Disorder




Figure 1. Weighted Mean SDQ Scores for Cumulative Numbers of Conditions in Each Physical Health Cluster.