



Reporting of Observational Studies Explicitly Aiming to Emulate Randomized Trials A Systematic Review

Harrison J. Hansford, BSc(Hons); Aidan G. Cashin, PhD; Matthew D. Jones, PhD; Sonja A. Swanson, ScD; Nazrul Islam, MD, PhD; Susan R. G. Douglas, BExPhys; Rodrigo R. N. Rizzo, PhD; Jack J. Devonshire, BSc(Hons); Sam A. Williams, BSc(Hons); Issa J. Dahabreh, MD, ScD; Barbra A. Dickerman, PhD; Matthias Egger, MD, MSc; Xabier Garcia-Albeniz, MD, PhD; Robert M. Golub, MD; Sara Lodi, PhD; Margarita Moreno-Betancur, PhD; Sallie-Anne Pearson, PhD; Sebastian Schneeweiss, MD, ScD; Jonathan A. C. Sterne, PhD; Melissa K. Sharp, PhD; Elizabeth A. Stuart, PhD; Miguel A. Hernán, MD, DrPh; Hopin Lee, PhD; James H. McAuley, PhD

Abstract

IMPORTANCE Observational (nonexperimental) studies that aim to emulate a randomized trial (ie, the target trial) are increasingly informing medical and policy decision-making, but it is unclear how these studies are reported in the literature. Consistent reporting is essential for quality appraisal, evidence synthesis, and translation of evidence to policy and practice.

OBJECTIVE To assess the reporting of observational studies that explicitly aimed to emulate a target trial.

EVIDENCE REVIEW We searched Medline, Embase, PsycINFO, and Web of Science for observational studies published between March 2012 and October 2022 that explicitly aimed to emulate a target trial of a health or medical intervention. Two reviewers double-screened and -extracted data on study characteristics, key predefined components of the target trial protocol and its emulation (eligibility criteria, treatment strategies, treatment assignment, outcome[s], follow-up, causal contrast[s], and analysis plan), and other items related to the target trial emulation.

FINDINGS A total of 200 studies that explicitly aimed to emulate a target trial were included. These studies included 26 subfields of medicine, and 168 (84%) were published from January 2020 to October 2022. The aim to emulate a target trial was explicit in 70 study titles (35%). Forty-three studies (22%) reported use of a published reporting guideline (eg, Strengthening the Reporting of Observational Studies in Epidemiology). Eighty-five studies (43%) did not describe all key items of how the target trial was emulated and 113 (57%) did not describe the protocol of the target trial and its emulation.

CONCLUSIONS AND RELEVANCE In this systematic review of 200 studies that explicitly aimed to emulate a target trial, reporting of how the target trial was emulated was inconsistent. A reporting guideline for studies explicitly aiming to emulate a target trial may improve the reporting of the target trial protocols and other aspects of these emulation attempts.

JAMA Network Open. 2023;6(9):e2336023.

Corrected on October 16, 2023. doi:10.1001/jamanetworkopen.2023.36023

Introduction

Analyses of observational (nonexperimental) data can be used to estimate the causal effect of interventions when randomized clinical trials are unavailable or infeasible. Bias in observational analyses may be limited by conceptualizing them as attempts to emulate target trials, ie, hypothetical randomized trials that would answer causal questions of interest.¹⁻³ Hernán and Robins⁴ have outlined a framework for this approach, which involves first specifying the protocol of the target trial

Open Access. This is an open access article distributed under the terms of the CC-BY License.

Key Points

Question How are studies that explicitly aim to emulate a target trial reported?

Findings In this systematic review of 200 studies that explicitly aimed to emulate a target trial, reporting was inconsistent, and studies often did not report all necessary information related to the emulation of the target trial.

Meaning Inconsistent reporting of studies that explicitly aim to emulate a target trial may impair the appraisal, synthesis, and implementation of study findings.

+ Supplemental content

Author affiliations and article information are listed at the end of this article.

and then emulating the trial as closely as possible using observational data.^{4,5} The target trial framework may help reduce common biases in observational analyses and enhance transparency regarding design and analytic decisions. Moreover, it facilitates the interpretation of effect estimates and promotes meaningful discourse concerning potentially discrepant findings observed across studies.

Since at least the 1950s, the notion of observational analyses as attempts to approximate the goals of randomized clinical trials has underpinned many comparative studies in health, medicine, and related fields.⁶⁻¹⁰ The target trial emulation framework, introduced by Hernán and Robins⁴ in 2016, provided a template for reporting and conducting studies that aim to emulate target trials. The framework outlines items to be reported in the protocol of a target trial and its emulation, including: eligibility criteria, treatment strategies, assignment procedures, follow-up period, outcome(s), causal contrast(s), and analysis plan. Since the introduction of the framework, several articles have been published to assist researchers in conducting these studies and educating clinicians and other end users to interpret their findings.^{3,5,11-14} However, there is limited understanding of how researchers have implemented the target trial framework when reporting observational analyses with the explicit aim to emulate a target trial.

This review aimed to (1) describe how studies that explicitly aimed to emulate a target trial were reported and (2) examine whether these used published reporting guidelines. The findings of this review will be used to inform the development of a reporting guideline for studies explicitly aiming to emulate a target trial.¹⁵

Methods

This review is reported according to the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) reporting guideline.¹⁶ The protocol was prospectively registered on the Open Science Framework (OSF).^{17,18} Protocol deviations are reported in eAppendix 1 in [Supplement 1](#).

Searches

We searched 4 electronic databases of published literature from March 13, 2012, to October 20, 2022, including Medline, Embase, PsycINFO, and Web of Science. Our search included terms such as *emulat* trial*, *target trial*, *target trial emulat**, *real world data*, and *causal inference*. Our complete search strategy is provided in eAppendix 2 in [Supplement 1](#). To supplement the search strategy, we used citationchaser¹⁹ to conduct forward citation tracking of 5 selected seminal papers describing the target trial emulation framework.^{3,4,8,20,21} We also included papers known to the authorship team.

Eligibility Criteria

We included observational studies that explicitly aimed to emulate a target trial of a medical intervention; eAppendix 3 in [Supplement 1](#) provides all terms deemed sufficient for an explicit target trial emulation. We restricted our inclusion of studies published from March 13, 2012 (10 years prior to registration of our protocol¹⁸ to capture recent trends in reporting) to October 20, 2022. We excluded studies that did not investigate a medical intervention; did not include human participants; were not written in English; only described the protocol of a study emulating a target trial, ie, a protocol of a planned study without results; or for which the full text was unavailable.

Record Management and Screening

We de-duplicated all records identified through searches in Endnote version 20 and imported into Excel version 2206 (Microsoft Corp). In duplicate, reviewers (H.J.H., A.G.C., M.D.J., and S.R.G.D.) independently performed screening of identified records for eligibility at the level of title and abstract and full text. Disagreements were resolved through discussion.

Data Extraction

Data were extracted in duplicate (H.J.H., M.D.J., S.R.G.D., J.J.D., S.A.W., R.R.N.R., A.G.C. performed this task independently) and compiled into a standardized spreadsheet piloted with 3 included studies. Disagreements were resolved by the lead author (H.J.H.) or through discussion. We did not blind reviewers to the journal article or study authors.

Data Items

We extracted information about the (1) characteristics of the included studies, (2) key protocol components that characterize the target trial approach, and (3) further items that may be important to report in studies emulating a target trial. The complete data extraction spreadsheet and code used are available on OSF.¹⁸

Characteristics of the included studies were year of publication, subfield of medicine defined based upon included population, data source (prospective cohort, electronic health records, claims data, registry, randomized clinical trial, or linked data, ie, where data sources were combined), sample size (unique individuals included and analyzed, rather than simulated or duplicated persons, such as in sequential trial designs), primary outcome, and type of treatment strategy being compared. *Treatment strategy* refers to any health care intervention including treatments, preventative interventions, and no change to current treatments, remaining consistent with the language used by Hernán and Robins.⁴ If a study investigated prevention of a given outcome in healthy individuals, the subfield of medicine was designated based on the outcome investigated. Each treatment strategy included in an article (eg, ≥ 2) was counted separately. We classified treatment strategies defined by the authors as no treatment or usual care as no change to current treatment approach(es).

We extracted whether each study reported the eligibility criteria, treatment strategies, assignment procedures, follow-up period, outcome(s), causal contrast(s), and analysis plan of the emulation of the target trial. These items and their definitions were informed by the target trial framework from Hernán and Robins.⁴ We considered a study to have specified how the target trial was emulated if all the previously listed protocol items were reported; these items are operationalized in eAppendix 4 in Supplement 1. We extracted whether the protocol of the target trial or its emulation were presented in a table or in text, with table being prioritized if reported in both table and text format. We also stated whether the protocol of the target trial and how it was emulated were reported. We extracted whether the study reported a baseline in the target trial emulation where eligibility criteria, start of follow-up, and treatment assignment were aligned.

Further details of specific protocol components that may be important to report in studies emulating a target trial were chosen based on expert knowledge and recommendations from methodological papers on the target trial emulation framework.²⁻⁴ These included:

- Treatment strategies: type of treatment strategy (eg, pharmacological, surgical; all studies are expected to include 2 or more treatment strategies), aspects of treatment strategies described (type of treatment, frequency, dose, and duration of treatment strategy).
- Analysis plan: method(s) used to emulate randomization, description and selection of potential confounding variables, statistical and causal assumptions that relate to analyses, sensitivity analyses.
- Other: study registration, rationale for the target trial emulation, reporting guideline used (referred to as a guideline hereafter). We only included a guideline if it was referenced as guiding the reporting of a study.

We deemed a study to report the assumptions underlying their analyses only when the assumption(s) were described in the text or in a cited reference. When authors reported that no residual confounding was assumed, we took this as equivalent to reporting an assumption of conditional exchangeability. We did not regard practices that may assess a causal assumption (eg, truncation of weights to satisfy the assumption of positivity) as reporting the assumption. We did not

assess the appropriateness of authors' reported assumptions. Items we extracted that were not included in commonly used guidelines are listed in eAppendix 5 in Supplement 1.

Data Analysis

We cleaned and analyzed data in R version 4.2.0 (R Project for Statistical Computing) using tableone, openxlsx, tidyverse, and readxl packages for data management and visualization. We summarized categorical variables using counts and percentages. Continuous variables were summarized using mean and SD or median and interquartile range. Post hoc, we assessed the reporting of how the target trial was emulated stratified by whether a guideline was used.

Results

We retrieved 3133 unique records, of which 200 were included in the review (Figure 1).^{2,22-220} All reasons for excluding records after full-text review are given in eAppendix 6 in Supplement 1.

Figure 1. Study Flow Diagram





Figure 2. Number of Explicit Emulations of a Target Trial Included in Review Published per Year



^a Publications until October 2022.

Characteristics of Included Studies

Of the 200 included studies,^{2,22-220} 168 (84%) were published from January 2020 to October 2022 (Figure 2).^{22, 23, 25-46, 48-51, 53, 57, 58, 60-68, 72-77, 79, 80, 82-90, 94-97, 99-113, 115-143, 145-165, 167-179, 181, 183-195, 197-199, 201, 203, 205, 206, 208, 210, 211, 213-217} The included studies spanned 26 fields of medicine, predominately infectious disease (43 [22%]; 27 [14%] on COVID-19),^{26, 27, 30, 35-39, 47, 53-56, 59, 66-68, 77, 82, 90, 94, 97, 99, 100, 108-110, 130, 136, 139-141, 145, 148, 157, 166, 186, 197, 205, 217, 218} cardiology (30 [15%]),^{22, 28, 31, 41, 60, 70, 71, 73, 85, 105, 106, 116, 117, 121, 122, 143, 149-152, 161, 179, 184, 187, 193, 198, 206, 214} and oncology (30 [15%]).^{2, 32, 40, 44-46, 51, 76, 78, 81, 89, 91-93, 95, 103, 104, 129, 132, 138, 153, 168, 173, 174, 178, 182, 190, 192, 204, 211} One hundred and thirty-one studies (66%)^{2, 22, 23, 25, 31, 32, 34-36, 39, 41-47, 51-53, 58, 59, 61-63, 65, 67, 68, 70-77, 80, 81, 85, 86, 89, 90, 92-96, 100-106, 108, 109, 111, 112, 116, 117, 119-131, 133-137, 141, 143, 144, 146-158, 160-162, 164, 167, 168, 170, 171, 173, 177-181, 183-185, 187-189, 193, 195, 199-201, 205, 207, 210, 212, 213, 216, 217, 219, 220} used data from electronic health records, claims databases, or linked data sources. The treatment strategies most frequently investigated were pharmacological (228 of 435 [52%]) and no change to current treatment approach(es) (usual care or noninitiation of study treatment; 82 of 435 [19%]). All extracted characteristics of included studies are displayed in Table 1.

Table 1. Characteristics of 200 Included Studies

Characteristic	Count, No. (%)
Domain	
Infectious diseases	43 (22)
Cardiology	30 (15)
Oncology	30 (15)
Nephrology	14 (7)
Endocrinology	11 (6)
Rheumatology	10 (5)
Internal medicine	9 (5)
Neurology	7 (4)
Psychiatry	6 (3)
Other	40 (20)
Data source	
Electronic health record data	49 (25)
Linked data ^a	46 (23)
Claims data	36 (18)
Registry	32 (16)
Prospective cohort ^b	30 (15)
Randomized clinical trial	6 (3)
Not reported	1 (1)
Sample size, median (IQR)^c	
Sample eligible	11 253 (2157-101 078)
Sample analyzed	9799 (1995-98 718)
Primary outcome	
Death	72 (36)
Major adverse cardiovascular event ^d	19 (10)
Cancer	8 (4)
Other	101 (51)
Treatment strategies compared, No.	
2	187 (94)
3	5 (3)
4	4 (2)
5	2 (1)
6	2 (1)
10	1 (1)

(continued)

Table 1. Characteristics of 200 Included Studies (continued)

Characteristic	Count, No. (%)
Type of treatment strategy, No./total No. (%) ^e	
Pharmacological	228/435 (52)
No change to current treatment	82/435 (19)
Other ^f	61/435 (14)
Surgical	37/435 (9)
Vaccine	19/435 (4)
Medical device	8/435 (2)

^a Data in which 2 or more data sources are combined, eg, a registry is linked to a claims database.

^b Studies could only contribute to 1 data source item; if data collection for a cohort was conducted prospectively, the data source was classified only as a prospective cohort, even if data collection took place in the form of electronic health records or other data source listed.

^c The sample size includes the number of unique participants.

^d As major adverse cardiovascular event is often described heterogeneously, myocardial infarction, stroke, or major adverse cardiovascular event as defined by the authors were included; where the primary outcome was death, even if from cardiovascular events, the outcome was classified as death.

^e A given study may contribute 2 or more treatment strategies, which may be different, ie, 2 pharmacological treatment strategies compared with 2 no change to current treatment strategies.

^f The other category includes health care consultations, health care programs, organ transplants, and other interventions that would not fall under the other categories listed.

Characteristics of Target Trials and How They Were Emulated

One-hundred and fifteen studies (58%)^{2, 22, 23, 25, 26, 30-32, 34, 35, 42, 45-51, 54-57, 59, 61, 64-66, 69-72, 74-78, 85, 87-91, 93, 95, 96, 98, 100-104, 106, 116, 127, 128, 130, 132-135, 137-139, 141-143, 148, 149, 151, 152, 154, 156-158, 160-162, 167, 168, 172, 174, 176-180, 182-184, 187-192, 195, 196, 198-201, 203-206, 208-211, 213-216, 219, 220} completely reported how the target trial protocol was emulated. Eighty-seven studies (44%)^{2, 24, 25, 28, 30, 32, 34, 42, 46, 51, 52, 55-57, 59, 61, 62, 64-66, 70-72, 76-79, 82, 85, 87-89, 91-94, 98, 100, 103, 104, 107, 113, 115-117, 119, 120, 122, 125, 126, 130, 133-135, 142, 143, 147, 149, 150, 152, 158, 162, 165, 168, 170, 172, 176, 179, 185, 187, 188, 190-192, 195, 202, 203, 209, 212-214, 216, 219, 220} provided both the protocol of the target trial and described how it was emulated (**Table 2**). The following items of the emulation were frequently reported: eligibility criteria (193 [97%]), treatment strategies (191 [96%]), assignment procedures (173 [87%]), primary outcome (196 [98%]), the follow-up period (186 [93%]), a causal contrast (146 [73%]), and an analysis plan (194 [97%]) (Table 2).

Reporting of Further Items That Relate to the Target Trial Emulation

Seventy studies (35%)^{23-25, 29, 30, 32, 34, 37, 43, 48, 50, 51, 55-57, 59, 61, 62, 65, 66, 70, 71, 74-76, 82, 86, 96, 98, 100, 103, 107, 109, 110, 113, 116, 126, 129, 130, 134, 135, 139, 143, 145-147, 149, 153, 154, 158, 159, 163, 165, 167, 168, 171, 172, 174, 175, 177, 179, 189, 199, 203, 206-208, 213, 216, 218} reported in the title that the study aimed to emulate a target trial; 180 (90%)^{2, 22-27, 30-46, 48-84, 86-89, 91-97, 99-104, 106, 107, 109-113, 115-119, 121-128, 130-144, 146-154, 156-158, 160-162, 164-171, 173, 174, 176-195, 197-214, 216-220} did so in the Methods section. Twenty studies (10%)^{24, 25, 34, 35, 43, 52, 68, 80, 84, 86, 117, 131, 133, 147, 156, 161, 169, 171, 192, 216} reported the study was prospectively registered, 16 of these 20 (80%)^{24, 25, 34, 43, 52, 68, 80, 86, 117, 131, 133, 156, 161, 169, 171, 216} also provided information on how to access the registration. One hundred and twenty-six studies (63%)^{2, 22-30, 34, 35, 37, 39, 40, 44-46, 48-50, 52, 54, 57, 59, 60, 62, 64-76, 78, 79, 82, 83, 86-89, 91-93, 99, 102-104, 108, 109, 111-115, 117, 120-127, 130, 133-135, 138, 139, 141, 142, 145-147, 149-153, 155, 156, 158, 159, 161, 162, 164, 168, 169, 171-173, 175, 176, 179, 180, 182, 184, 185, 187, 190-192, 194, 195, 197-201, 205, 208-211, 213, 215-217} specified whether a randomized clinical trial could be feasibly conducted; 61 (31%)^{28, 29, 34, 39, 45, 48, 49, 54, 57, 62, 66-69, 73, 75, 79, 83, 86-88, 91, 99, 102-104, 108, 109, 112, 115, 117, 121-123, 125, 126, 138, 139, 146, 152, 153, 156, 158, 162, 171, 184, 185, 187, 192, 198, 199, 205, 210, 211, 213, 215, 216} stated that the randomized clinical trial was possible. Of the

studies that stated a randomized clinical trial was possible, uncertainty in the generalizability of available trial findings was the most common reason for the target trial emulation (22 of 61 [36%]).^{28, 45, 52, 57, 68, 85, 86, 103, 114, 115, 117, 122, 123, 152, 158, 184, 187, 192, 198, 210, 215} Forty-three studies (22%)^{2, 27, 39, 45, 48, 53, 61, 62, 67, 75, 80, 87, 99, 101, 102, 104, 107, 111, 112, 114, 117, 118, 122, 123, 128, 130, 135, 144, 151, 152, 155, 158-161, 168, 183, 189, 194, 195, 208, 210, 215} reported using a guideline, most commonly (29 of 43 [67%]) the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guideline.^{221, 222} There were no qualitative differences between the reporting of the target trial emulation when studies were stratified by guideline use (eAppendix 7 in Supplement 1).

Most studies (187 [94%])^{2, 22-29, 31-40, 42-51, 53-61, 63-79, 81-83, 87, 89-126, 128-174, 176-184, 186-192, 194-220} reported the set of variables that authors had decided to adjust for (eg, because they were potential confounders) in analyses, and 77 (39%)^{22, 24-26, 28, 35, 41, 44, 47-52, 58, 61, 68-70, 77, 80, 81, 84, 87, 95, 99, 100, 104, 105, 107, 110, 111, 114, 124, 126, 130, 135, 138, 140, 141, 147-149, 151, 155, 156, 158-162, 165, 166, 168, 170, 171, 173, 175, 182, 183, 185, 186, 197, 199, 205, 208, 212, 214, 216-218} reported how these variables were selected. One hundred and thirty-one studies (66%)^{2, 25, 26, 28, 31, 34, 36, 41-44, 46-49, 52, 54, 56-59, 61, 64-66, 68-70, 72, 73, 75-78, 80, 82-88, 90, 92-95, 99-101, 105-107, 109, 112-117, 119, 120, 123, 125-131, 133-138, 141-144, 146, 152-154, 157, 158, 160-166, 168-174, 177, 179-181, 187-189, 192, 194-196, 198-201, 204-208, 210-220} reported conducting a sensitivity analysis for statistical or causal assumptions; the most frequent (42 of 131 [32%]) was the use of a different approach to confounding adjustment (eg, using weighting rather than outcome regression). One hundred and fifty-eight studies (79%)^{2, 22, 25-30, 32, 36, 38-41, 44-52, 54-59, 61, 63-72, 75-85, 87-90, 93, 94, 97, 99-101, 103, 105-109, 111-114, 116-123, 126-130, 132-147, 149, 150, 152, 153, 155, 156, 158-163, 165-177, 180-183, 186-192, 194, 195, 197-208, 210-216, 219} reported that causal interpretation rests on the assumption that the comparison groups were comparable (ie, exchangeable) given the variables included in the analysis (Table 3). Twenty-six studies (13%)^{29, 42, 48, 49, 56, 70, 71, 82, 85, 100, 119, 129, 135, 153, 162, 163, 165, 170, 172, 175, 187, 194, 200, 203, 208, 213} reported reliance on more than 1 causal assumption.

Discussion

This systematic review summarized items reported in observational studies that explicitly aimed to emulate a target trial. We included 200 studies^{2, 22-220} published from 2013 to 2022, the majority of which (168 [84%]) were published between January 2020 and October 2022. The studies spanned 26 fields of medicine and mostly used sources of data that were routinely collected, such as electronic health records, health insurance claims data, or these data linked with other data sources. While the publication of studies explicitly aiming to emulate a target trial is increasing, only 58% of included studies completely reported how the target trial protocol was emulated.

Our finding that studies aiming to emulate a target trial inconsistently reported the emulation of the target trial is similar with results of previous systematic reviews of observational studies that did not explicitly aim to emulate a target trial.²²³ Nguyen et al²²³ systematically reviewed the risk of bias in observational studies investigating the effectiveness of interventions using the ROBINS-I tool,²²⁴ a risk of bias tool informed by the target trial framework. The authors found that only 3% of these observational studies (2 of 77) completely specified all items of the protocol of the (implicit or explicit) target trial. A much larger proportion of our sample of studies that explicitly aimed to emulate a target trial reported how the target trial was emulated; however, many were still incompletely reported. It appears the guidance from Hernán and Robins⁴ and previous work²²⁵⁻²²⁸ has been used inconsistently or perhaps misinterpreted, leaving key elements of the target trial and its emulation unreported.³

Our review shows there has been an increase in the publication of studies that explicitly aim to emulate a target trial. This trend could be attributed to the growing influence of such studies in shaping policy and regulatory decisions.²²⁹⁻²³² For example, in mid-2022, the UK National Institute of Health and Care Excellence released “Real-World Evidence Framework,”²²⁹ which emphasizes the importance of using the framework of a target trial when estimating treatment effects for regulatory decision-making using observational data.²²⁹ Considering the emerging role of studies explicitly

Table 2. Characteristics of Target Trials and How They Were Emulated

Characteristic	Count, No. (%)
How the protocol of the emulated target trial was reported	
Not fully described	85 (42)
Only in text	59 (30)
Table	56 (28)
Both target trial protocol and its emulation described explicitly as such	87 (44)
Description of how the target trial was emulated ^a	
Eligibility criteria	193 (97)
Treatment strategies	191 (96)
Assignment procedures	173 (87)
Outcome(s)	196 (98)
Follow-up	186 (93)
Causal contrast(s)	146 (73)
Analysis plan	194 (97)
Specification of time zero (ie, baseline)	165 (83)

^a Operational definitions of target trial protocol items are described in eAppendix 4 in Supplement 1.

Table 3. Reporting of Further Items That Relate to the Target Trial Emulation

Item	Count, No. (%)
Where aim to emulate a target trial was described ^a	
Title	70 (35)
Abstract	148 (74)
Introduction	119 (60)
Methods	180 (90)
Results	55 (28)
Discussion	142 (71)
Study prospectively registered	20 (10)
Reason given why a randomized clinical trial could not be conducted	
Not reported	74 (37)
NA, trial possible	61 (31)
Unethical	16 (8)
Long-term follow-up	7 (4)
Rare outcomes	7 (4)
Too costly	5 (3)
Not timely	5 (3)
Other	25 (13)
When randomized clinical trial was reported as being possible, primary reason given for emulating a target trial, No./total No. (%)	
Generalizability of available trial findings	22/61 (36)
Replicate published trial	14/61 (23)
Trial ongoing	8/61 (13)
Comparative effectiveness not previously investigated	5/61 (8)
Previous conflicting results reported	3/61 (5)
Other	26/61 (43)
Data source cited	125 (63)
Reporting guideline reported	43 (22)
Reporting guideline used, No./total No. (%)	
STROBE	29/43 (67)
ISPOR Good Research Practices for Comparative Effectiveness Research ^b	5/43 (12)
RECORD	4/43 (9)
Nature Research Reporting Summary	3/43 (7)
RECORD-PE	2/43 (5)
TRIPOD	1/43 (2)
Aspects of treatment strategies described ^a	
Type	417 (96)
Dose	83 (19)
Duration	57 (13)
Frequency	54 (12)
None	18 (4)
Other	13 (3)
Variables adjusted in analyses listed	187 (94)
Potential unmeasured confounders listed	73 (37)
Method for selection of variables adjusted for described	77 (39)
Analytic and causal assumptions stated ^a	
Exchangeability given selected confounders	158 (79)
Positivity	27 (14)
Consistency	13 (7)
Statistical assumptions	24 (12)
Other	4 (2)
None	35 (18)

(continued)

Table 3. Reporting of Further Items That Relate to the Target Trial Emulation (continued)

Item	Count, No. (%)
Sensitivity analyses attempting to assess robustness to analytic or causal assumption(s) violations given	131 (66)
Sensitivity analyses as reported by authors, No./total No. (%)	
Different approach to confounding adjustment	42/131 (32)
Negative control	23/131 (18)
Additional adjustment for confounding	19/131 (15)
E-value	15/131 (11)
Different censoring procedure	10/131 (8)
Different approach to handling missing data	4/131 (3)
Other	24/131 (18)
Table describing baseline characteristics of groups presented	171 (86)

Abbreviations: ISPOR, International Society of Pharmacoeconomics and Outcomes Research; NA, not applicable; RECORD, Reporting of Studies Conducted Using Observational Routinely Collected Health Data; RECORD-PE, Reporting of Studies Conducted Using Observational Routinely Collected Health Data-Statement for Pharmacoepidemiology; STROBE, Strengthening the Reporting of Observational Studies in Epidemiology; TRIPOD, Transparent Reporting of a Multivariable Prediction Model for Individual Prognosis or Diagnosis.

^a Total exceeds 100% as multiple characteristics could be included in a single study.

^b The ISPOR Good Research Practices for Comparative Effectiveness Research are guidelines for the conduct of comparative effectiveness studies, not a reporting guideline, however, were commonly cited as being used for reporting, therefore have been included.

emulating a target trial within the health care decision-making framework, it is critical these studies are consistently and transparently reported. Once a target trial is emulated, unmeasured confounding may be a primary concern with observational analyses informing decision-making.^{229,233} We found that only 73 studies (37%) reported potential unmeasured confounders. It is unlikely all confounders would be measured in a given analysis, therefore the robustness of findings from a target trial emulation may be better assessed if potentially important unmeasured confounders are reported.

Guidelines have been developed to address inconsistent reporting,²³⁴ and if actively implemented, can improve reporting consistency and completeness.²³⁵⁻²³⁷ None of the included studies identified specific reporting guidance for studies that aimed to explicitly emulate a target trial, and the authors are not aware of any guidance for studies emulating a target trial published or under development,²³⁸ suggesting no formal guidance has been published. Twenty-two percent of studies cited general (eg, STROBE)^{221,222} and potentially inappropriate guidelines (eg, Transparent Reporting of a Multivariable Prediction Model for Individual Prognosis or Diagnosis).²³⁹ The use of guidelines was comparatively lower than seen in similar reviews of other types of observational studies, in which observed rates of guideline use ranged from 46% (67 of 147)²⁴⁰ to 47% (68 of 88).²⁴¹ The lower use of guidelines observed in our review may reflect authors' uncertainty on the most appropriate guideline when reporting a study that used the target trial framework.

Implications

Despite the growing number of studies using the target trial framework, reporting was inconsistent. Consistent and transparent reports are important for these studies given their emerging role in decision-making. For example, critical appraisal²²⁴ of the quality and robustness of findings from a study emulating a target trial is impaired when such analyses are poorly reported, leaving readers

unable to understand the quality and conduct of the emulation. Similarly, findings of studies emulating a target trial are frequently compared with those of randomized clinical trials.^{2,86,242-245} Differences in effect estimates between target trial emulations and randomized clinical trials may arise due to various factors.²⁴⁶ Transparent reporting of the target trial protocol and how it was emulated may aid in understanding these differences and optimize the usefulness of these studies for decision-making.

No established, consensus-based¹⁵ guidelines are available to support authors reporting studies emulating a target trial. Commonly used guidelines (eg, STROBE)^{221,222} do not include items that relate to the protocol of the target trial⁴ or key items of the target trial emulation (causal contrast and items that relate to defining time-zero). Reporting of these items was not improved when authors followed guidelines such as STROBE (eAppendix 7 in [Supplement 1](#)). A new guideline for studies that explicitly aim to emulate a target trial is needed to provide detailed recommendations for the minimum set of items to be reported for these studies. Improved reporting of studies emulating a target trial may facilitate peer review by helping to ensure publications are complete, accurate, transparent, and reproducible. Improved reporting could also facilitate scientific discourse, support decision-making, reduce research waste, and ultimately improve health care.^{247,248}

Strengths and Limitations

We used a sensitive search strategy to ensure all relevant studies were captured and followed recommended systematic review methods,²⁴⁹ including screening studies and extracting data in duplicate. We prospectively registered this systematic review¹⁸ and reported the findings in line with the PRISMA 2020 reporting guideline.¹⁶

This study has several limitations. First, we only included studies that explicitly stated that they aimed to emulate a target trial; therefore, our findings may present a more positive view of reporting practices compared with all observational analyses comparing interventions.^{250,251} Using the target trial framework is neither necessary nor sufficient for obtaining valid causal effect estimates from observational analyses; however, the framework may guide the implementation of sound principles of causal inference and study design. Second, we prespecified the reporting items to be extracted based on published recommendations for the specification of the target trial protocol and its emulation.⁴ Therefore, our ratings for these items may be skewed toward a particular way of reporting studies explicitly emulating a target trial. Third, we did not assess the appropriateness of the methods of included studies, only their reporting.

Conclusions

In this systematic review, reporting of studies that explicitly emulate a target trial was inconsistent, with several opportunities to improve the reporting of key items. A guideline expanding on the current recommendations may facilitate consistent and transparent reporting, improving the appraisal, synthesis, and implementation of study findings in clinical practice and health policy.

ARTICLE INFORMATION

Accepted for Publication: August 22, 2023.

Published: September 27, 2023. doi:10.1001/jamanetworkopen.2023.36023

Correction: This article was corrected on October 16, 2023, to fix the degrees listed for Dr Egger in the byline.

Open Access: This is an open access article distributed under the terms of the [CC-BY License](#). © 2023 Hansford HJ et al. *JAMA Network Open*.

Corresponding Author: Harrison J. Hansford, BSc(Hons), School of Health Sciences, Faculty of Medicine and Health, UNSW Sydney, Wallace Wurth Building (C27), 2052 Sydney, Australia (h.hansford@unsw.edu.au).

Author Affiliations: School of Health Sciences, Faculty of Medicine and Health, UNSW Sydney, Sydney, Australia

(Hansford, Cashin, Jones, Douglas, Rizzo, McAuley); Centre for Pain IMPACT, Neuroscience Research Australia, Sydney, Australia (Hansford, Cashin, Jones, Rizzo, Devonshire, Williams, McAuley); Department of Epidemiology, University of Pittsburgh, Pittsburgh, Pennsylvania (Swanson); Oxford Population Health, Big Data Institute, University of Oxford, Oxford, United Kingdom (Islam); Faculty of Medicine, University of Southampton, Southampton, United Kingdom (Islam); CAUSALab, Harvard T.H. Chan School of Public Health, Boston, Massachusetts (Swanson, Dahabreh, Dickerman, Lodi, Hernán); Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts (Swanson, Dahabreh, Dickerman, Garcia-Albeniz, Hernán); Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, Massachusetts (Dahabreh, Hernán); Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland (Egger); Centre for Infectious Disease Epidemiology and Research, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa (Egger); Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, United Kingdom (Egger); RTI Health Solutions, Barcelona, Spain (Garcia-Albeniz); Department of Medicine, Northwestern University Feinberg School of Medicine, Chicago, Illinois (Golub); Department of Biostatistics, Boston University School of Public Health, Boston, Massachusetts (Lodi); Clinical Epidemiology & Biostatistics Unit, Murdoch Children's Research Institute, Royal Children's Hospital, Parkville, Victoria, Australia (Moreno-Betancur); Department of Paediatrics, The University of Melbourne, Parkville, Victoria, Australia (Moreno-Betancur); School of Population Health, Faculty of Medicine and Health, UNSW Sydney, New South Wales, Australia (Pearson); Division of Pharmacoepidemiology, Department of Medicine, Brigham & Women's Hospital, Harvard Medical School, Boston, Massachusetts (Schneeweiss); Department of Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, United Kingdom (Sterne); NIHR Bristol Biomedical Research Centre, Bristol, United Kingdom (Sterne); Health Data Research UK South-West, Bristol, United Kingdom (Sterne); Department of Public Health and Epidemiology, RCSI University of Medicine and Health Sciences, Dublin, Ireland (Sharp); Department of Biostatistics, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland (Stuart); University of Exeter Medical School, Exeter, United Kingdom (Lee); EMEA Methods & Evidence Generation, IQVIA, London, United Kingdom (Lee).

Author Contributions: Mr Hansford and Dr McAuley had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Hansford, Cashin, Jones, Swanson, Islam, Golub, Schneeweiss, Sterne, Stuart, Hernan, Lee, McAuley.

Acquisition, analysis, or interpretation of data: Hansford, Cashin, Jones, Swanson, Islam, Douglas, Rizzo, Devonshire, Williams, Dahabreh, Dickerman, Egger, Garcia-Albeniz, Lodi, Moreno Betancur, Pearson, Sharp, Lee, McAuley.

Drafting of the manuscript: Hansford, Cashin, Egger, Golub, Moreno Betancur, McAuley.

Critical review of the manuscript for important intellectual content: All authors.

Statistical analysis: Hansford, Schneeweiss, Stuart, McAuley.

Obtained funding: Egger.

Administrative, technical, or material support: Islam, Williams, Lee, McAuley.

Supervision: Cashin, Jones, Swanson, Schneeweiss, Hernan, Lee, McAuley.

Conflict of Interest Disclosures: Mr Hansford reported receiving grants from National Health and Medical Research Council (NHMRC) and Neuroscience Research Australia during the conduct of the study. Dr Dahabreh reported receiving grants to the institution from Sanofi and consulting fees from Moderna outside the submitted work. Dr Dickerman reported receiving grants from the National Institutes of Health (NIH) during the conduct of the study. Prof Egger reported receiving grants from Swiss National Science Foundation during the conduct of the study. Dr Garcia-Albeniz reported being employed by RTI Health Solutions. Dr Schneeweiss reported receiving consulting fees from Aetion, Inc. and grants from UCB Pharma, Takeda, and Boehringer Ingelheim outside the submitted work. Dr Stuart reported receiving grants from the NIH during the conduct of the study. Dr Hernán reported receiving grants from NIH during the conduct of the study and receiving personal fees from ProPublica, Cytel, and ADIA Lab outside the submitted work. No other disclosures were reported.

Funding/Support: There is no specific funding for this study. Mr Hansford was supported by an Australian NHMRC Postgraduate Scholarship and a PhD Top-Up Scholarship from Neuroscience Research Australia and was a Neuroscience Research Australia PhD Pearl sponsored by Mrs Sandra Salteri. Dr Cashin was supported by an Australian NHMRC Investigator Grant (identification No. 2010088). Dr Moreno Betancur was supported by an Australian NHMRC Investigator Grant (identification No. 2009572). Dr McAuley was supported by an Australian NHMRC Investigator Grant (identification No. 2010128). Dr Dickerman was supported by a grant from the NIH (grant No. R00 CA248335). Prof Egger was supported by grants from the NIH (grants Nos. R01 AI152772-01 and 5U01-AI069924-05) and the Swiss National Science Foundation (32FP30-174281). Dr Islam was supported by grants from the National Institute for Health and Care Research (grant No. HDRUK2022.0313) and the UK Office

for National Statistics (No. 2002563). Dr Hernán was supported by a grant from the NIH (grant No. R37 AI102634).

Dr Lodi was supported by the Providence/Boston Center for AIDS Research (grant No. P30AI042853).

Role of the Funder/Sponsor: The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Data Sharing Statement: See Supplement 2.

Additional Contributions: We acknowledge Nia Roberts, MSc, FD, outreach librarian and information specialist at the University of Oxford for assistance designing the literature search. She was not compensated for her time outside of her salary.

REFERENCES

1. Kuehne F, Arvandi M, Hess LM, et al. Causal analyses with target trial emulation for real-world evidence removed large self-inflicted biases: systematic bias assessment of ovarian cancer treatment effectiveness. *J Clin Epidemiol*. 2022;152:269-280. doi:10.1016/j.jclinepi.2022.10.005
2. Dickerman BA, García-Albéniz X, Logan RW, Denaxas S, Hernán MA. Avoidable flaws in observational analyses: an application to statins and cancer. *Nat Med*. 2019;25(10):1601-1606. doi:10.1038/s41591-019-0597-x
3. Hernán MA, Sauer BC, Hernández-Díaz S, Platt R, Shrier I. Specifying a target trial prevents immortal time bias and other self-inflicted injuries in observational analyses. *J Clin Epidemiol*. 2016;79:70-75. doi:10.1016/j.jclinepi.2016.04.014
4. Hernán MA, Robins JM. Using big data to emulate a target trial when a randomized trial is not available. *Am J Epidemiol*. 2016;183(8):758-764. doi:10.1093/aje/kwv254
5. Hernán MA, Wang W, Leaf DE. Target trial emulation: a framework for causal inference from observational data. *JAMA*. 2022;328(24):2446-2447. doi:10.1001/jama.2022.21383
6. Cochran WG. Observational studies. *Statistical Papers in Honor of George W Snedecor*. Iowa State University Press. 1972:77-90. doi:10.2307/2346927
7. Dorn HF. Philosophy of inferences from retrospective studies. *Am J Public Health Nations Health*. 1953;43(6 Pt 1):677-683. doi:10.2105/AJPH.43.6.Pt_1.677
8. Robins J. A new approach to causal inference in mortality studies with a sustained exposure period—application to control of the healthy worker survivor effect. *Math Model*. 1986;7(9-12):1393-1512. doi:10.1016/0270-0255(86)90088-6
9. Rubin DB. Estimating causal effects of treatments in randomized and nonrandomized studies. *J Educ Psychol*. 1974;66(5):688. doi:10.1037/h0037350
10. Wold H. Causality and econometrics. *Econometrica*. 1954;22(2):162-177. doi:10.2307/1907540
11. Hernán MA. Methods of public health research—strengthening causal inference from observational data. *N Engl J Med*. 2021;385(15):1345-1348. doi:10.1056/NEJMp2113319
12. Kutcher SA, Brophy JM, Banack HR, Kaufman JS, Samuel M. Emulating a randomized controlled trial with observational data: an introduction to the target trial framework. *Can J Cardiol*. 2021;37(9):1365-1377. doi:10.1016/j.cjca.2021.05.012
13. Matthews AA, Danaei G, Islam N, Kurth T. Target trial emulation: applying principles of randomized trials to observational studies. *BMJ*. 2022;378:e071108. doi:10.1136/bmj-2022-071108
14. Gomes M, Latimer N, Soares M, et al. Target trial emulation for transparent and robust estimation of treatment effects for health technology assessment using real-world data: opportunities and challenges. *Pharmacoeconomics*. 2022;40(6):577-586. doi:10.1007/s40273-022-01141-x
15. Moher D, Schulz KF, Simera I, Altman DG. Guidance for developers of health research reporting guidelines. *PLoS Med*. 2010;7(2):e1000217. doi:10.1371/journal.pmed.1000217
16. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372(71):n71. doi:10.1136/bmj.n71
17. Foster ED, Deardorff A. Open Science Framework (OSF). *J Med Libr Assoc*. 2017;105(2):203-206. doi:10.5195/jmla.2017.88
18. Hansford H, Lee H, Cashin A, et al. What is reported in target trial emulations: a systematic review protocol. March 13, 2022. Accessed August 29, 2023. <https://osf.io/uj56m/>
19. Haddaway H. citationchaser: An R package for forward and backward citations chasing in academic searching. Version 0.0.3. 2021. Accessed August 28, 2023. <https://github.com/nealhaddaway/citationchaser>

20. Hernán MA. How to estimate the effect of treatment duration on survival outcomes using observational data. *BMJ*. 2018;360:k182. doi:10.1136/bmj.k182
21. Robins JM, Hernan MJ. Estimation of the causal effects of time-varying exposures. In: Fitzmaurice G, Davidian M, Verbeke G, Molenberghs G, eds. *Longitudinal Data Analysis*. CRC press; 2008. doi:10.1201/9781420011579.ch23
22. Aakjaer M, Werther SK, De Bruin ML, Andersen M. Serious arrhythmia in initiators of citalopram, escitalopram, and other selective serotonin reuptake inhibitors: a population-based cohort study in older adults. *Clin Transl Sci*. 2022;15(9):2105-2115. doi:10.1111/cts.13319
23. Abrahami D, Pradhan R, Yin H, Honig P, Baumfeld Andre E, Azoulay L. Use of real-world data to emulate a clinical trial and support regulatory decision making: assessing the impact of temporality, comparator choice, and method of adjustment. *Clin Pharmacol Ther*. 2021;109(2):452-461. doi:10.1002/cpt.2012
24. Admon AJ, Donnelly JP, Casey JD, et al. Emulating a novel clinical trial using existing observational data. predicting results of the PreVent Study. *Ann Am Thorac Soc*. 2019;16(8):998-1007. doi:10.1513/AnnalsATS.201903-241OC
25. Ahn N, Nolde M, Günter A, et al. Emulating a target trial of proton pump inhibitors and dementia risk using claims data. *Eur J Neurol*. 2022;29(5):1335-1343. doi:10.1111/ene.15284
26. Al-Samkari H, Gupta S, Leaf RK, et al; STOP-COVID Investigators. Thrombosis, bleeding, and the observational effect of early therapeutic anticoagulation on survival in critically ill patients with COVID-19. *Ann Intern Med*. 2021;174(5):622-632. doi:10.7326/M20-6739
27. Prats-Urbe A, Tobed M, Villacampa JM, et al; TraqueoCOVID SEORL Group. Timing of elective tracheotomy and duration of mechanical ventilation among patients admitted to intensive care with severe COVID-19: a multicenter prospective cohort study. *Head Neck*. 2021;43(12):3743-3756. doi:10.1002/hed.26863
28. Althunian TA, de Boer A, Groenwold RHH, Rengerink KO, Souverein PC, Klungel OH. Rivaroxaban was found to be noninferior to warfarin in routine clinical care: a retrospective noninferiority cohort replication study. *Pharmacoepidemiol Drug Saf*. 2020;29(10):1263-1272. doi:10.1002/pds.5065
29. Ankarcrona V, Zhao H, Jacobsson B, Brismar Wendel S. Obstetric anal sphincter injury after episiotomy in vacuum extraction: an epidemiological study using an emulated randomized trial approach. *BJOG*. 2021;128(10):1663-1671. doi:10.1111/1471-0528.16663
30. Atkinson A, Zwahlen M, Barger D, et al; Opportunistic Infections Project Working Group of the Collaboration of Observational HIV Epidemiological Research Europe (COHERE) in EuroCoord. Withholding primary pneumocystis pneumonia prophylaxis in virologically suppressed patients with human immunodeficiency virus: an emulation of a pragmatic trial in COHERE. *Clin Infect Dis*. 2021;73(2):195-202. doi:10.1093/cid/ciaa615
31. Aubert CE, Sussman JB, Hofer TP, Cushman WC, Ha JK, Min L. Adding a new medication versus maximizing dose to intensify hypertension treatment in older adults : a retrospective observational study. *Ann Intern Med*. 2021;174(12):1666-1673. doi:10.7326/M21-1456
32. Bacic J, Liu T, Thompson RH, et al. Emulating target clinical trials of radical nephrectomy with or without lymph node dissection for renal cell carcinoma. *Urology*. 2020;140:98-106. doi:10.1016/j.urology.2020.01.039
33. Bai L, Scott MKD, Steinberg E, et al. Computational drug repositioning of atorvastatin for ulcerative colitis. *J Am Med Inform Assoc*. 2021;28(11):2325-2335. doi:10.1093/jamia/ocab165
34. Barbulescu A, Asklung J, Saevarsdottir S, Kim SC, Frisell T. Combined conventional synthetic disease modifying therapy vs. infliximab for rheumatoid arthritis: emulating a randomized trial in observational data. *Clin Pharmacol Ther*. 2022;112(4):836-845. doi:10.1002/cpt.2673
35. Barda N, Dagan N, Ben-Shlomo Y, et al. Safety of the BNT162b2 mRNA COVID-19 vaccine in a nationwide setting. *N Engl J Med*. 2021;385(12):1078-1090. doi:10.1056/NEJMoa2110475
36. Barda N, Dagan N, Cohen C, et al. Effectiveness of a third dose of the BNT162b2 mRNA COVID-19 vaccine for preventing severe outcomes in Israel: an observational study. *Lancet*. 2021;398(10316):2093-2100. doi:10.1016/S0140-6736(21)02249-2
37. Becker WC, Li Y, Caniglia EC, et al. Cannabis use, pain interference, and prescription opioid receipt among persons with HIV: a target trial emulation study. *AIDS Care*. 2022;34(4):469-477. doi:10.1080/09540121.2021.1944597
38. Bell Gorrod H, Court R, Schomaker M, Maartens G, Murphy RA. Increased mortality with delayed and missed switch to second-line antiretroviral therapy in South Africa. *J Acquir Immune Defic Syndr*. 2020;84(1):107-113. doi:10.1097/QAI.0000000000002313

39. Bergqvist R, Ahlqvist VH, Lundberg M, et al. HMG-CoA reductase inhibitors and COVID-19 mortality in Stockholm, Sweden: a registry-based cohort study. *PLoS Med*. 2021;18(10):e1003820. doi:10.1371/journal.pmed.1003820
40. Bharadwaj M, Kaul S, Fleishman A, et al. Adjuvant chemotherapy versus observation following radical cystectomy for locally advanced urothelial carcinoma of the bladder. *Urol Oncol*. 2022;40(6):274.e15-274.e23. doi:10.1016/j.urolonc.2022.02.002
41. Biscaglia S, Erriquez A, Serenelli M, et al. Complete versus culprit-only strategy in older MI patients with multivessel disease. *Catheter Cardiovasc Interv*. 2022;99(4):970-978. doi:10.1002/ccd.30075
42. Börnhorst C, Reinders T, Rathmann W, et al. Avoiding time-related biases: a feasibility study on antidiabetic drugs and pancreatic cancer applying the parametric g-formula to a large German healthcare database. *Clin Epidemiol*. 2021;13(13):1027-1038. doi:10.2147/CLEP.S328342
43. Bosch NA, Law AC, Vail EA, et al. Inhaled nitric oxide vs epoprostenol during acute respiratory failure: an observational target trial emulation. *Chest*. 2022;162(6):1287-1296. doi:10.1016/j.chest.2022.08.001
44. Boström P, Hultberg DK, Häggström J, et al. Oncological impact of high vascular tie after surgery for rectal cancer: a nationwide cohort study. *Ann Surg*. 2021;274(3):e236-e244. doi:10.1097/SLA.0000000000003663
45. Boyne DJ, Cheung WY, Hilsden RJ, et al. Association of a shortened duration of adjuvant chemotherapy with overall survival among individuals with stage III colon cancer. *JAMA Netw Open*. 2021;4(3):e213587. doi:10.1001/jamanetworkopen.2021.3587
46. Braitmaier M, Schwarz S, Kollhorst B, Senore C, Didelez V, Haug U. Screening colonoscopy similarly prevented distal and proximal colorectal cancer: a prospective study among 55-69-year-olds. *J Clin Epidemiol*. 2022;149:118-126. doi:10.1016/j.jclinepi.2022.05.024
47. Brouwer ES, Napravnik S, Eron JJ Jr, et al. Effects of combination antiretroviral therapies on the risk of myocardial infarction among HIV patients. *Epidemiology*. 2014;25(3):406-417. doi:10.1097/EDE.0000000000000041
48. Bruun-Rasmussen P, Andersen PK, Banasik K, Brunak S, Johansson PI. Estimating the effect of donor sex on red blood cell transfused patient mortality: a retrospective cohort study using a targeted learning and emulated trials-based approach. *EClinicalMedicine*. 2022;51:101628. doi:10.1016/j.eclinm.2022.101628
49. Bruun-Rasmussen P, Kragh Andersen P, Banasik K, Brunak S, Johansson PI. Intervening on the storage time of RBC units and its effects on adverse recipient outcomes using real-world data. *Blood*. 2022;139(25):3647-3654. doi:10.1182/blood.2022015892
50. Bujkiewicz S, Singh J, Wheaton L, et al. Bridging disconnected networks of first and second lines of biologic therapies in rheumatoid arthritis with registry data: andomiz evidence synthesis with target trial emulation. *J Clin Epidemiol*. 2022;150:171-178. doi:10.1016/j.jclinepi.2022.06.011
51. Buranupakorn T, Thangsuk P, Patumanond J, Phinyo P. Emulation of a target trial to evaluate the causal effect of palliative care consultation on the survival time of patients with hepatocellular carcinoma. *Cancers (Basel)*. 2021;13(5):992. doi:10.3390/cancers13050992
52. Burn E, Weaver J, Morales DR, et al. Opioid use, postoperative complications, and implant survival after unicompartmental versus total knee replacement: a population-based network study. *Lancet Rheumatol*. 2019;1(4):229-236. doi:10.1016/S2665-9913(19)30075-X
53. Butt AA, Talisa VB, Shaikh OS, Omer SB, Mayr FB. Relative vaccine effectiveness of a severe acute respiratory syndrome coronavirus 2 messenger RNA vaccine booster dose against the Omicron variant. *Clin Infect Dis*. 2022;75(12):2161-2168. doi:10.1093/cid/ciac328
54. Cain LE, Hernán MA; HIV-CAUSAL Collaboration. The effect of efavirenz versus nevirapine-containing regimens in the HIV-CAUSAL Collaboration: reply to Llibre and Podzamczar and additional results. *AIDS*. 2013;27(13):2169-2170. doi:10.1097/01.aids.0000432446.15061.27
55. Cain LE, Saag MS, Petersen M, et al; Antiretroviral Therapy Cohort Collaboration, the Centers for AIDS Research Network of Integrated Clinical Systems, and the HIV-CAUSAL Collaboration. Using observational data to emulate a randomized trial of dynamic treatment-switching strategies: an application to antiretroviral therapy. *Int J Epidemiol*. 2016;45(6):2038-2049. doi:10.1093/ije/dyv295
56. Caniglia EC, Robins JM, Cain LE, et al. Emulating a trial of joint dynamic strategies: an application to monitoring and treatment of HIV-positive individuals. *Stat Med*. 2019;38(13):2428-2446. doi:10.1002/sim.8120
57. Caniglia EC, Rojas-Saunero LP, Hilal S, et al. Emulating a target trial of statin use and risk of dementia using cohort data. *Neurology*. 2020;95(10):e1322-e1332. doi:10.1212/WNL.0000000000010433

58. Caniglia EC, Stevens ER, Khan M, et al. Does reducing drinking in patients with unhealthy alcohol use improve pain interference, use of other substances, and psychiatric symptoms? *Alcohol Clin Exp Res*. 2020;44(11):2257-2265. doi:10.1111/acer.14455
59. Caniglia EC, Zash R, Jacobson DL, et al. Emulating a target trial of antiretroviral therapy regimens started before conception and risk of adverse birth outcomes. *AIDS*. 2018;32(1):113-120. doi:10.1097/QAD.0000000000001673
60. Chauré-Pardos A, Aguilar-Palacio I, Rabanaque MJ, et al. Effectiveness of statins for primary prevention of cardiovascular disease in low- and medium-risk males: a causal inference approach with observational data. *J Pers Med*. 2022;12(5):658. doi:10.3390/jpm12050658
61. Chen YB, Liang CS, Wang LJ, et al. Comparative effectiveness of valproic acid in different serum concentrations for maintenance treatment of bipolar disorder: a retrospective cohort study using target trial emulation framework. *EClinicalMedicine*. 2022;54:101678. doi:10.1016/j.eclinm.2022.101678
62. Chen Z, Zhang H, Guo Y, et al. Exploring the feasibility of using real-world data from a large clinical data research network to simulate clinical trials of Alzheimer's disease. *NPJ Digit Med*. 2021;4(1):84. doi:10.1038/s41746-021-00452-1
63. Cheng Y, Zamrini E, Ahmed A, Wu WC, Shao Y, Zeng-Treitler Q. Medication-wide association study plus (MWAS+): a proof of concept study on drug repurposing. *Med Sci (Basel)*. 2022;10(3):48. doi:10.3390/medsci10030048
64. Chiu Y-H, Chavarro JE, Dickerman BA, et al. Estimating the effect of nutritional interventions using observational data: the American Heart Association's 2020 Dietary Goals and mortality. *Am J Clin Nutr*. 2021;114(2):690-703. doi:10.1093/ajcn/nqab100
65. Chiu YH, Yland JJ, Rinaudo P, et al. Effectiveness and safety of intrauterine insemination vs. assisted reproductive technology: emulating a target trial using an observational database of administrative claims. *Fertil Steril*. 2022;117(5):981-991. doi:10.1016/j.fertnstert.2022.02.003
66. Cho K, Keithly SC, Kurgansky KE, et al. Early convalescent plasma therapy and mortality among US veterans hospitalized with nonsevere COVID-19: an observational analysis emulating a target trial. *J Infect Dis*. 2021;224(6):967-975. doi:10.1093/infdis/jiab330
67. Dagan N, Barda N, Biron-Shental T, et al. Effectiveness of the BNT162b2 mRNA COVID-19 vaccine in pregnancy. *Nat Med*. 2021;27(10):1693-1695. doi:10.1038/s41591-021-01490-8
68. Dagan N, Barda N, Kepten E, et al. BNT162b2 mRNA COVID-19 vaccine in a nationwide mass vaccination setting. *N Engl J Med*. 2021;384(15):1412-1423. doi:10.1056/NEJMoa2101765
69. Danaei G, García Rodríguez LA, Fernández Cantero O, Hernán MA. Statins and risk of diabetes: an analysis of electronic medical records to evaluate possible bias due to differential survival. *Diabetes Care*. 2013;36(5):1236-1240. doi:10.2337/dc12-1756
70. Danaei G, Rodríguez LAG, Cantero OF, Logan R, Hernán MA. Observational data for comparative effectiveness research: an emulation of randomized trials of statins and primary prevention of coronary heart disease. *Stat Methods Med Res*. 2013;22(1):70-96. doi:10.1177/0962280211403603
71. Danaei G, García Rodríguez LA, Cantero OF, Logan RW, Hernán MA. Electronic medical records can be used to emulate target trials of sustained treatment strategies. *J Clin Epidemiol*. 2018;96:12-22. doi:10.1016/j.jclinepi.2017.11.021
72. Dave CV, Brittenham GM, Carson JL, Setoguchi S. Risks for anaphylaxis with intravenous iron formulations: a retrospective cohort study. *Ann Intern Med*. 2022;175(5):656-664. doi:10.7326/M21-4009
73. Dawwas GK, Cuker A, Barnes GD, Lewis JD, Hennessy S. Apixaban versus rivaroxaban in patients with atrial fibrillation and valvular heart disease: a population-based study. *Ann Intern Med*. 2022;175(11):1506-1514. doi:10.7326/M22-0318
74. Delapaz NR, Hor WK, Gilbert M, et al. An emulation of randomized trials of administering antipsychotics in PTSD patients for outcomes of suicide-related events. *J Pers Med*. 2021;11(3):178. doi:10.3390/jpm11030178
75. Deng Y, Polley EC, Wallach JD, et al. Emulating the GRADE trial using real world data: retrospective comparative effectiveness study. *BMJ*. 2022;379:e070717. doi:10.1136/bmj-2022-070717
76. Dickerman BA, García-Albéniz X, Logan RW, Denaxas S, Hernán MA. Emulating a target trial in case-control designs: an application to statins and colorectal cancer. *Int J Epidemiol*. 2020;49(5):1637-1646. doi:10.1093/ije/dyaa144
77. Dickerman BA, Gerlovin H, Madenci AL, et al. Comparative effectiveness of BNT162b2 and mRNA-1273 vaccines in U.S. veterans. *N Engl J Med*. 2022;386(2):105-115. doi:10.1056/NEJMoa2115463

78. Dickerman BA, Giovannucci E, Pernar CH, Mucci LA, Hernán MA. Guideline-Based physical activity and survival among US men with nonmetastatic prostate cancer. *Am J Epidemiol*. 2019;188(3):579-586. doi:10.1093/aje/kwy261
79. Dickerman BA, Madenci AL, Gerlovin H, et al. Comparative safety of BNT162b2 and mRNA-1273 vaccines in a nationwide cohort of US veterans. *JAMA Intern Med*. 2022;182(7):739-746. doi:10.1001/jamainternmed.2022.2109
80. Don EE, Mijatovic V, van Eekelen R, Huirne JAF. The effect of myomectomy on reproductive outcomes in patients with uterine fibroids: a retrospective cohort study. *Reprod Biomed Online*. 2022;45(5):970-978. doi:10.1016/j.rbmo.2022.05.025
81. Emilsson L, García-Albéniz X, Logan RW, Caniglia EC, Kalager M, Hernán MA. Examining bias in studies of statin treatment and survival in patients with cancer. *JAMA Oncol*. 2018;4(1):63-70. doi:10.1001/jamaoncol.2017.2752
82. Evans RN, Harris J, Rogers CA, MacGowan A. Emulating the MERINO randomized control trial using data from an observational cohort and trial of rapid diagnostic (BSI-FOO). *PLoS One*. 2022;17(5):e0268807. doi:10.1371/journal.pone.0268807
83. Farjat AE, Virdone S, Thomas LE, Kakkar AK, Pieper KS, Piccini JP. The importance of the design of observational studies in comparative effectiveness research: lessons from the GARFIELD-AF and ORBIT-AF registries. *Am Heart J*. 2022;243:110-121. doi:10.1016/j.ahj.2021.09.003
84. Fischer U, Branca M, Bonati LH, et al; Investigators of the Swiss Stroke Registry. Magnetic resonance imaging or computed tomography for suspected acute stroke: association of admission image modality with acute recanalization therapies, workflow metrics, and outcomes. *Ann Neurol*. 2022;92(2):184-194. doi:10.1002/ana.26413
85. Fotheringham J, Latimer N, Froissart M, et al. Survival on four compared with three times per week haemodialysis in high ultrafiltration patients: an observational study. *Clin Kidney J*. 2020;14(2):665-672. doi:10.1093/ckj/sfaa250
86. Franklin JM, Patorno E, Desai RJ, et al. Emulating randomized clinical trials with nonrandomized real-world evidence studies: first results from the RCT DUPLICATE Initiative. *Circulation*. 2021;143(10):1002-1013. doi:10.1161/CIRCULATIONAHA.120.051718
87. Fu EL, Evans M, Carrero JJ, et al. Timing of dialysis initiation to reduce mortality and cardiovascular events in advanced chronic kidney disease: nationwide cohort study. *BMJ*. 2021;375:e066306. doi:10.1136/bmj-2021-066306
88. Fu EL, Evans M, Clase CM, et al. Stopping renin-angiotensin system inhibitors in patients with advanced CKD and risk of adverse outcomes: a nationwide study. *J Am Soc Nephrol*. 2021;32(2):424-435. doi:10.1681/ASN.2020050682
89. Gaber CE, Shaheen NJ, Edwards JK, et al. Trimodality therapy vs definitive chemoradiation in older adults with locally advanced esophageal cancer. *JNCI Cancer Spectr*. 2022;6(6):pkac069. doi:10.1093/jncics/pkac069
90. Galloway L, Tran V-T, Perrodeau E, et al; COCO-OLD (Collaborative cOhort COrticosteroids for OLD patients with COvid-19) study Group. Fourteen-day survival among older adults with severe infection with severe acute respiratory syndrome coronavirus 2 treated with corticosteroid: a cohort study. *Clin Microbiol Infect*. 2021;27(8):1145-1150. doi:10.1016/j.cmi.2021.03.021
91. Garcia-Albeniz X, Chan JM, Paciorko A, et al. Immediate versus deferred initiation of androgen deprivation therapy in prostate cancer patients with PSA-only relapse: an observational follow-up study. *Eur J Cancer*. 2015;51(7):817-824. doi:10.1016/j.ejca.2015.03.003
92. García-Albéniz X, Hsu J, Bretthauer M, Hernán MA. Effectiveness of screening colonoscopy to prevent colorectal cancer among Medicare beneficiaries aged 70 to 79 years: a prospective observational study. *Ann Intern Med*. 2017;166(1):18-26. doi:10.7326/M16-0758
93. Garcia-Albeniz X, Hernan M, Logan R, Price M, Hsu J. Continuation of annual screening mammograms and breast-cancer mortality in women over 70. *Ann Oncol*. 2019;30(suppl 5):v675-v676. doi:10.1093/annonc/mdz263.016
94. Gazit S, Shlezinger R, Perez G, et al. The incidence of SARS-CoV-2 reinfection in persons with naturally acquired immunity with and without subsequent receipt of a single dose of BNT162b2 vaccine: a retrospective cohort study. *Ann Intern Med*. 2022;175(5):674-681. doi:10.7326/M21-4130
95. Ghiani M, Maywald U, Wilke T, Heeg B. Bridging the gap between oncology clinical trials and real-world data: evidence on replicability of efficacy results using German claims data. *J Comp Eff Res*. 2022;11(7):513-521. doi:10.2217/cer-2021-0224

96. Gilbert M, Dinh La A, Romulo Delapaz N, et al. An emulation of randomized trials of administering benzodiazepines in PTSD patients for outcomes of suicide-related events. *J Clin Med*. 2020;9(11):3492. doi:10.3390/jcm9113492
97. Girardi E, Caro-Vega Y, Cozzi-Lepri A, et al. The contribution of late HIV diagnosis on the occurrence of HIV-associated tuberculosis: a 5-year estimate using real-world data. *AIDS*. 2022;36(14):2005-2013. doi:10.1097/QAD.0000000000003321
98. Grøn KL, Grintborg B, Nørgaard M, et al. Comparative effectiveness of certolizumab pegol, abatacept, and biosimilar infliximab in patients with rheumatoid arthritis treated in routine care: observational data from the Danish DANBIO registry emulating a randomized trial. *Arthritis Rheumatol*. 2019;71(12):1997-2004. doi:10.1002/art.41031
99. Gupta S, Wang W, Hayek SS, et al; STOP-COVID Investigators. Association between early treatment with tocilizumab and mortality among critically ill patients with COVID-19. *JAMA Intern Med*. 2021;181(1):41-51. doi:10.1001/jamainternmed.2020.6252
100. Hajage D, Combes A, Guervilly C, et al; COVID-ICU Investigators. Extracorporeal membrane oxygenation for severe acute respiratory distress syndrome associated with COVID-19: an emulated target trial analysis. *Am J Respir Crit Care Med*. 2022;206(3):281-294. doi:10.1164/rccm.202111-2495OC
101. Hassan W, Shrestha P, Sumida K, et al. Association of uric acid-lowering therapy with incident chronic kidney disease. *JAMA Netw Open*. 2022;5(6):e2215878. doi:10.1001/jamanetworkopen.2022.15878
102. Hedderson MM, Badon SE, Pimentel N, et al. Association of glyburide and subcutaneous insulin with perinatal complications among women with gestational diabetes. *JAMA Netw Open*. 2022;5(3):e225026. doi:10.1001/jamanetworkopen.2022.5026
103. Heil TC, Verdaasdonk EGG, Maas HAAM, et al. Improved postoperative outcomes after prehabilitation for colorectal cancer surgery in older patients: an emulated target trial. *Ann Surg Oncol*. 2023;30(1):244-254. doi:10.1245/s10434-022-12623-9
104. Hou J, Zhao R, Cai T, et al. Temporal trends in clinical evidence of 5-year survival within electronic health records among patients with early-stage colon cancer managed with laparoscopy-assisted colectomy vs open colectomy. *JAMA Netw Open*. 2022;5(6):e2218371. doi:10.1001/jamanetworkopen.2022.18371
105. Htoo PT, Buse J, Cavender M, et al. Cardiovascular effectiveness of sodium-glucose cotransporter 2 inhibitors and glucagon-like peptide-1 receptor agonists in older patients in routine clinical care with or without history of atherosclerotic cardiovascular diseases or heart failure. *J Am Heart Assoc*. 2022;11(4):e022376. doi:10.1161/JAHA.121.022376
106. Huang H-K, Liu PP-S, Lin S-M, et al. Diabetes-related complications and mortality in patients with atrial fibrillation receiving different oral anticoagulants: a nationwide analysis. *Ann Intern Med*. 2022;175(4):490-498. doi:10.7326/M21-3498
107. Huang JY, Cai S, Huang Z, et al. Analyses of child cardiometabolic phenotype following assisted reproductive technologies using a pragmatic trial emulation approach. *Nat Commun*. 2021;12(1):5613. doi:10.1038/s41467-021-25899-4
108. Hulme WJ, Williamson EJ, Green ACA, et al. Comparative effectiveness of ChAdOx1 versus BNT162b2 COVID-19 vaccines in health and social care workers in England: cohort study using OpenSAFELY. *BMJ*. 2022;378:e068946. doi:10.1136/bmj-2021-068946
109. Ioannou GN, Locke ER, Green PK, Berry K. Comparison of Moderna versus Pfizer-BioNTech COVID-19 vaccine outcomes: a target trial emulation study in the U.S. Veterans Affairs healthcare system. *EClinicalMedicine*. 2022;45:101326. doi:10.1016/j.eclinm.2022.101326
110. Ioannou GN, Locke ER, O'Hare AM, et al. COVID-19 vaccination effectiveness against infection or death in a national US health care system: a target trial emulation study. *Ann Intern Med*. 2022;175(3):352-361. doi:10.7326/M21-3256
111. Izano MA, Lo JC, Adams AL, et al. Bisphosphonate treatment beyond 5 years and hip fracture risk in older women. *JAMA Netw Open*. 2020;3(12):e2025190. doi:10.1001/jamanetworkopen.2020.25190
112. Jang HY, Kim IW, Oh JM. Using real-world data for supporting regulatory decision making: comparison of cardiovascular and safety outcomes of an empagliflozin randomized clinical trial versus real-world data. *Front Pharmacol*. 2022;13:928121. doi:10.3389/fphar.2022.928121
113. Jin X, Ding C, Hunter DJ, Gallego B. Effectiveness of vitamin D supplementation on knee osteoarthritis: a target trial emulation study using data from the Osteoarthritis Initiative cohort. *Osteoarthritis Cartilage*. 2022;30(11):1495-1505. doi:10.1016/j.joca.2022.06.005

114. Johri M, Sylvestre M-P, Koné GK, Chandra D, Subramanian SV. Effects of improved drinking water quality on early childhood growth in rural Uttar Pradesh, India: a propensity-score analysis. *PLoS One*. 2019;14(1):e0209054. doi:10.1371/journal.pone.0209054
115. Karoboyas A, Morgenstern H, Fleischer NL, Schaubel DE, Robinson BM. Replicating randomized trial results with observational data using the parametric g-formula: an application to intravenous iron treatment in hemodialysis patients. *Clin Epidemiol*. 2020;12:1249-1260. doi:10.2147/CLEP.S283321
116. Katsoulis M, Stavola BD, Diaz-Ordaz K, et al. Weight change and the onset of cardiovascular diseases: emulating trials using electronic health records. *Epidemiology*. 2021;32(5):744-755. doi:10.1097/EDE.0000000000001393
117. Kaura A, Sterne JAC, Trickey A, et al. Invasive versus non-invasive management of older patients with non-ST elevation myocardial infarction (SENIOR-NSTEMI): a cohort study based on routine clinical data. *Lancet*. 2020;396(10251):623-634. doi:10.1016/S0140-6736(20)30930-2
118. Kazda L, McGeechan K, Bell K, Thomas R, Barratt A. Association of attention-deficit/hyperactivity disorder diagnosis with adolescent quality of life. *JAMA Netw Open*. 2022;5(10):e2236364. doi:10.1001/jamanetworkopen.2022.36364
119. Keil AP, Buckley JP, Kalkbrenner AE. Bayesian G-computation for estimating impacts of interventions on exposure mixtures: demonstration with metals from coal-fired power plants and birth weight. *Am J Epidemiol*. 2021;190(12):2647-2657. doi:10.1093/aje/kwab053
120. Keshet A, Rossman H, Shilo S, et al. Estimating the effect of cesarean delivery on long-term childhood health across two countries. *PLoS One*. 2022;17(10):e0268103. doi:10.1371/journal.pone.0268103
121. Keyhani S, Cheng EM, Hoggatt K, et al. Comparative effectiveness of carotid stenting to medical therapy among patients with asymptomatic carotid stenosis. *Stroke*. 2022;53(4):1157-1166. doi:10.1161/STROKEAHA.121.036178
122. Keyhani S, Cheng EM, Hoggatt KJ, et al. Comparative effectiveness of carotid endarterectomy vs initial medical therapy in patients with asymptomatic carotid stenosis. *JAMA Neurol*. 2020;77(9):1110-1121. doi:10.1001/jamaneurol.2020.1427
123. Khosrow-Khavar F, Desai RJ, Lee H, Lee SB, Kim SC. Tofacitinib and risk of malignancy: results from the Safety of Tofacitinib in Routine Care Patients With Rheumatoid Arthritis (STAR-RA) Study. *Arthritis Rheumatol*. 2022;74(10):1648-1659. doi:10.1002/art.42250
124. Kim Y, Zhang K, Savitz SI, Chen L, Schulz PE, Jiang X. Counterfactual analysis of differential comorbidity risk factors in Alzheimer's disease and related dementias. *PLoS Digit Health*. 2022;1(3):e0000018. doi:10.1371/journal.pdig.0000018
125. Kirchgerner J, Desai RJ, Beaugerie L, Kim SC, Schneeweiss S. Calibrating real-world evidence studies against randomized trials: treatment effectiveness of infliximab in Crohn's disease. *Clin Pharmacol Ther*. 2022;111(1):179-186. doi:10.1002/cpt.2304
126. Kirchgerner J, Desai RJ, Schneeweiss MC, Beaugerie L, Kim SC, Schneeweiss S. Emulation of a randomized controlled trial in ulcerative colitis with US and French claims data: infliximab with thiopurines compared to infliximab monotherapy. *Pharmacoepidemiol Drug Saf*. 2022;31(2):167-175. doi:10.1002/pds.5356
127. Kirchgerner J, Desai RJ, Schneeweiss MC, Beaugerie L, Schneeweiss S, Kim SC. Decreased risk of treatment failure with vedolizumab and thiopurines combined compared with vedolizumab monotherapy in Crohn's disease. *Gut*. 2022;71(9):1781-1789. doi:10.1136/gutjnl-2022-327002
128. Laroche MR, Lodi S, Yan S, Clothier BA, Goldsmith ES, Bohnert ASB. Comparative effectiveness of opioid tapering or abrupt discontinuation vs no dosage change for opioid overdose or suicide for patients receiving stable long-term opioid therapy. *JAMA Netw Open*. 2022;5(8):e2226523. doi:10.1001/jamanetworkopen.2022.26523
129. Lazzati A, Epaud S, Ortala M, Katsahian S, Lanoy E. Effect of bariatric surgery on cancer risk: results from an emulated target trial using population-based data. *Br J Surg*. 2022;109(5):433-438. doi:10.1093/bjs/znac003
130. Lim C, Mo Y, Teparukkul P, et al. Effect of delays in concordant antibiotic treatment on mortality in patients with hospital-acquired *Acinetobacter* species bacteremia: emulating a target randomized trial with a 13-year retrospective cohort. *Am J Epidemiol*. 2021;190(11):2395-2404. doi:10.1093/aje/kwab158
131. LoMartire R, Björk M, Dahlström Ö, et al. The value of interdisciplinary treatment for sickness absence in chronic pain: a nationwide register-based cohort study. *Eur J Pain*. 2021;25(10):2190-2201. doi:10.1002/ejp.1832
132. Lu Y, Gehr AW, Meadows RJ, et al. Timing of adjuvant chemotherapy initiation and mortality among colon cancer patients at a safety-net health system. *BMC Cancer*. 2022;22(1):593. doi:10.1186/s12885-022-09688-w
133. Lyu H, Yoshida K, Zhao SS, et al. Delayed denosumab injections and fracture risk among patients with osteoporosis: a population-based cohort study. *Ann Intern Med*. 2020;173(7):516-526. doi:10.7326/M20-0882

134. Madenci AL, Wanis KN, Cooper Z, et al. Strengthening health services research using target trial emulation: an application to volume-outcomes studies. *Am J Epidemiol*. 2021;190(11):2453-2460. doi:10.1093/aje/kwab170
135. Madenci AL, Wanis KN, Cooper Z, et al. Comparison of mortality risk with different surgeon and hospital operative volumes among individuals undergoing pancreatectomy by emulating target trials in US Medicare beneficiaries. *JAMA Netw Open*. 2022;5(3):e221766. doi:10.1001/jamanetworkopen.2022.1766
136. Magen O, Waxman JG, Makov-Assif M, et al. Fourth dose of BNT162b2 mRNA COVID-19 vaccine in a nationwide setting. *N Engl J Med*. 2022;386(17):1603-1614. doi:10.1056/NEJMoa2201688
137. Maguen S, Madden E, Holder N, et al. Effectiveness and comparative effectiveness of evidence-based psychotherapies for posttraumatic stress disorder in clinical practice. *Psychol Med*. 2023;53(2):419-428. doi:10.1017/S0033291721001628
138. Martin T, Krishnan A, Yong K, et al. Comparative effectiveness of ciltacabtagene autoleucel in CARTITUDE-1 versus physician's choice of therapy in the Flatiron Health multiple myeloma cohort registry for the treatment of patients with relapsed or refractory multiple myeloma. *EJHaem*. 2021;3(1):97-108. doi:10.1002/jha2.312
139. Martínez-Alés G, Domingo-Relloso A, Quintana-Díaz M, Fernández-Capitán C, Hernán MA; COVID@HULP Group. Thromboprophylaxis with standard-dose vs. flexible-dose heparin for hospitalized COVID-19 patients: a target trial emulation. *J Clin Epidemiol*. 2022;151:96-103. doi:10.1016/j.jclinepi.2022.08.006
140. Martínez-Sanz J, Ron R, Moreno E, et al. Similar CD4/CD8 ratio recovery after initiation of dolutegravir plus lamivudine versus dolutegravir or bictegravir-based three-drug regimens in adults with HIV. *Front Immunol*. 2022;13:873408. doi:10.3389/fimmu.2022.873408
141. Mathews KS, Soh H, Shaefi S, et al; STOP-COVID Investigators. Prone positioning and survival in mechanically ventilated patients with coronavirus disease 2019-related respiratory failure. *Crit Care Med*. 2021;49(7):1026-1037. doi:10.1097/CCM.0000000000004938
142. Matthews AA, Dahabreh IJ, Fröbert O, et al. Benchmarking observational analyses before using them to address questions trials do not answer: an application to coronary thrombus aspiration. *Am J Epidemiol*. 2022;191(9):1652-1665. doi:10.1093/aje/kwac098
143. Matthews AA, Szummer K, Dahabreh IJ, et al. Comparing effect estimates in randomized trials and observational studies from the same population: an application to percutaneous coronary intervention. *J Am Heart Assoc*. 2021;10(11):e020357. doi:10.1161/JAHA.120.020357
144. Mattishent K, Richardson K, Dhatariya K, Savva GM, Fox C, Loke YK. The effects of hypoglycaemia and dementia on cardiovascular events, falls and fractures and all-cause mortality in older individuals: a retrospective cohort study. *Diabetes Obes Metab*. 2019;21(9):2076-2085. doi:10.1111/dom.13769
145. Mazzotta V, Cozzi-Lepri A, Colavita F, et al. Emulation of a target trial from observational data to compare effectiveness of casirivimab/imdevimab and bamlanivimab/etesevimab for early treatment of non-hospitalized patients with COVID-19. *Front Immunol*. 2022;13:868020. doi:10.3389/fimmu.2022.868020
146. McDermott G, Fu X, Cook C, et al. The effect of achieving serological remission on subsequent risk of relapse, end-stage renal disease and mortality in ANCA-associated vasculitis: a target trial emulation study. *Ann Rheum Dis*. 2022;81(10):1438-1444. doi:10.1136/annrheumdis-2022-222439
147. McGrath LJ, Nielson C, Saul B, et al. Lessons learned using real-world data to emulate randomized trials: a case study of treatment effectiveness for newly diagnosed immune thrombocytopenia. *Clin Pharmacol Ther*. 2021;110(6):1570-1578. doi:10.1002/cpt.2399
148. Medrinal C, Gillet A, Boujibar F, et al. Role of non-invasive respiratory supports in COVID-19 acute respiratory failure patients with do not intubate orders. *J Clin Med*. 2021;10(13):2783. doi:10.3390/jcm10132783
149. Mei H, Wang J, Ma S. An emulated target trial analysis based on Medicare data suggested non-inferiority of dabigatran versus rivaroxaban. *J Clin Epidemiol*. 2021;139:28-37. doi:10.1016/j.jclinepi.2021.07.001
150. Mei H, Xu Y, Wang J, Ma S. Evaluation of survival outcomes of endovascular versus open aortic repair for abdominal aortic aneurysms with a big data approach. *Entropy (Basel)*. 2020;22(12):1349. doi:10.3390/e22121349
151. Meid AD, Wirbka L, Groll A, Haefeli WE; ARMIN Study Group. Can machine learning from real-world data support drug treatment decisions? a prediction modeling case for direct oral anticoagulants. *Med Decis Making*. 2022;42(5):587-598. doi:10.1177/0272989X211064604
152. Melgaard L, Overvad TF, Jensen M, et al. Effectiveness and safety of NOAC versus warfarin in patients with atrial fibrillation and aortic stenosis. *J Am Heart Assoc*. 2021;10(23):e022628. doi:10.1161/JAHA.121.022628
153. Merola D, Young J, Schrag D, Lin KJ, Robert N, Schneeweiss S. Oncology drug effectiveness from electronic health record data calibrated against RCT evidence: the PARSIFAL trial emulation. *Clin Epidemiol*. 2022;14:1135-1144. doi:10.2147/CLEP.S373291

154. Meyer A, Neumann A, Drouin J, Weill A, Carbonnel F, Dray-Spira R. Benefits and risks associated with continuation of anti-tumor necrosis factor after 24 weeks of pregnancy in women with inflammatory bowel disease: a nationwide emulation trial. *Ann Intern Med*. 2022;175(10):1374-1382. doi:10.7326/M22-0819
155. Miranda R, Smets T, De Schreye R, et al. Improved quality of care and reduced healthcare costs at the end-of-life among older people with dementia who received palliative home care: a nationwide propensity score-matched decedent cohort study. *Palliat Med*. 2021;35(9):1701-1712. doi:10.1177/02692163211019321
156. Moler-Zapata S, Grieve R, Lugo-Palacios D, et al. Local instrumental variable methods to address confounding and heterogeneity when using electronic health records: an application to emergency surgery. *Med Decis Making*. 2022;42(8):1010-1026. doi:10.1177/0272989X221100799
157. Monge S, Rojas-Benedicto A, Olmedo C, et al; IBERCovid. Effectiveness of mRNA vaccine boosters against infection with the SARS-CoV-2 Omicron (B.1.1.529) variant in Spain: a nationwide cohort study. *Lancet Infect Dis*. 2022;22(9):1313-1320. doi:10.1016/S1473-3099(22)00292-4
158. Moreno-Betancur M, Lynch JW, Pilkington RM, et al. Emulating a target trial of intensive nurse home visiting in the policy-relevant population using linked administrative data. *Int J Epidemiol*. 2023;52(1):119-131. doi:10.1093/ije/dyac092
159. Nabi H, Georgiadis S, Loft AG, et al. Comparative effectiveness of two adalimumab biosimilars in 1318 real-world patients with inflammatory rheumatic disease mandated to switch from originator adalimumab: nationwide observational study emulating a randomized clinical trial. *Ann Rheum Dis*. 2021;80(11):1400-1409. doi:10.1136/annrheumdis-2021-219951
160. Neugebauer R, Schroeder EB, Reynolds K, et al. Comparison of mortality and major cardiovascular events among adults with type 2 diabetes using human vs analogue insulins. *JAMA Netw Open*. 2020;3(1):e1918554. doi:10.1001/jamanetworkopen.2019.18554
161. Nolde M, Ahn N, Dreischulte T, et al. The long-term risk for myocardial infarction or stroke after proton pump inhibitor therapy (2008-2018). *Aliment Pharmacol Ther*. 2021;54(8):1033-1040. doi:10.1111/apt.16565
162. O'Donnell A, Pham N, Battisti L, et al. Estimating the causal effect of treatment with direct-acting antivirals on kidney function among individuals with hepatitis C virus infection. *PLoS One*. 2022;17(5):e0268478. doi:10.1371/journal.pone.0268478
163. Olarte Parra C, Waernbaum I, Schön S, Goetghebeur E. Trial emulation and survival analysis for disease incidence registers: a case study on the causal effect of pre-emptive kidney transplantation. *Stat Med*. 2022;41(21):4176-4199. doi:10.1002/sim.9503
164. Ozery-Flato M, Goldschmidt Y, Shaham O, Ravid S, Yanover C. Framework for identifying drug repurposing candidates from observational healthcare data. *JAMIA Open*. 2020;3(4):536-544. doi:10.1093/jamiaopen/ooaa048
165. Reijnders TDY, Peters-Sengers H, van Vught LA, et al; MARS consortium. Effect of erythromycin on mortality and the host response in critically ill patients with sepsis: a target trial emulation. *Crit Care*. 2022;26(1):151. doi:10.1186/s13054-022-04016-x
166. Petersen ML, Tran L, Geng EH, et al. Delayed switch of antiretroviral therapy after virologic failure associated with elevated mortality among HIV-infected adults in Africa. *AIDS*. 2014;28(14):2097-2107. doi:10.1097/QAD.0000000000000349
167. Peterson RG, Xiao R, Katcoff H, Fisher BT, Weiss PF. Effect of first-line biologic initiation on glucocorticoid exposure in children hospitalized with new-onset systemic juvenile idiopathic arthritis: emulation of a pragmatic trial using observational data. *Pediatr Rheumatol Online J*. 2021;19(1):109. doi:10.1186/s12969-021-00597-z
168. Petito LC, García-Albéniz X, Logan RW, et al. Estimates of overall survival in patients with cancer receiving different treatment regimens: emulating hypothetical target trials in the Surveillance, Epidemiology, and End Results (SEER)-Medicare linked database. *JAMA Netw Open*. 2020;3(3):e200452. doi:10.1001/jamanetworkopen.2020.0452
169. Porcher R, Desguerre I, Amthor H, et al. Association between prophylactic angiotensin-converting enzyme inhibitors and overall survival in Duchenne muscular dystrophy-analysis of registry data. *Eur Heart J*. 2021;42(20):1976-1984. doi:10.1093/eurheartj/ehab054
170. Prats-Urbe A, Kolovos S, Berencsi K, et al. Unicompartamental compared with total knee replacement for patients with multimorbidities: a cohort study using propensity score stratification and inverse probability weighting. *Health Technol Assess*. 2021;25(66):1-126. doi:10.3310/hta25660

171. Pufulete M, Harris J, Pouwels K, et al. Real-world bleeding in patients with acute coronary syndrome (ACS) undergoing percutaneous coronary intervention (PCI) and prescribed different combinations of dual antiplatelet therapy (DAPT) in England: a population-based cohort study emulating a 'target trial'. *Open Heart*. 2022;9(2): e001999. doi:10.1136/openhrt-2022-001999
172. Qiao N, Shen M, He W, et al. Comparative effectiveness of endoscopic versus microscopic transsphenoidal surgery for patients with growth hormone secreting pituitary adenoma: an emulated trial. *Clin Neurol Neurosurg*. 2021;207:106781. doi:10.1016/j.clineuro.2021.106781
173. Rahman S, Thomas B, Maynard N, et al. Impact of postoperative chemotherapy on survival for oesophagogastric adenocarcinoma after preoperative chemotherapy and surgery. *Br J Surg*. 2022;109(2): 227-236. doi:10.1093/bjs/znab427
174. Reitblat C, Fleishman A, Kaplan IA, et al. Radical prostatectomy versus external beam radiation therapy for high-grade, clinically localized prostate cancer: emulation of a target clinical trial. *Urol Oncol*. 2021;39(11):785.e1-785.e10. doi:10.1016/j.urolonc.2021.03.017
175. Rogawski McQuade ET, Benjamin-Chung J, Westreich D, Arnold BF. Population intervention effects in observational studies to emulate target trial results: reconciling the effects of improved sanitation on child growth. *Int J Epidemiol*. 2022;51(1):279-290. doi:10.1093/ije/dyab070
176. Rojas-Saunero LP, Hilal S, Murray EJ, Logan RW, Ikram MA, Swanson SA. Hypothetical blood-pressure-lowering interventions and risk of stroke and dementia. *Eur J Epidemiol*. 2021;36(1):69-79. doi:10.1007/s10654-020-00694-5
177. Rossides M, Kullberg S, Di Giuseppe D, et al. Infection risk in sarcoidosis patients treated with methotrexate compared to azathioprine: a retrospective 'target trial' emulated with Swedish real-world data. *Respirology*. 2021; 26(5):452-460. doi:10.1111/resp.14001
178. Sánchez A, Roos VH, Navarro M, et al. Quality of colonoscopy is associated with adenoma detection and postcolonoscopy colorectal cancer prevention in Lynch syndrome. *Clin Gastroenterol Hepatol*. 2022;20(3):611-621.e9, E9. doi:10.1016/j.cgh.2020.11.002
179. Schmidt M, Sørensen HT, Pedersen L. Cardiovascular risks of diclofenac versus other older COX-2 inhibitors (meloxicam and etodolac) and newer COX-2 inhibitors (celecoxib and etoricoxib): a series of nationwide emulated trials. *Drug Saf*. 2022;45(9):983-994. doi:10.1007/s40264-022-01211-1
180. Schmidt M, Sørensen HT, Pedersen L. Diclofenac use and cardiovascular risks: series of nationwide cohort studies. *BMJ*. 2018;362:k3426. doi:10.1136/bmj.k3426
181. Schneeweiss MC, Kim SC, Wyss R, Schneeweiss S, Merola JF. Dupilumab and the risk of conjunctivitis and serious infection in patients with atopic dermatitis: a propensity score-matched cohort study. *J Am Acad Dermatol*. 2021;84(2):300-311. doi:10.1016/j.jaad.2020.09.084
182. Schoenfeld AJ, Losina E, Ferrone ML, et al. Ambulatory status after surgical and nonsurgical treatment for spinal metastasis. *Cancer*. 2019;125(15):2631-2637. doi:10.1002/cncr.32140
183. Schroeder EB, Neugebauer R, Reynolds K, et al. Association of cardiovascular outcomes and mortality with sustained long-acting insulin only vs long-acting plus short-acting insulin treatment. *JAMA Netw Open*. 2021;4(9): e2126605. doi:10.1001/jamanetworkopen.2021.26605
184. Secora AM, Shin JI, Qiao Y, et al. Hyperkalemia and acute kidney injury with spironolactone use among patients with heart failure. *Mayo Clin Proc*. 2020;95(11):2408-2419. doi:10.1016/j.mayocp.2020.03.035
185. Seeger JD, Nunes A, Loughlin AM. Using RWE research to extend clinical trials in diabetes: an example with implications for the future. *Diabetes Obes Metab*. 2020;22(Suppl 3)(suppl 3):35-44. doi:10.1111/dom.14021
186. Shaefi S, Brenner SK, Gupta S, et al; STOP-COVID Investigators. Extracorporeal membrane oxygenation in patients with severe respiratory failure from COVID-19. *Intensive Care Med*. 2021;47(2):208-221. doi:10.1007/s00134-020-06331-9
187. Shin H, Schneeweiss S, Glynn RJ, Paterno E. Cardiovascular outcomes in patients initiating first-line treatment of type 2 diabetes with sodium-glucose cotransporter-2 inhibitors versus metformin: a cohort study. *Ann Intern Med*. 2022;175(7):927-937. doi:10.7326/M21-4012
188. Shin J-I, Fine DM, Sang Y, et al. Association of rosuvastatin use with risk of hematuria and proteinuria. *J Am Soc Nephrol*. 2022;33(9):1767-1777. doi:10.1681/ASN.2022020135
189. Simmonds KP, Burke J, Kozłowski AJ, Andary M, Luo Z, Reeves MJ. Emulating 3 clinical trials that compare stroke rehabilitation at inpatient rehabilitation facilities with skilled nursing facilities. *Arch Phys Med Rehabil*. 2022; 103(7):1311-1319. doi:10.1016/j.apmr.2021.12.029
190. Softness K, Kaul S, Fleishman A, et al. Radical cystectomy versus trimodality therapy for muscle-invasive urothelial carcinoma of the bladder. *Urol Oncol*. 2022;40(6):272.e1-272.e9. doi:10.1016/j.urolonc.2021.12.015

191. Solomon DH, Ruppert K, Habel LA, et al. Prescription medications for sleep disturbances among midlife women during 2 years of follow-up: a SWAN retrospective cohort study. *BMJ Open*. 2021;11(5):e045074. doi:10.1136/bmjopen-2020-045074
192. Stattin P, Westerberg M, Lissbrant IF, et al. Real world outcomes in patients with metastatic, castration-resistant prostate cancer treated with radium-223 in routine clinical practice in Sweden. *Clin Genitourin Cancer*. 2023;21(1):107.e1-107.e9. doi:10.1016/j.clgc.2022.09.002
193. Stokes JW, Gannon WD, Sherrill WH, et al. Bleeding, thromboembolism, and clinical outcomes in venovenous extracorporeal membrane oxygenation. *Crit Care Explor*. 2020;2(11):e0267. doi:10.1097/CCE.000000000000267
194. Strohmaier S, Wallisch C, Kammer M, et al. Survival benefit of first single-organ deceased donor kidney transplantation compared with long-term dialysis across ages in transplant-eligible patients with kidney failure. *JAMA Netw Open*. 2022;5(10):e2234971. doi:10.1001/jamanetworkopen.2022.34971
195. Sun JW, Young JG, Sarvet AL, et al. Comparison of rates of type 2 diabetes in adults and children treated with anticonvulsant mood stabilizers. *JAMA Netw Open*. 2022;5(4):e226484. doi:10.1001/jamanetworkopen.2022.6484
196. Sung RS, Zhang M, Schaubel DE, Shu X, Magee JC. A reassessment of the survival advantage of simultaneous kidney-pancreas versus kidney-alone transplantation. *Transplantation*. 2015;99(9):1900-1906. doi:10.1097/TP.0000000000000663
197. Swaminathan L, Kaatz S, Chubb H, et al. Impact of early corticosteroids on preventing clinical deterioration in non-critically ill patients hospitalized with COVID-19: a multi-hospital cohort study. *Infect Dis Ther*. 2022;11(2):887-898. doi:10.1007/s40121-022-00615-x
198. Szummer K, Montez-Rath ME, Alfredsson J, et al. Comparison between ticagrelor and clopidogrel in elderly patients with an acute coronary syndrome: insights from the SWEDEHEART registry. *Circulation*. 2020;142(18):1700-1708. doi:10.1161/CIRCULATIONAHA.120.050645
199. Takeuchi Y, Kumamaru H, Hagiwara Y, et al. Sodium-glucose cotransporter-2 inhibitors and the risk of urinary tract infection among diabetic patients in Japan: target trial emulation using a nationwide administrative claims database. *Diabetes Obes Metab*. 2021;23(6):1379-1388. doi:10.1111/dom.14353
200. Takeuchi Y, Shinozaki T, Kumamaru H, Hiramatsu T, Matsuyama Y. Analyzing intent-to-treat and per-protocol effects on safety outcomes using a medical information database: an application to the risk assessment of antibiotic-induced liver injury. *Expert Opin Drug Saf*. 2018;17(11):1071-1079. doi:10.1080/14740338.2018.1528224
201. Tapper EB, Zhao Z, Winder GS, Parikh ND. Deprescribing zolpidem reduces falls and fractures in patients with cirrhosis. *JHEP Rep*. 2022;4(6):100478. doi:10.1016/j.jhepr.2022.100478
202. Thamer M, Zhang Y, Kaufman J, Cotter D, Hernán MA. Similar outcomes for two anemia treatment strategies among elderly hemodialysis patients with diabetes. *J Endocrinol Diabetes*. 2014;1(2):6. doi:10.15226/2374-6890/1/2/00111
203. Thomas DS, Lee AY, Müller PL, et al; UK AMD EMR Users Group. Contextualizing single-arm trials with real-world data: an emulated target trial comparing therapies for neovascular age-related macular degeneration. *Clin Transl Sci*. 2021;14(3):1166-1175. doi:10.1111/cts.12974
204. Thomsen FB, Bosco C, Garmo H, et al. Anti-androgen monotherapy versus gonadotropin-releasing hormone agonists in men with advanced, non-metastatic prostate cancer: a register-based, observational study. *Acta Oncol*. 2019;58(1):110-118. doi:10.1080/0284186X.2018.1529427
205. Tran VT, Mahévas M, Bani-Sadr F, et al; COCORICO Collaborative Group. Corticosteroids in patients hospitalized for COVID-19 pneumonia who require oxygen: observational comparative study using routine care data. *Clin Microbiol Infect*. 2021;27(4):603-610. doi:10.1016/j.cmi.2020.11.035
206. Trevisan M, Fu EL, Xu Y, et al. Stopping mineralocorticoid receptor antagonists after hyperkalaemia: trial emulation in data from routine care. *Eur J Heart Fail*. 2021;23(10):1698-1707. doi:10.1002/ehf.2287
207. Tsilidis KK, Capothanassi D, Allen NE, et al. Metformin does not affect cancer risk: a cohort study in the U.K. Clinical Practice Research Datalink analyzed like an intention-to-treat trial. *Diabetes Care*. 2014;37(9):2522-2532. doi:10.2337/dc14-0584
208. van Santen DK, Boyd A, Matser A, et al. The effect of needle and syringe program and opioid agonist therapy on the risk of HIV, hepatitis B and C virus infection for people who inject drugs in Amsterdam, the Netherlands: findings from an emulated target trial. *Addiction*. 2021;116(11):3115-3126. doi:10.1111/add.15503
209. Wanis KN, Madenci AL, Dokus MK, et al. The meaning of confounding adjustment in the presence of multiple versions of treatment: an application to organ transplantation. *Eur J Epidemiol*. 2019;34(3):225-233. doi:10.1007/s10654-019-00484-8

210. Wei J, Choi HK, Neogi T, et al. Allopurinol initiation and all-cause mortality among patients with gout and concurrent chronic kidney disease: a population-based cohort study. *Ann Intern Med*. 2022;175(4):461-470. doi:10.7326/M21-2347
211. Weisel K, Martin T, Krishnan A, et al. Comparative efficacy of ciltacabtagene autoleucel in CARTITUDE-1 vs physician's choice of therapy in the long-term follow-up of POLLUX, CASTOR, and EQUULEUS clinical trials for the treatment of patients with relapsed or refractory multiple myeloma. *Clin Drug Investig*. 2022;42(1):29-41. doi:10.1007/s40261-021-01100-y
212. Xie Y, Bowe B, Yan Y, Xian H, Li T, Al-Aly Z. Estimates of all cause mortality and cause specific mortality associated with proton pump inhibitors among US veterans: cohort study. *BMJ*. 2019;365:l1580. doi:10.1136/bmj.l1580
213. Xie Y, Bowe B, Gibson AK, et al. Comparative effectiveness of SGLT2 inhibitors, GLP-1 receptor agonists, DPP-4 inhibitors, and sulfonylureas on risk of kidney outcomes: emulation of a target trial using health care databases. *Diabetes Care*. 2020;43(11):2859-2869. doi:10.2337/dc20-1890
214. Xu Y, Fu EL, Trevisan M, et al. Stopping renin-angiotensin system inhibitors after hyperkalemia and risk of adverse outcomes. *Am Heart J*. 2022;243:177-186. doi:10.1016/j.ahj.2021.09.014
215. an ZZN, Mason KJ, Hampton PJ, et al; BADBIR Study Group. Randomized trial replication using observational data for comparative effectiveness of secukinumab and ustekinumab in psoriasis: a study from the British Association of Dermatologists Biologics and Immunomodulators register. *JAMA Dermatol*. 2021;157(1):66-73. doi:10.1001/jamadermatol.2020.4202
216. Yland JJ, Chiu YH, Rinaudo P, Hsu J, Hernán MA, Hernández-Díaz S. Emulating a target trial of the comparative effectiveness of clomiphene citrate and letrozole for ovulation induction. *Hum Reprod*. 2022;37(4):793-805. doi:10.1093/humrep/deac005
217. Young J, Wong S, Janjua NZ, Klein MB. Comparing direct acting antivirals for hepatitis C using observational data—why and how? *Pharmacol Res Perspect*. 2020;8(5):e00650. doi:10.1002/prp2.650
218. Young J, Scherrer AU, Calmy A, et al; Swiss HIV Cohort Study. The comparative effectiveness of NRTI-sparing dual regimens in emulated trials using observational data from the Swiss HIV Cohort Study. *Antivir Ther*. 2019;24(5):343-353. doi:10.3851/IMP3310
219. Zhang Y, Thamer M, Kaufman J, Cotter D, Hernán MA. Comparative effectiveness of two anemia management strategies for complex elderly dialysis patients. *Med Care*. 2014;52(0 3)(suppl 3):S132-S139. doi:10.1097/MLR.0b013e3182a53ca8
220. Zhang Y, Young JG, Thamer M, Hernán MA. Comparing the effectiveness of dynamic treatment strategies using electronic health records: an application of the parametric g-formula to anemia management strategies. *Health Serv Res*. 2018;53(3):1900-1918. doi:10.1111/1475-6773.12718
221. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Bull World Health Organ*. 2007;85(11):867-872. doi:10.2471/BLT.07.045120
222. Vandenbroucke JP, von Elm E, Altman DG, et al; STROBE Initiative. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *PLoS Med*. 2007;4(10):e297. doi:10.1371/journal.pmed.0040297
223. Nguyen VT, Engleton M, Davison M, Ravaut P, Porcher R, Boutron I. Risk of bias in observational studies using routinely collected data of comparative effectiveness research: a meta-research study. *BMC Med*. 2021;19(1):279. doi:10.1186/s12916-021-02151-w
224. Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*. 2016;355:i4919. doi:10.1136/bmj.i4919
225. Rubin DB. The design versus the analysis of observational studies for causal effects: parallels with the design of randomized trials. *Stat Med*. 2007;26(1):20-36. doi:10.1002/sim.2739
226. Hernán MA. With great data comes great responsibility: publishing comparative effectiveness research in epidemiology. *Epidemiology*. 2011;22(3):290-291. doi:10.1097/EDE.0b013e3182114039
227. Hernán MA, Robins JM, García Rodríguez LA. Discussion on 'Statistical Issues Arising in the Women's Health Initiative'. *Biometrics*. 2005;61(4):922-930. doi:10.1111/j.0006-341X.2005.454.7.x
228. Hernán MA, Alonso A, Logan R, et al. Observational studies analyzed like randomized experiments: an application to postmenopausal hormone therapy and coronary heart disease. *Epidemiology*. 2008;19(6):766-779. doi:10.1097/EDE.0b013e3181875e61
229. National Institute of Health and Care Excellence. NICE real-world evidence framework. June 23, 2022. Accessed August 28, 2023. <https://www.nice.org.uk/corporate/ecd9/chapter/overview>

230. European Medicines Agency. European Medicines Agencies network strategy to 2025. Accessed August 28, 2023. <https://www.ema.europa.eu/en/about-us/how-we-work/andomiz-medicines-regulatory-network/andomiz-medicines-agencies-network-strategy>
231. Government of Canada. Optimizing the Use of Real World Evidence to Inform Regulatory Decision-Making. April 16, 2019. Accessed August 28, 2023. <https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/announcements/optimizing-real-world-evidence-regulatory-decisions.html>
232. Therapeutic Goods Administration. Real world evidence and patient reported outcomes in the regulatory context. February 28, 2023. Accessed August 28, 2023. <https://www.tga.gov.au/review-real-world-evidence-and-patient-reported-outcomes>
233. Canada's Drug and Health Technology Agency. Guidance for Reporting Real-World Evidence. May 2023. Accessed August 28, 2023. <https://www.cadth.ca/sites/default/files/RWE/MGO020/MGO020-RWE-Guidance-Report.pdf>
234. Altman DG, Simera I, Hoey J, Moher D, Schulz K. EQUATOR: reporting guidelines for health research. *Lancet*. 2008;371(9619):1149-1150. doi:10.1016/S0140-6736(08)60505-X
235. Moher D, Jones A, Lepage L; CONSORT Group (Consolidated Standards for Reporting of Trials). Use of the CONSORT statement and quality of reports of randomized trials: a comparative before-and-after evaluation. *JAMA*. 2001;285(15):1992-1995. doi:10.1001/jama.285.15.1992
236. Turner L, Shamseer L, Altman DG, et al. Consolidated standards of reporting trials (CONSORT) and the completeness of reporting of andomized controlled trials (RCTs) published in medical journals. *Cochrane Database Syst Rev*. 2012;11(11):MR000030. doi:10.1002/14651858.MR000030.pub2
237. Cobo E, Cortés J, Ribera JM, et al. Effect of using reporting guidelines during peer review on quality of final manuscripts submitted to a biomedical journal: masked andomized trial. *BMJ*. 2011;343:d6783. doi:10.1136/bmj.d6783
238. EQUATOR Network. Reporting guidelines under development for observational studies. Accessed May 29, 2023. <https://www.equator-network.org/library/reporting-guidelines-under-development/reporting-guidelines-under-development-for-observational-studies/>
239. Collins GS, Reitsma JB, Altman DG, Moons KG. Transparent Reporting of a multivariable prediction model for Individual Prognosis or Diagnosis (TRIPOD): the TRIPOD statement. *Ann Intern Med*. 2015;162(1):55-63. doi:10.7326/M14-0697
240. Ziemann S, Paetzolt I, Grüßer L, Coburn M, Rossaint R, Kowark A. Poor reporting quality of observational clinical studies comparing treatments of COVID-19—a retrospective cross-sectional study. *BMC Med Res Methodol*. 2022;22(1):23. doi:10.1186/s12874-021-01501-9
241. Wiehn E, Ricci C, Alvarez-Perea A, et al; Task Force 'Adherence to reporting guidelines in articles published in EAACI Journals: a systematic review' of the EAACI Working Group on Epidemiology. Adherence to the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) checklist in articles published in EAACI Journals: A bibliographic study. *Allergy*. 2021;76(12):3581-3588. doi:10.1111/all.14951
242. Wang SV, Schneeweiss S, Franklin JM, et al; RCT-DUPLICATE Initiative. Emulation of randomized clinical trials with nonrandomized database analyses: results of 32 clinical trials. *JAMA*. 2023;329(16):1376-1385. doi:10.1001/jama.2023.4221
243. Dahabreh IJ, Steingrimsson JA, Robins JM, Hernán MA. Randomized trials and their observational emulations: a framework for benchmarking and joint analysis. *arXiv*. Preprint posted online March 28, 2022. doi:10.48550/arXiv.2203.14857
244. Dahabreh IJ, Robins JM, Hernán MA. Benchmarking observational methods by comparing randomized trials and their emulations. *Epidemiology*. 2020;31(5):614-619. doi:10.1097/EDE.0000000000001231
245. Dahabreh IJ, Matthews A, Steingrimsson JA, Scharfstein DO, Stuart EA. Using trial and observational data to assess effectiveness: trial emulation, transportability, benchmarking, and joint analysis. *Epidemiol Rev*. 2023; mxac011. doi:10.1093/epirev/mxac011
246. Gershman B, Guo DP, Dahabreh IJ. Using observational data for personalized medicine when clinical trial evidence is limited. *Fertil Steril*. 2018;109(6):946-951. doi:10.1016/j.fertnstert.2018.04.005
247. Chalmers I, Glasziou P. Avoidable waste in the production and reporting of research evidence. *Lancet*. 2009; 374(9683):86-89. doi:10.1016/S0140-6736(09)60329-9
248. Glasziou P, Altman DG, Bossuyt P, et al. Reducing waste from incomplete or unusable reports of biomedical research. *Lancet*. 2014;383(9913):267-276. doi:10.1016/S0140-6736(13)62228-X
249. Higgins JP, Thomas J, Chandler J, et al. *Cochrane Handbook for Systematic Reviews of Interventions*. John Wiley & Sons; 2019. doi:10.1002/9781119536604

250. Lund JL, Richardson DB, Stürmer T. The active comparator, new user study design in pharmacoepidemiology: historical foundations and contemporary application. *Curr Epidemiol Rep*. 2015;2(4):221-228. doi:10.1007/s40471-015-0053-5
251. Ray WA. Evaluating medication effects outside of clinical trials: new-user designs. *Am J Epidemiol*. 2003;158(9):915-920. doi:10.1093/aje/kwg231

SUPPLEMENT 1.

Appendix 1. Methodological Differences to the Protocol

Appendix 2. Complete Search Strategies for All Databases

Appendix 3. List of Phrases Deemed to Indicate Explicit Emulation of a Target Trial

Appendix 4. Operational Definitions for Items of the Emulation of the Target Trial

Appendix 5. Items Extracted That Are Not Included in Commonly Used Reporting Guidelines (STROBE, RECORD, ISPOR)

Appendix 6. Reasons for Study Exclusion at Level of Full Text

Appendix 7. Subgroup Analysis of Reporting of Target Trial Protocol and How It Was Emulated, by Reporting Guideline Use

SUPPLEMENT 2.

Data Sharing Statement