

1 **Diagnosis and Evaluation of Prognosis of Myelofibrosis: A British Society for**

2 **Haematology Guideline**

3 Donal. P McLornan¹, Anna L. Godfrey², Anna Green³, Rebecca Frewin⁴, Siamak
4 Arami⁵, Jessica Brady⁶, Nauman M. Butt⁷, Catherine Cargo⁸, Joanne Ewing⁹,
5 Sebastian Francis¹⁰ Mamta Garg¹¹, Claire Harrison¹², Andrew Innes¹³, Alesia Khan⁸,
6 Steve Knapper¹⁴, Jonathan Lambert¹, Adam Mead¹⁵, Andrew McGregor¹⁶, Pratap
7 Neelakantan¹⁷, Bethan Psaila¹⁵, Tim CP Somervaille¹⁸, Claire Woodley¹², Jyoti
8 Nangalia¹⁹, Nicholas C.P Cross²⁰, Mary Frances McMullin²¹

- 9 1, Department of Haematology, University College London Hospitals, London
10 2. Haematopathology & Oncology Diagnostics Service, Department of
11 Haematology, Cambridge University Hospitals NHS Foundation Trust
12 3. Department of Histopathology, Guy's and St. Thomas' NHS Foundation Trust,
13 London
14 4. Department of Haematology, Gloucestershire Hospitals NHS Foundation Trust
15 5. Department of Haematology, London Northwest Healthcare University NHS Trust
16 6. Department of Clinical Oncology, Guy's and St. Thomas' NHS Foundation Trust,
17 7. Department of Haematology, The Clatterbridge Cancer Centre NHS Foundation
18 Trust, Liverpool
19 8. Department of Haematology, Leeds Teaching Hospitals NHS Foundation Trust
20 9. Department of Haematology, University Hospitals Birmingham Trust
21 10. Department of Haematology, Sheffield Teaching Hospital NHS Foundation Trust
22 11. Department of Haematology, University Hospitals Leicester NHS Trust
23 12. Department of Haematology, Guy's and St. Thomas' NHS Foundation Trust,
24 London
25 13. Department of Haematology, Imperial College, London
26 14. Department of Haematology, Cardiff University, Cardiff
27 15. MRC Weatherall Institute of Molecular Medicine, University of Oxford and
28 Department of Haematology, Churchill Hospital, Oxford University NHS Trust
29 16. Department of Haematology, Freeman Hospital, Newcastle
30 17. Department of Haematology, Royal Berkshire NHS Foundation Trust
31 18. Cancer Research UK Manchester Institute & The Christie NHS Foundation Trust,
32 Manchester
33 19. Wellcome Sanger Institute, University of Cambridge
34 20. Faculty of Medicine, University of Southampton, Southampton, UK
35 21. Centre for Medical Education, Queen's University, Belfast
36

Correspondence: BSH Administrator, British Society for Haematology, 100 White
Lion Street, London, N1 9PF, UK. E-mail: bshguidelines@b-s-h.org.uk

Authorship: All authors contributed to guideline writing, review and editing.

Conflict of Interest Statements: All conflict of interest statements have been
registered with the British Society of Haematology.

Funding: There was no funding associated with this work

Data availability statement: NA

Ethics approval statement: NA

Patient consent statement: NA

Permission to reproduce material from other sources: NA

Clinical trial registration (including trial number): NA

Summary and Aims

This document represents an update of the British Society for Haematology (BSH) guideline on myelofibrosis first published in 2012 and updated in 2015 (1). This guideline aims to provide healthcare professionals with clear guidance on the diagnosis and prognostic evaluation of primary myelofibrosis (PMF), as well as post-polycythaemia myelofibrosis (post-PV MF) and post-essential thrombocythaemia myelofibrosis (post-ET MF). A section on pre-fibrotic myelofibrosis is also included. A separate BSH guideline covers the management of myelofibrosis and is published alongside this guideline.

Methodology

These guidelines were compiled according to the BSH process <https://b-s-h.org.uk/media/16732/bsh-guidance-development-process-dec-5-18.pdf>. The Grading of Recommendations, Assessment, Development and Evaluation (GRADE) nomenclature was used to evaluate levels of evidence and to assess the strength of recommendations. The GRADE criteria can be found at <http://www.gradeworkinggroup.org>. Recommendations are based on a review of the relevant myelofibrosis related literature using Medline, PubMed/Medline and Cochrane searches beginning from 2012 up to mid 2022. Filters were applied to include only publications written in English, studies carried out in humans, clinical conferences, congresses, clinical trials, clinical studies, meta-analyses, multicentre studies and randomised controlled trials. Exclusion criteria included papers published in non-English journals and those publications without an abstract.

Review of the manuscript

Review of the manuscript was performed by the BSH Guidelines Committee Haemato-oncology Task Force, the BSH Guidelines Committee and the Haemato-oncology sounding board of the BSH. We invited two global expert external reviewers to review contents – Professor Ruben Mesa and Professor Alessandro Vannucchi. This guideline has also been reviewed by patient representatives from MPN Voice.

Introduction

Myelofibrosis (MF) encompasses primary myelofibrosis (PMF), post-essential thrombocythaemia (ET) MF, and post-polycythaemia vera (PV) MF. It is characterised by clonal haemopoietic stem cell proliferation and elevated levels of pro-inflammatory cytokines, resulting in reticulin deposition and collagen fibrosis. The annual incidence is estimated at 1-2 individuals per 100,000 of the population in the United Kingdom, with an equal sex incidence (2). All patients newly diagnosed with MF should be reported to the National Cancer Registry, via the multidisciplinary meeting, and to MF-specific registries if available.

Clinical Features

Clinical features of MF are heterogeneous and may include anaemia, leucocytosis and extramedullary haemopoiesis, with progressive splenomegaly. Patients may experience constitutional symptoms, consequences of progressive splenomegaly (pain, early satiety, portal hypertension and dyspnoea), progressive marrow failure and have an inherent risk of leukaemic transformation.

105 Palpable splenomegaly is present in up to 80% of patients. Clinical palpation is the
106 easiest method to evaluate spleen size, with the patient in the supine position. Ideally,
107 a simple measuring tape can be used to record the size of an enlarged spleen below
108 the left costal margin in centimetres (cm). Ultrasound may aid spleen size
109 determination in a more uniform manner. In MF trials, the International Working Group
110 Myeloproliferative Neoplasm Research and Treatment (IWG-MRT) criteria utilised
111 spleen volume (SV) as part of the clinical improvement (CI) response; this can be
112 derived from computed tomography (CT) or magnetic resonance imaging (MRI). This
113 is not a requirement in routine clinical practice.

114 MF frequently has a significant symptom burden that can negatively impact on quality
115 of life, activities of daily living and functional status. Validated tools that have been
116 developed to objectively measure symptom burden include the MF Symptoms
117 Assessment Form (MF SAF), MPN Symptoms Assessment Form (MPN SAF) and
118 MPN-SAF Total Symptom Score (MPN SAF TSS – MPN 10) (3,4). The MPN SAF TSS
119 is an abbreviated symptom assessment tool that measures 10 symptoms through
120 patient self-assessment on a linear scale from 0 (absent) to 10 (worst imaginable),
121 namely: fatigue, early satiety, abdominal discomfort, inactivity, concentration
122 problems, night sweats, pruritus, bone pain, fever and weight loss. Regular use of
123 MPN SAF TSS provides an indication of symptom status and treatment response and
124 should be performed at each clinical review as appropriate.

125 Thrombosis risk is often underestimated in MF, in particular for those in the so-called
126 lower prognostic groups with a *JAK2* V617F mutation (5). An individualised risk
127 assessment is warranted.

Lastly, it is well established that some MF patients are at risk of non-cirrhotic portal hypertension, in particular those with bulky splenomegaly (6). Where clinical signs (e.g., the presence of ascites, anterior abdominal wall dilated veins or signs associated with liver impairment) or liver imaging/transient elastography suggest the presence of portal hypertension, consideration should be given, in appropriate cases, for an OesophagoGastroDuodenoscopy (OGD) to be performed to rule out occult varices.

Recommendation

- **Spleen assessment by palpation, recorded as cm below the left costal margin, is the most straightforward method but may be limited by body habitus. Ultrasound aids spleen size determination in a more uniform manner, if required (Grade 1C).**
- **Assessment of symptoms using a validated tool e.g., the MPN-SAF TSS (MPN-10) is recommended at baseline, followed by dynamic assessment of symptom burden during follow-up (Grade 1B).**
- **An individualised risk assessment of thrombosis is warranted for all patients, in particular for those with the *JAK2* V617F mutation (Grade 1C).**
- **Where clinical signs or liver imaging/transient elastography suggest the presence of myelofibrosis-related portal hypertension, consideration should be given for an OesophagoGastroDuodenoscopy (OGD) to be performed to rule out occult varices (Grade 2B).**

Establishing a Diagnosis of Myelofibrosis

Patients classically present with progressive anaemia, a leucoerythroblastic blood film with teardrop poikilocytes, splenomegaly and constitutional symptoms. These, along with pathogenic mutations (see below) and typical bone marrow morphological findings, form the basis of the diagnostic criteria proposed by both the World Health Organisation (WHO) and International Consensus Classifications (ICC; **Table 1**)(7,8). Any patient being investigated for potential MPN with suspicious clinical features or atypical peripheral blood (PB) findings (cytopenia, left shifted granulopoiesis, circulating blasts) should proceed to bone marrow (BM) examination, which is essential for the diagnosis(9).

A key morphological finding in the BM is a proliferation of atypical megakaryocytes(7), showing clustering and abnormal localisation with hyperchromatic or bulbous nuclei, lying within an increased reticulin network with focal or diffuse collagen(10). In advanced stages, osteosclerosis can be extensive. To establish a diagnosis and define the disease, reticulin grading is essential with a minimum of grade 2 (0-3 grading system) (11). Commercial reticulin staining kits detect only reticulin fibrosis necessitating additional staining for collagen (trichrome stain)(10–12). Separate scoring systems for collagen fibrosis and osteosclerosis have been recommended (7,12). These may enable more accurate response assessment in patients receiving disease modifying therapies although their clinical impact is yet to be established (12,13). The BM should be reported in a Specialist Integrated Haematological Malignancy Diagnostic Service (SIHMDS) and as per either WHO or ICC criteria. The classification used should be stated in the report and consideration should be given to stating the diagnosis according to both classifications.

It is essential to distinguish MF from other myeloid malignancies and careful morphological assessment for features such as the degree of dysplasia is required(14). In particular, it is important to highlight that systemic mastocytosis can be associated with significant marrow fibrosis. In patients with monocytosis, distinguishing between MF and chronic myelomonocytic leukaemia (CMML) can be challenging though genomic testing with targeted myeloid sequencing panels may help. This will not only detect mutations which are more specific to either disease but will also provide a *JAK2* mutant allele burden which is frequently reported to be higher in PMF than CMML(15,16).

Genetic tests for assessment of patients with MPN have been previously described (17). Suspected PMF cases should be screened for common MPN driver mutations (affecting the *JAK2*, *CALR* and *MPL* genes), on either PB- or BM-derived DNA. Between 50 and 60% of PMF cases are positive for *JAK2* V617F, with the remaining 15-35% and 6-9% of cases testing positive for *CALR* exon 9 or *MPL* exon 10 mutations, respectively(17). Type 1/Type 1-like *CALR* mutations are much more prevalent in PMF than Type 2/Type-2-like mutations (18).

Patients with BM histology and clinical features consistent with PMF or pre-PMF who test negative for *JAK2*, *CALR* or *MPL* mutations should be tested further using a myeloid gene panel and, ideally, karyotyping or genome wide single nucleotide polymorphism (SNP) array(17). Patients without a typical driver mutation in *JAK2*, *MPL* or *CALR* may be diagnosed as triple negative (TN) PMF, although this should prompt careful evaluation of the clinical picture and morphology to exclude the diagnosis of another myeloid neoplasm as highlighted above. In the absence of a clonal marker of disease, causes of secondary fibrosis also require exclusion (**Table 2**)(7,8,19). In

addition, exclusion of *BCR::ABL1* is important for all TN patients with thrombocytosis and/or atypical features.

For patients with a confirmed diagnosis of PMF or post PV/ET-MF, a myeloid panel and karyotyping/SNP array, performed on either PB or BM, provide important prognostic and potentially additional therapeutic target information and are generally recommended for allogeneic haematopoietic cell transplantation (allo-HSCT) candidates. In other patients, including those with pre-PMF, testing may be considered for prognostic purposes and/or if additional genomic data will guide clinical management.

As a minimum, myeloid gene panels should include *ASXL1*, *CBL*, *CSF3R*, *DNMT3A*, *EZH2*, *KIT*, *KRAS*, *IDH1/2*, *NRAS*, *RUNX1*, *SETBP1*, *SF3B1*, *SH2B3*, *SRSF2*, *TET2*, *TP53* and *U2AF1*, together with full coding sequence coverage of *JAK2*, *MPL* and *CALR* exon 9. Broader mutation screens provide additional personalised prognostic information (20–22). All cases should be discussed in a specialist multi-disciplinary meeting. This should be a quorate meeting (e.g., clinical haematologists and representatives of the Specialist Integrated Haematological Malignancy Diagnostic Service etc) as per national/ local guidance on haemato-oncology multi-disciplinary meetings.

Recommendations:

- **All patients with suspected myelofibrosis should undergo a diagnostic bone marrow biopsy and molecular testing for *JAK2*, *CALR* or *MPL* variants as appropriate (Grade 1A).**

- The bone marrow trephine biopsy should have a reticulin stain and grade (Grade 1A) and consideration of trichrome staining. Ideally, the bone marrow should be reported in a Specialist Integrated Haematological Malignancy Diagnostic Service (SIHMDS) and as per either WHO or ICC criteria (Grade 2B).
- A myeloid gene panel, cytogenetic analysis and/or SNP array, and careful morphological examination is recommended for patients with bone marrow histology and clinical features consistent with PMF who test negative for *JAK2*, *CALR* and *MPL* (Grade 1B).
- *BCR::ABL1* should be excluded in cases with persistent thrombocytosis negative for *JAK2*, *CALR* and *MPL* variants, or those with atypical features (Grade 1B).
- Secondary causes of myelofibrosis require exclusion in patients without typical MPN morphology or in those lacking an MPN-associated mutation (Grade 1A).
- Myeloid gene panel testing and conventional karyotyping and/or SNP array are recommended for all patients with PMF, post-PV or post-ET MF who are candidates for allogeneic stem cell transplant, or if it would be useful to guide patient management/ prognostic assessment (Grade 1B).
- All diagnoses should be discussed in a specialist multi-disciplinary meeting (Grade 1B).

Prognostic Evaluation in Primary and post-PV/post-ET Myelofibrosis

Overall survival in MF varies widely and clinicians should be aware of the strengths and limitations of the many prognostic scores available to inform clinical use and guide discussions on therapy and management. These are summarised in **Table 3**, with suggestions where each score may be best utilised.

The International Prognostic Scoring System (IPSS) was the first risk stratification model to consider a large PMF cohort (23). IPSS identified five factors associated with reduced patient survival: age >65yrs, presence of constitutional symptoms (> 10% weight loss in 6 months, night sweats, unexplained fever higher than 37.5°C), haemoglobin <100 g/l, white cell count >25 × 10⁹/l and ≥1% circulating blast cells. Use of the same five factors led to generation of the Dynamic IPSS (DIPSS) score, facilitating dynamic assessment during the disease course. This was further refined in the DIPSS-plus risk stratification by three additional risk factors (unfavourable karyotype, thrombocytopenia (platelets <100 × 10⁹/l) and red cell transfusion dependence)(24,25).

Approximately 40% of patients with PMF have an abnormal karyotype (25–28). Patients with inv(3), -5/5q, -7/7q-, +8, 11q23 and 12p-, i(17q), or complex karyotypes (>2 abnormalities) have significantly poorer outcomes (26,28–30). *JAK2* V617F and *MPL* mutations have been associated with a worse prognosis compared to *CALR* mutations in several studies. Prognostic advantage of a *CALR* mutation may, however, only be confined to Type-1 or Type 1-like mutations (31–34). Overall survival for patients with TN MF appears worse than for those patients with a *JAK2*- or *MPL*-mutation. (31,33). So-called ‘high-molecular risk’ (HMR) pathogenic mutations in five

genes (*ASXL1*, *SRSF2*, *EZH2*, *IDH1* and *IDH2*) have been shown to adversely impact life expectancy and increase the likelihood of leukaemic transformation in MF (35). Patients with >1 HMR mutation have a particularly poor prognosis. Mutations in *TP53*, *U2AF1*, *RUNX1*, *CBL*, *NRAS* and *KRAS* can also confer adverse outcomes (20,22).

The Mutation-enhanced IPSS (MIPSS70+ v2.0) score combines typical haematological features together with karyotype and mutations in “HMR” genes and *U2AF1* Q157(21). MIPSS70+ v2.0 (<http://www.mipss70score.it/>) takes into account varying severity of anaemia. The model included mainly patients <70 years with PMF and pre-PMF and is more accurate than IPSS. Both DIPSS and MIPSS70+ v2.0 appear relevant to those patients eligible for transplant as the risk score correlates with post-transplant outcomes (36,37). The so-called ‘RR6 model’ predicts survival in MF based on clinical response after 6 months of ruxolitinib (considers spleen length reduction, dose density of ruxolitinib and transfusion requirements; (<http://www.rr6.eu/>)) (38). For transplant-eligible patients, the clinical-molecular myelofibrosis transplant scoring system (MTSS) combines age, haematological and molecular parameters, patient fitness and degree of HLA matching to predict survival after allo-HSCT (39). The MYSEC-PM score was developed specifically for patients with post-PV MF and post-ET MF(40).

A personalised prognosis calculator for MPN patients, (Predict blood; <https://blood.predict.nhs.uk/>), takes into account 63 patient demographic, clinical and molecular variables to predict personally-tailored risk for both disease transformation and survival. The model incorporates many more variables than the risk scoring systems described above, does not dichotomise continuous risk variables (such as

increasing age or worsening blood counts), and can predict several different disease outcomes simultaneously. It was shown to provide improved accuracy and greater discrimination over both DIPSS and IPSS, even when incomplete information on molecular variables was available (22).

Prognostication can aid treatment decisions including allo-HSCT. No model can currently predict which patients may benefit from any particular therapy. In general, it is advisable to repeat dynamic prognostication for patients at regular intervals, for example annually, or particularly if there is clinical concern or change in disease phenotype.

Recommendations

- **All patients with myelofibrosis should have a prognostic evaluation performed using one of the currently available validated scores (Grade 1B).**
- **Use of validated risk scores for prognostication can aid treatment decisions, including consideration for allogeneic haemopoietic stem cell transplantation (Grade 1B).**
- **Dynamic prognostic assessment should be performed appropriate to patient characteristics, particularly if there is a change in disease phenotype or loss of response to therapy (Grade 2C)**

Focus on Pre-fibrotic Myelofibrosis

Diagnostic classification and Prognostication

The diagnosis of pre-PMF and its distinction from other MPNs is also based on a combination of clinical, morphological and genomic features (**Table 4**)(7,8). It is important to note that pre-PMF is entirely distinct from low-risk overt MF. Distinction of ET from pre-PMF often causes the most diagnostic difficulty. Compared to ET, patients with pre-PMF tend to have higher white cell and platelet counts, lower haemoglobin levels, higher lactate dehydrogenase (LDH) and greater splenomegaly, and less favourable outcomes: reduced survival, increased leukaemic transformation and increased progression to overt MF (41,42). Pre-PMF tends to have milder clinical features and better survival than overt PMF (43,44).

There is recognised inter-observer variability in distinguishing histological features of pre-PMF and ET (45–49), albeit not fully consistent across studies (50,51). This variability, together with the proportion of patients diagnosed as unclassifiable MPN (47,52) has led to the utility of the WHO criteria being questioned. Although the IPSET thrombosis score from ET has been validated for thrombotic risk in pre-PMF (53), other conventional MF prognostic scores are not fully applicable. Novel prognostic modelling methods have been proposed for pre-PMF including mutational profiles(54). A myeloid gene panel and cytogenetic evaluation is recommended at diagnosis in patients with pre-PMF who are considered to be future allo-HSCT candidates, or where more accurate prognostic information would aid management. Most patients are currently treated pragmatically according to clinical phenotype. There is a risk of thrombosis associated with pre-PMF which must be considered.

351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376

Acknowledgements

All the authors contributed to the writing and review of these guidelines. The writing committee would like to thank Professor Ruben Mesa and Professor Alessandro Vannucchi for their external expert review of this guideline. We would also like to thank the members of MPN Voice who expertly appraised these guidelines and the BSH Haemato-oncology Task Force, the BSH sounding board and the BSH Guidelines Committee for their guidance and expertise.

Conflicts of interest

All authors have made a declaration of interests to the BSH and Task Force Chairs which may be viewed on request.

Disclaimer

While the advice and information in this guidance is believed to be true and accurate at the time of going to press, neither the authors, the BSH nor the publishers accept any legal responsibility for the content of this guidance.

WHO 5th edition(7) ICC(8) 2022 Post-PV/ET MF		WHO 5th edition ICC 2022 Overt PMF	
Major	Documentation of a previous established diagnosis of PV or ET.	Major	Bone marrow biopsy showing megakaryocytic proliferation and atypia, accompanied by reticulin and/or collagen fibrosis grades 2 or 3
	Bone marrow fibrosis of grade 2–3 on a 0–3 scale.		<i>JAK2</i> , <i>CALR</i> or <i>MPL</i> mutation assessed by sensitive technique OR presence of another clonal marker (<i>ASXL1</i> , <i>EZH2</i> , <i>TET2</i> , <i>IDH1</i> , <i>IDH2</i> , <i>SRSF2</i> , and <i>SF3B1</i>) OR absence of minor reactive bone marrow reticulin fibrosis.
			Diagnostic criteria for <i>BCR::ABL1</i> -positive CML, PV, ET, MDS or other myeloid neoplasms not met ^x
Minor	Anaemia and a >20 g/l decrease from baseline haemoglobin concentration. A sustained loss of requirement of either phlebotomy (in the absence of cytoreductive therapy) or cytoreductive treatment for erythrocytosis.	Minor	Anaemia not attributed to a comorbid condition
	Development of any 2 (or all 3) of the following constitutional symptoms: >10% weight loss in 6 months, night sweats, unexplained fever (>37.5 °C)		Leucocytosis $\geq 11 \times 10^9/l$
	Increased palpable splenomegaly >5 cm from the baseline or newly palpable		Splenomegaly detected clinically and/or by imaging
	Elevated LDH (for post-ET MF only).		Elevated LDH
	Leucoerythroblastosis		Leucoerythroblastosis
Diagnosis requires both major criteria and at least 2 minor criteria confirmed in 2 consecutive determinations.		Diagnosis requires all 3 major criteria and at least 1 minor criterion confirmed in 2 consecutive determinations	

Table 1: WHO diagnostic criteria for post-PV/ET myelofibrosis and overt PMF

^xMyeloproliferative neoplasms can be associated with monocytosis or they can develop it during the course of the disease; these cases may mimic chronic myelomonocytic leukaemia (CMML); in these rare instances, a history of MPN excludes CMML, whereas the presence of MPN features in the bone marrow and/or MPN-associated mutations (in *JAK2*, *CALR* or *MPL*) tend to support the diagnosis of MPN with monocytosis rather than CMML.

Causes of secondary bone marrow fibrosis	Examples
Infections	<ul style="list-style-type: none"> • HIV • Visceral leishmaniasis • Tuberculosis • Epstein–Barr virus infection
Autoimmune disorders	<ul style="list-style-type: none"> • Systemic lupus erythematosus • Sjögren syndrome • Antiphospholipid syndrome • Juvenile idiopathic arthritis
Chronic inflammatory condition	
Hairy cell leukaemia	
Other haematological disorders	<ul style="list-style-type: none"> • Myelodysplastic syndromes • Hodgkin lymphoma • Chronic myeloid leukaemia • Some cases of acute myelomonocytic leukemia • Paroxysmal nocturnal haemoglobinuria • Systemic mastocytosis • TAFRO (thrombocytopenia, anasarca, reticulin fibrosis of the bone marrow, renal dysfunction, and organomegaly)
Metastatic malignancy	
Toxic chronic myelopathy	
Treatment with growth factors	<ul style="list-style-type: none"> • Recombinant human thrombopoietin agonists, e.g., romiplostim • IL-11 e.g., oprelvekin
Osseous or other metabolic disease	<ul style="list-style-type: none"> • Vitamin D deficiency • Hyperparathyroidism
Other causes may result in focal fibrosis	<ul style="list-style-type: none"> • Osteonecrosis/myelitis • Bone marrow irradiation • Previous trephine biopsy site • Grey platelet syndrome

Table 2: Potential secondary causes of marrow fibrosis

Prognostic model	Patients (n)	When to use Score	Type of patients included	Variables	Risk Groups (Median overall survival)
IPSS(23)	1054	Newly diagnosed PMF patients	Newly diagnosed PMF	Age >65 y (1) Hb <100 g/l (2) WBC >25 x 10 ⁹ /l (1) Circulating blasts ≥1% (1)	<ul style="list-style-type: none"> Low (0) = 11.3 y Intermediate-1 (1) = 7.9 y Intermediate-2 (2) = 4 y High (3-5) = 2.3 y
DIPSS(24)	525	PMF patients and can be applied at any stage in the disease course	Newly diagnosed PMF and follow up.	Age >65 y (1) Hb <100 g/l (2) WBC >25 x 10 ⁹ /l (1) Circulating blasts ≥1% (1) Constitutional symptoms (1)	<ul style="list-style-type: none"> Low (0) = not reached Intermediate-1 (1-2) = 14.2 y Intermediate-2 (3-4) = 4 y High (>4) = 1.5 y
DIPSS-PLUS(25)	793	PMF patients and can be applied at any stage in the disease course.	Newly diagnosed PMF and follow up.	Age >65 y (1) Hb <100 g/l (2) WBC >25 x 10 ⁹ /l (1) Circulating blasts ≥1% (1) Constitutional symptoms (1) Unfavourable karyotype (1) Red cell transfusion need (1) Platelets <100 x 10 ⁹ /l (1)	<ul style="list-style-type: none"> Low (0) = 15.4 y Intermediate-1 (1-2) = 6.5 y Intermediate-2 (3-4) = 2.9 y High (>4) = 1.3 y
MIPSS 70 v 2.0(21)	406	PMF patients; validated up to the age of 70 years	Patients <70 y (n=311) with PMF. Included prefibrotic MF	VHR karyotype (4) Unfavorable karyotype (3) ≥2 HMR mutations (3) 1 HMR mutation (2) Type 1/like CALR absent (2) Hb <80 g/l females, Hb < 90 g/l Male (2) Hb 80-99 g/l Females, Hb 90-109 g/l Male (1) Circulating blasts ≥2% (1) Constitutional symptoms (2)	<ul style="list-style-type: none"> Very low (0) = Not reached Low (1-2) = 16.4 y Intermediate (3-4) = 7.7 y High (5-8) = 4.1 y Very high (≥9) = 1.8 y
MYSEC-PM(40)	685	Patients with Post-PV and Post ET MF	Post PV and post ET MF	Hb <110 g/l (2) Platelets <150 x 10 ⁹ /l (1) Circulating blasts ≥3% (2) CALR mutation Absent (2) Constitutional symptoms (1) Age (0.15 per yr of age)	<ul style="list-style-type: none"> Low (<11) = Not reached Intermediate-1 (11-13) = 9.3 y Intermediate-2 (14-16) = 4.4 y High (>16) = 2.0 y
MTSS(39)	361	PMF or Post-PV/ ET MF planned for allogeneic stem cell transplantation	Patients presenting for first allogeneic stem cell transplantation. 206 had primary myelofibrosis, 101 had post-ET or post-PV myelofibrosis	Age >57 y (1) WBC >25 x 10 ⁹ /l (1) Platelets <150 x 10 ⁹ /l (1) ASXL1 mutated (1) Karnofsky performance status <90% (1) HLA-mismatched unrelated donor (2) Not CALR/MPL mutated	<ul style="list-style-type: none"> Low (0-2) = 83% 5 y OS Intermediate (3-4) = 64% 5 y OS High (5) = 37% 5yr OS Very high (6-9) = 22% 5 yr OS
Predictblood(22)	2035 MPN 309 MF Validation cohort (515 MPN, 190 MF)	PMF and post-PV/ET MF both at diagnosis and during disease course.	At diagnosis or first referral	Multistate Cox proportional hazards algorithm incorporating 63 clinical and genomic variables to predict risk of survival and disease transformation to myelofibrosis and acute leukaemia	Individualised results for <ul style="list-style-type: none"> Development of MF from ET/PV Development of acute myeloid leukaemia from either chronic phase or any MF (either PMF or secondary MF) Survival in ET/PV, PMF and secondary MF
RR6 Model(38)	209 MF 40 MF in validation cohort	PMF and post PV/ET MF patients, at least 6 months of therapy with RUX and may prompt therapy switch.	PMF and post-ET/PV patients treated with ruxolitinib for at least 6 months	(1) RUX dose <20 mg twice daily at baseline, months 3 and 6 (2) palpable spleen length reduction from baseline ≤30% at months 3 and 6 (3) transfusion need at months 3 and/or 6 (4) transfusion need at all time points (i.e., baseline and months 3 and 6)	Response to RUX after 6 months (RR6), dissected 3 risk categories regarding OS: <ul style="list-style-type: none"> Low (median OS, not reached) Intermediate (median OS, 61 months; 95% CI, 43-80) High (median OS, 33 months; 95% CI, 21-50)

Table 3: Summary of prognostication models validated in patients with myelofibrosis.

PMF=primary myelofibrosis, PV=polycythaemia vera, ET=essential thrombocythaemia, RUX=ruxolitinib, Hb=haemoglobin concentration, WBC=white blood cell count, VHR=very high risk, HMR=high molecular risk, RBC=red blood cell, Human Leucocyte Antigen (HLA), CALR (calreticulin), RUX (ruxolitinib)

WHO 5th edition(7)		ICC 2022(8)	
Major	Megakaryocyte proliferation and atypia, without reticulin fibrosis grade >1, accompanied by increased age-adjusted bone marrow cellularity, granulocytic proliferation and (often) decreased erythropoiesis	Major	Bone marrow biopsy showing megakaryocytic proliferation and atypia, bone marrow fibrosis grade <2, increased age-adjusted BM cellularity, granulocytic proliferation and (often) decreased erythropoiesis
	Diagnostic criteria for <i>BCR::ABL1</i> -positive CML, PV, ET, MDS or other myeloid neoplasms not met		<i>JAK2</i> , <i>CALR</i> or <i>MPL</i> mutation or presence of another clonal marker (assessed by cytogenetic analysis or sensitive NGS techniques, i.e., mutations associated with myeloid neoplasms)
	<i>JAK2</i> , <i>CALR</i> or <i>MPL</i> mutation OR presence of another clonal marker (i.e., mutations associated with other myeloid neoplasms) OR absence of minor reactive bone marrow reticulin fibrosis		Diagnostic criteria for <i>BCR::ABL1</i> -positive CML, PV, ET, MDS or other myeloid neoplasms not met
Minor	Anaemia not attributed to a comorbid condition	Minor	Anaemia not attributed to a comorbid condition
	Leucocytosis $\geq 11 \times 10^9/L$		Leucocytosis $\geq 11 \times 10^9/L$
	Splenomegaly detected clinically and/or by imaging		Palpable splenomegaly
	Elevated LDH		Elevated LDH
	Leucoerythroblastosis		
Diagnosis requires all 3 major criteria and at least 1 minor criterion confirmed in 2 consecutive determinations		Diagnosis requires all 3 major criteria and at least 1 minor criterion confirmed in 2 consecutive determinations	

Table 4: Diagnostic criteria for pre-fibrotic myelofibrosis as per WHO and ICC classifications

BM= bone marrow, PV=polycythaemia vera, ET =essential thrombocythaemia, MDS=myelodysplastic syndromes, LDH =lactate dehydrogenase, NGS=next generation sequencing, WHO=World Health Organization and ICC= International Consensus Classification.

References

1. Reilly JT, McMullin MF, Beer PA, Butt N, Conneally E, Duncombe AS, et al. Use of JAK inhibitors in the management of myelofibrosis: a revision of the British Committee for Standards in Haematology Guidelines for Investigation and Management of Myelofibrosis 2012. *Br J Haematol*. 2014 Nov;167(3):418–20.
2. Titmarsh GJ, Duncombe AS, McMullin MF, O'Rourke M, Mesa R, De Vocht F, et al. How common are myeloproliferative neoplasms? A systematic review and meta-analysis. *Am J Hematol*. 2014 Jun;89(6):581–7.
3. Emanuel RM, Dueck AC, Geyer HL, Kiladjian JJ, Slot S, Zweegman S, et al. Myeloproliferative neoplasm (MPN) symptom assessment form total symptom score: prospective international assessment of an abbreviated symptom burden scoring system among patients with MPNs. *J Clin Oncol Off J Am Soc Clin Oncol*. 2012 Nov 20;30(33):4098–103.
4. Scherber R, Dueck AC, Johansson P, Barbui T, Barosi G, Vannucchi AM, et al. The Myeloproliferative Neoplasm Symptom Assessment Form (MPN-SAF): International Prospective Validation and Reliability Trial in 402 patients. *Blood*. 2011 Jul 14;118(2):401–8.
5. Barbui T, Ghirardi A, Carobbio A, Masciulli A, Carioli G, Rambaldi A, et al. Increased risk of thrombosis in JAK2 V617F-positive patients with primary myelofibrosis and interaction of the mutation with the IPSS score. *Blood Cancer J*. 2022 Nov 16;12(11):156.

6. Yan M, Geyer H, Mesa R, Atallah E, Callum J, Bartoszko J, et al. Clinical Features of Patients With Philadelphia-Negative Myeloproliferative Neoplasms Complicated by Portal Hypertension. *Clin Lymphoma Myeloma Leuk*. 2015 Jan;15(1):e1–5.
7. Khoury JD, Solary E, Abla O, Akkari Y, Alaggio R, Apperley JF, et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. *Leukemia*. 2022 Jul;36(7):1703–19.
8. Arber DA, Orazi A, Hasserjian RP, Borowitz MJ, Calvo KR, Kvasnicka HM, et al. International Consensus Classification of Myeloid Neoplasms and Acute Leukemias: integrating morphologic, clinical, and genomic data. *Blood*. 2022 Sep 15;140(11):1200–28.
9. Passamonti F, Maffioli M. Update from the latest WHO classification of MPNs: a user's manual. *Hematol Am Soc Hematol Educ Program*. 2016 Dec 2;2016(1):534–42.
10. Fujiwara H. Histological evaluation of myeloproliferative neoplasms. *J Clin Exp Hematop JCEH*. 2018;58(2):45–50.
11. Thiele J, Kvasnicka HM, Facchetti F, Franco V, van der Walt J, Orazi A. European consensus on grading bone marrow fibrosis and assessment of cellularity. *Haematologica*. 2005 Aug;90(8):1128–32.
12. Kvasnicka HM, Beham-Schmid C, Bob R, Dirnhofer S, Hussein K, Kreipe H, et al. Problems and pitfalls in grading of bone marrow fibrosis, collagen deposition and

- osteosclerosis - a consensus-based study. *Histopathology*. 2016 May;68(6):905–15.
13. Kröger N, Zabelina T, Alchalby H, Stübig T, Wolschke C, Ayuk F, et al. Dynamic of Bone Marrow Fibrosis Regression Predicts Survival after Allogeneic Stem Cell Transplantation for Myelofibrosis. *Biol Blood Marrow Transplant*. 2014 Jun;20(6):812–5.
14. Tefferi A. Primary myelofibrosis: 2021 update on diagnosis, risk-stratification and management. *Am J Hematol*. 2021 Jan;96(1):145–62.
15. Chapman J, Geyer JT, Khanlari M, Moul A, Casas C, Connor ST, et al. Myeloid neoplasms with features intermediate between primary myelofibrosis and chronic myelomonocytic leukemia. *Mod Pathol Off J U S Can Acad Pathol Inc*. 2018 Mar;31(3):429–41.
16. Hu Z, Ramos CEB, Medeiros LJ, Zhao C, Yin CC, Li S, et al. Utility of JAK2 V617F allelic burden in distinguishing chronic myelomonocytic Leukemia from Primary myelofibrosis with monocytosis. *Hum Pathol*. 2019 Mar;85:290–8.
17. Cross NCP, Godfrey AL, Cargo C, Garg M, Mead AJ, A British Society for Haematology Good Practice Paper. The use of genetic tests to diagnose and manage patients with myeloproliferative and myeloproliferative/myelodysplastic neoplasms, and related disorders. *Br J Haematol*. 2021 Nov;195(3):338–51.
18. Tefferi A, Lasho TL, Finke C, Belachew AA, Wassie EA, Ketterling RP, et al. Type 1 vs type 2 calreticulin mutations in primary myelofibrosis: differences in phenotype and prognostic impact. *Leukemia*. 2014 Jul;28(7):1568–70.

19. Kuter DJ, Bain B, Mufti G, Bagg A, Hasserjian RP. Bone marrow fibrosis: pathophysiology and clinical significance of increased bone marrow stromal fibres. *Br J Haematol*. 2007 Nov;139(3):351–62.
20. Luque Paz D, Riou J, Verger E, Cassinat B, Chauveau A, Ianotto JC, et al. Genomic analysis of primary and secondary myelofibrosis redefines the prognostic impact of *ASXL1* mutations: a FIM study. *Blood Adv*. 2021 Mar 9;5(5):1442–51.
21. Tefferi A, Guglielmelli P, Lasho TL, Gangat N, Ketterling RP, Pardanani A, et al. MIPSS70+ Version 2.0: Mutation and Karyotype-Enhanced International Prognostic Scoring System for Primary Myelofibrosis. *J Clin Oncol Off J Am Soc Clin Oncol*. 2018;36(17):1769–70.
22. Grinfeld J, Nangalia J, Baxter EJ, Wedge DC, Angelopoulos N, Cantrill R, et al. Classification and Personalized Prognosis in Myeloproliferative Neoplasms. *N Engl J Med*. 2018 Oct 11;379(15):1416–30.
23. Cervantes F, Dupriez B, Pereira A, Passamonti F, Reilly JT, Morra E, et al. New prognostic scoring system for primary myelofibrosis based on a study of the International Working Group for Myelofibrosis Research and Treatment. *Blood*. 2009 Mar 26;113(13):2895–901.
24. Passamonti F, Cervantes F, Vannucchi AM, Morra E, Rumi E, Pereira A, et al. A dynamic prognostic model to predict survival in primary myelofibrosis: a study by the IWG-MRT (International Working Group for Myeloproliferative Neoplasms Research and Treatment). *Blood*. 2010 Mar 4;115(9):1703–8.

25. Gangat N, Caramazza D, Vaidya R, George G, Begna K, Schwager S, et al.
DIPSS Plus: A Refined Dynamic International Prognostic Scoring System for Primary Myelofibrosis That Incorporates Prognostic Information From Karyotype, Platelet Count, and Transfusion Status. *J Clin Oncol*. 2011 Feb 1;29(4):392–7.
26. Hussein K, Huang J, Lasho T, Pardanani A, Mesa RA, Williamson CM, et al.
Karyotype complements the International Prognostic Scoring System for primary myelofibrosis. *Eur J Haematol*. 2009 Apr;82(4):255–9.
27. Tefferi A, Guglielmelli P, Nicolosi M, Mannelli F, Mudireddy M, Bartalucci N, et al.
GIPSS: genetically inspired prognostic scoring system for primary myelofibrosis. *Leukemia*. 2018 Jul;32(7):1631–42.
28. Tefferi A, Nicolosi M, Mudireddy M, Lasho TL, Gangat N, Begna KH, et al.
Revised cytogenetic risk stratification in primary myelofibrosis: analysis based on 1002 informative patients. *Leukemia*. 2018 May;32(5):1189–99.
29. Tam CS, Abruzzo LV, Lin KI, Cortes J, Lynn A, Keating MJ, et al. The role of cytogenetic abnormalities as a prognostic marker in primary myelofibrosis: applicability at the time of diagnosis and later during disease course. *Blood*. 2009 Apr 30;113(18):4171–8.
30. Caramazza D, Begna KH, Gangat N, Vaidya R, Siragusa S, Van Dyke DL, et al.
Refined cytogenetic-risk categorization for overall and leukemia-free survival in primary myelofibrosis: a single center study of 433 patients. *Leukemia*. 2011 Jan;25(1):82–8.

31. Rumi E, Pietra D, Pascutto C, Guglielmelli P, Martínez-Trillos A, Casetti I, et al. Clinical effect of driver mutations of JAK2, CALR, or MPL in primary myelofibrosis. *Blood*. 2014 Aug 14;124(7):1062–9.
32. Tefferi A, Lasho TL, Tischer A, Wassie EA, Finke CM, Belachew AA, et al. The prognostic advantage of calreticulin mutations in myelofibrosis might be confined to type 1 or type 1-like CALR variants. *Blood*. 2014 Oct 9;124(15):2465–6.
33. Tefferi A, Lasho TL, Finke CM, Knudson RA, Ketterling R, Hanson CH, et al. CALR vs JAK2 vs MPL-mutated or triple-negative myelofibrosis: clinical, cytogenetic and molecular comparisons. *Leukemia*. 2014 Jul;28(7):1472–7.
34. Tefferi A, Lasho TL, Finke C, Belachew AA, Wassie EA, Ketterling RP, et al. Type 1 vs type 2 calreticulin mutations in primary myelofibrosis: differences in phenotype and prognostic impact. *Leukemia*. 2014 Jul;28(7):1568–70.
35. Vannucchi AM, Lasho TL, Guglielmelli P, Biamonte F, Pardanani A, Pereira A, et al. Mutations and prognosis in primary myelofibrosis. *Leukemia*. 2013 Sep;27(9):1861–9.
36. Kröger N, Giorgino T, Scott BL, Ditschkowski M, Alchalby H, Cervantes F, et al. Impact of allogeneic stem cell transplantation on survival of patients less than 65 years of age with primary myelofibrosis. *Blood*. 2015 May 21;125(21):3347–50.
37. Ali H, Aldoss I, Yang D, Mokhtari S, Khaled S, Aribi A, et al. MIPSS70+ v2.0 predicts long-term survival in myelofibrosis after allogeneic HCT with the Flu/Mel conditioning regimen. *Blood Adv*. 2019 Jan 8;3(1):83–95.

38. Maffioli M, Mora B, Ball S, Iurlo A, Elli EM, Finazzi MC, et al. A prognostic model to predict survival after 6 months of ruxolitinib in patients with myelofibrosis. *Blood Adv.* 2022 Mar 22;6(6):1855–64.
39. Gagelmann N, Ditschkowski M, Bogdanov R, Bredin S, Robin M, Cassinat B, et al. Comprehensive clinical-molecular transplant scoring system for myelofibrosis undergoing stem cell transplantation. *Blood.* 2019 May 16;133(20):2233–42.
40. Passamonti F, Giorgino T, Mora B, Guglielmelli P, Rumi E, Maffioli M, et al. A clinical-molecular prognostic model to predict survival in patients with post polycythemia vera and post essential thrombocythemia myelofibrosis. *Leukemia.* 2017 Dec;31(12):2726–31.
41. Rumi E, Boveri E, Bellini M, Pietra D, Ferretti VV, Sant'Antonio E, et al. Clinical course and outcome of essential thrombocythemia and prefibrotic myelofibrosis according to the revised WHO 2016 diagnostic criteria. *Oncotarget.* 2017 Nov 24;8(60):101735–44.
42. Barbui T, Thiele J, Passamonti F, Rumi E, Boveri E, Ruggeri M, et al. Survival and disease progression in essential thrombocythemia are significantly influenced by accurate morphologic diagnosis: an international study. *J Clin Oncol.* 2011 Aug 10;29(23):3179–84.
43. Guglielmelli P, Pacilli A, Rotunno G, Rumi E, Rosti V, Delaini F, et al. Presentation and outcome of patients with 2016 WHO diagnosis of prefibrotic and overt primary myelofibrosis. *Blood.* 2017 Jun 15;129(24):3227–36.

44. Mudireddy M, Shah S, Lasho T, Barraco D, Hanson CA, Ketterling RP, et al. Prefibrotic versus overtly fibrotic primary myelofibrosis: clinical, cytogenetic, molecular and prognostic comparisons. *Br J Haematol*. 2018 Aug;182(4):594–7.
45. Wilkins BS, Erber WN, Bareford D, Buck G, Wheatley K, East CL, et al. Bone marrow pathology in essential thrombocythemia: interobserver reliability and utility for identifying disease subtypes. *Blood*. 2008 Jan 1;111(1):60–70.
46. Brousseau M, Parot-Schinkel E, Moles MP, Boyer F, Hunault M, Rousselet MC. Practical application and clinical impact of the WHO histopathological criteria on bone marrow biopsy for the diagnosis of essential thrombocythemia versus prefibrotic primary myelofibrosis. *Histopathology*. 2010 May;56(6):758–67.
47. Buhr T, Hebeda K, Kaloutsi V, Porwit A, Van der Walt J, Kreipe H. European Bone Marrow Working Group trial on reproducibility of World Health Organization criteria to discriminate essential thrombocythemia from prefibrotic primary myelofibrosis. *Haematologica*. 2012 Mar;97(3):360–5.
48. Alvarez-Larran A, Ancochea A, Garcia M, Climent F, Garcia-Pallarols F, Angona A, et al. WHO-histological criteria for myeloproliferative neoplasms: reproducibility, diagnostic accuracy and correlation with gene mutations and clinical outcomes. *Br J Haematol*. 2014 Sep;166(6):911–9.
49. Madelung AB, Bondo H, Stamp I, Loevgreen P, Nielsen SL, Falensteen A, et al. World Health Organization-defined classification of myeloproliferative neoplasms: morphological reproducibility and clinical correlations--the Danish experience. *Am J Hematol*. 2013 Dec;88(12):1012–6.

50. Thiele J, Kvasnicka HM, Müllauer L, Buxhofer-Ausch V, Gisslinger B, Gisslinger H. Essential thrombocythemia versus early primary myelofibrosis: a multicenter study to validate the WHO classification. *Blood*. 2011 May 26;117(21):5710–8.
51. Gianelli U, Bossi A, Cortinovis I, Sabattini E, Tripodo C, Boveri E, et al. Reproducibility of the WHO histological criteria for the diagnosis of Philadelphia chromosome-negative myeloproliferative neoplasms. *Mod Pathol Off J U S Can Acad Pathol Inc*. 2014 Jun;27(6):814–22.
52. Ochiai T, Yasuda H, Araki M, Misawa K, Morishita S, Nudejima M, et al. The 2014 BCSH criteria and the 2016 WHO criteria for essential thrombocythemia: A comparison in a large-scale cohort. *Eur J Haematol*. 2018 Jun;100(6):544–9.
53. Guglielmelli P, Carobbio A, Rumi E, De Stefano V, Mannelli L, Mannelli F, et al. Validation of the IPSET score for thrombosis in patients with prefibrotic myelofibrosis. *Blood Cancer J*. 2020 Feb 25;10(2):21.
54. Carobbio A, Guglielmelli P, Rumi E, Cavalloni C, De Stefano V, Betti S, et al. A multistate model of survival prediction and event monitoring in prefibrotic myelofibrosis. *Blood Cancer J*. 2020 Oct 14;10(10):100.