



The neurocognitive correlates of non-substance addictive behaviors

Erynn Christensen^{a,*}, Lucy Albertella^a, Samuel R. Chamberlain^{b,c}, Maja Brydevall^a, Chao Suo^a, Jon E. Grant^e, Murat Yücel^{a,d}, Rico Sze Chun Lee^{a,f}

^a BrainPark, Turner Institute for Brain and Mental Health, Monash University, Melbourne, VIC, Australia

^b Department of Psychiatry, University of Southampton, Southampton, the United Kingdom of Great Britain and Northern Ireland

^c Southern Health NHS Foundation Trust, Southampton, the United Kingdom of Great Britain and Northern Ireland

^d QIMR Berghofer Medical Research Institute, Herston, QLD, Australia

^e Department of Psychiatry & Behavioral Neuroscience, University of Chicago, Pritzker School of Medicine, Chicago, IL, USA

^f Melbourne School of Psychological Sciences, University of Melbourne, Melbourne, VIC, Australia

ARTICLE INFO

Keywords:

Addiction
Cognition
Neurocognition
Alcohol
Internet use
Pornography

ABSTRACT

Neurocognitive deficits have been implicated as transdiagnostic risk markers of substance use disorders. However, these have yet to be comprehensively evaluated in other, non-substance addictions. In a large, general community sample (N = 475) the present study evaluated the neurocognitive correlates of problem alcohol use and three non-substance-related addictive behaviors: addictive eating (AE), problematic pornography use (PPU), and problematic use of the internet (PUI), to identify potential shared and distinct neurocognitive correlates. A sample of Australian residents (54.4 % female M[SD] age = 32.4[11.9] years) completed a comprehensive online assessment of neurocognitive tasks tapping into eight distinct expert-endorsed domains purportedly associated with addiction. Multiple linear regressions with bootstrapping were used to examine associations among each addictive behavior of interest and neurocognition, trait impulsivity, and compulsivity, as well as key covariates. Neurocognition was differentially associated with each addictive behavior. None of the neurocognitive domains were significantly associated with problematic alcohol use or AE ($p > .05$), poorer performance monitoring was significantly associated with higher levels of PPU and PUI ($\beta = -0.10, p = .049$; $\beta = -0.09, p = .028$), and a preference for delayed gratification was associated with more severe PUI ($\beta = -0.10, p = .025$). Our findings have theoretical implications for how we understand non-substance addiction and suggest the need for a more nuanced approach to studying addictive behaviors that take into account the underlying neurocognitive mechanisms associated with each type of addiction.

1. Introduction

Addiction is a complex condition characterized by repeated engagement in substance use or other behavior despite negative consequences (Diagnostic Statistical Manual of Mental Disorders [DSM-5]: American Psychiatric Association, 2013). The literature has predominantly centered on substance use disorders, such as alcohol use disorder (AUD), whereas there has been an increasing focus on non-substance (behavioral) addictions (Chamberlain et al., 2016). Recent such efforts include addictive eating (AE), problematic pornography use (PPU), and problematic use of the internet (PUI) due to their associated psychological distress and poorer quality of life (Burmeister et al., 2013; Burrows et al., 2018; Camilleri et al., 2021; Fineberg et al., 2018; Floros & Ioannidis, 2021; Kuss et al., 2014; Raj et al., 2022). It is essential to

acknowledge that none of these addictive behaviors have agreed-upon clinical criteria. Consequently, it is more appropriate to view them as problem behaviors positioned on a severity continuum. AE, PPU, and PUI may share some important features with substance addiction, including loss of control, craving, and emotional distress (Adams et al., 2017; Fineberg et al., 2018; Grubbs et al., 2015; Tiego et al., 2021). Addictive behaviors can be conceptualized using current neurocognitive ‘dual-process’ models of addiction, typified by excessive drive and reward-seeking coupled with impaired executive ‘top-down’ control (Volkow et al., 2019; Volkow & Morales, 2015). Dual process models have been further extended in non-substance addictive behaviors (Brand, 2022; Brand et al., 2019; Wei et al., 2017). However, addiction-specific neurocognitive functions have yet to be comprehensively evaluated in non-substance addictions. For example, it is unclear whether

* Corresponding author at: Monash Biomedical Imaging, 770 Blackburn Rd, Clayton, VIC, 3168, Australia.

E-mail address: erynnchristensen@gmail.com (E. Christensen).

<https://doi.org/10.1016/j.addbeh.2023.107904>

Received 2 August 2023; Received in revised form 19 October 2023; Accepted 2 November 2023

Available online 7 November 2023

0306-4603/© 2023 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

non-substance addictive behaviors share neurocognitive features. Substance use research has also shown inconsistent neurocognitive correlates (Ekhtiari et al., 2017; Lundqvist, 2005; Smith et al., 2014). Identifying the neurocognitive correlates of addictive behaviors is essential to progressing etiological understanding and facilitating the development of effective prevention and intervention strategies. This study investigates the *trans*-behavioral neurocognitive correlates of non-substance addictive behaviors; namely, AE, PPU, and PUI, in comparison to problematic alcohol use, to identify potential shared and distinct neurocognitive mechanisms.

Current consensus is that some neurocognitive deficits can serve as transdiagnostic markers of addiction, regardless of the specific type of behavior (Yücel et al., 2019). Self-regulation is often defined as being comprised of: *Inhibition*, the ability to stop or inhibit an undesirable automatic response (Chambers et al., 2009); *Shifting*, the ability to flexibly shift between tasks, mental sets, or demands (Kiesel et al., 2010); and *Performance monitoring*, the ability to monitor and evaluate action outcomes to adjust performance (Franken et al., 2018). Poorer executive functioning in individuals with addiction is marked by reduced inhibitory control, poorer performance monitoring, and inflexible task shifting (Franken et al., 2018; Odlaug et al., 2011; Rodrigue et al., 2018; Smith et al., 2014; Zhou et al., 2013). By contrast, the 'bottom-up' motivational processes associated with addiction are underpinned by heightened reward-seeking; chief among this is an increased preference for immediate over larger but delayed rewards or poor 'delay discounting' (Amlung et al., 2017), and heightened incentive attribution towards addiction-related cues (Berridge & Robinson, 2016). Unfortunately, most studies to date have focused on individual neurocognitive factors in isolation. Therefore, there is a lack of empirical evidence regarding how each domain is independently associated with addictive behaviors. This is a critical area of further investigation since neurocognitive functions associated with executive control and heightened reward-seeking significantly overlap with each other (Criaud & Boulinguez, 2013; Ridderinkhof et al., 2004).

Few studies have examined neurocognitive functioning associated with AE, PPU, and PUI. Existing studies have generally identified neurocognitive markers implicated in problematic alcohol use, including heightened attentional bias toward reward cues (Adams et al., 2019; Jeromin et al., 2016; Mechelmans et al., 2014; Nikolaidou et al., 2019) and a preference for immediate gratification over larger, later rewards (Cheng et al., 2021; Kowalewska et al., 2017; VanderBroek-Stice et al., 2017). Individuals with AE, PPU, and PUI also present with executive dysfunction, but in differing domains. PUI but neither AE or PPU has been shown to be associated with impaired response inhibition (Antons & Brand, 2018; Antons & Matthias, 2020; Hardee et al., 2020; Ioannidis et al., 2022; Meule et al., 2012; VanderBroek-Stice et al., 2017), and both AE and PUI have been shown to be associated with impaired performance monitoring (Franken et al., 2018; Rodrigue et al., 2018; Zhou et al., 2013). It is unclear whether these differences arise from methodological differences, for example selection of neurocognitive measures, sample characteristics, and inclusion (or lack thereof) of covariates. Because many studies only look at a single neurocognitive measure, it is entirely plausible that differing executive control profiles among AE, PPU, and PUI may be due to common neurocognitive function(s) that drive performance across measures. Further, sample characteristics often vary, including differing illness severities, and demographic features (e.g. sex, age). These studies also do not address comorbidities of other addictive behaviors. Assessing the relationship between neurocognition and multiple addictive behaviors in the same sample would account for these disparities, and help us evaluate whether AE, PPU, and PUI share common underlying neurocognitive processes.

Examining addictive behaviors in general community samples using a dimensional approach can lead to a more nuanced understanding of the neurocognitive functions underlying addictive behaviors. Clinical presentations of addiction represent only the tip of the iceberg, as the

majority of addiction problems do not meet strict diagnostic thresholds (Grant et al., 2015; Hasin et al., 2013). Further, despite not meeting diagnostic thresholds, these addictive behaviors still contribute to burden of disease and are important to detect early. By adopting a dimensional approach and examining addictive behaviors across the full spectrum of severities, we can detect often more subtle effects, with direct implications for the development of early intervention strategies.

In addition to neurocognition, addictive behaviors are associated with trait impulsivity and compulsivity (Forsén Mantilla et al., 2022; Kuss et al., 2014; Murphy et al., 2014). Impulsivity can be defined as the tendency to rapidly react to a situation in a reward-driven manner, without forethought or consideration of the consequences (Moeller et al., 2001). Impulsivity is a multifaceted construct, commonly conceptualized and assessed in addiction research as being made up of four key facets: lack of planning, lack of perseverance, sensation seeking and emotion-driven rash action (urgency; Cyders et al., 2014). Compulsivity can be defined as repeated actions that are inappropriate to a given situation and lacks a clear connection to an overarching goal, often resulting in negative consequences (Dalley et al., 2011). Although previous research has found neurocognitive functions correspond with trait impulsivity (Christiansen et al., 2012) and compulsivity (Albertella et al., 2020), when assessed concurrently it is clear that they are not measuring the same underlying construct (Eisenberg et al., 2019). For example, convergent validity among task-related executive function and self-report questionnaires was only moderate in a large meta-analysis on self-regulation ability (Duckworth & Kern, 2011). Therefore, both trait-based and neurocognitive-based measures of impulsivity and compulsivity appear to contribute to addiction vulnerability in different ways.

This study sought to investigate the neurocognitive correlates of addictive behaviors. Using a tailored assessment battery that measures a comprehensive range of addiction-specific neurocognitive functions, the present study aims to evaluate the shared and distinct neurocognitive correlates of problematic alcohol use, AE, PPU, and PUI, adjusting for key covariates. A demographically targeted, general community sample was recruited to capture a broad spectrum of behavioral severity.

2. Methods

2.1. Participants and procedure

This study was embedded in a larger normative study for the BrainPark Assessment of Cognition (BrainPAC) neurocognitive battery (see [supplementary material](#)). Australian residents were recruited via Prolific, social media advertisements, and local community newsletters. Inclusion criteria were: 18 to 65 years old, not color blind, self-reported absence of a neurological disorder (i.e. stroke, brain injury, and dementia) or history of a psychotic disorder. Participants completed all measures online via the Qualtrics survey platform (<https://www.qualtrics.com>). The neurocognitive tasks were separated by self-report surveys (trait and behavior scales) and the order of task presentation was counterbalanced. The study was approved by the Monash University Human Research Ethics Committee [26088].

2.2. Neurocognitive measures

The neurocognitive battery (Table 1) was selected to measure expert-endorsed neurocognitive domains as associated with addiction and addiction-related outcomes (Yücel et al., 2019).

2.3. Self-report scales

Trait Compulsivity – The Cambridge-Chicago Compulsivity Trait Scale (CHI-T; Chamberlain & Grant, 2018; Tiego et al., 2023) is a 15-item scale. Items are summed into an overall score. Higher scores indicate higher compulsivity.

Trait Impulsivity – The Short UPPS Impulsive Behavior Scale (SUPPS-

Table 1
Neurocognitive assessment battery.

Function	Task	Brief description	Trials (practice)	Primary metric
Response inhibition	The BrainPAC Stop Signal Taks (SST) (Lee et al., 2023)	A gamified visual cue stop signal paradigm (Verbruggen et al., 2019) in a medieval war game format, with the goal of defeating a dragon. Players pass arrows to two archers (left/ right) as fast as they can (go signal) whilst avoiding the dragon's fire (stop signal) so they can shoot the arrows and defend their village.	150 (10)	Stop signal reaction time (SSRT). Higher RT indicates poorer response inhibition.
Reward learning (reward-related attentional bias)	The BrainPAC Value Modulated Attentional Capture (VMAC) Task (Lee et al., 2023)	A gamified version of the original VMAC task (Albertella et al., 2019; Le Pelley et al., 2015) following a soccer format. Players must kick the ball (left/right), with speed and accuracy to earn points. Some trials have players with different hair colors acting as distractors and indicating the potential reward value of that trial.	5x24 (6)	VMAC score averaged across the last two blocks of the task. Higher values reflect more reward-related attentional capture.
Reward learning (goal-directed vs habitual)	The BrainPAC Sequential Decision-Making Task (SDT) (Lee et al., 2023)	A gamified two-step choice task (Kool et al., 2016, 2017), presented in the form of an animal rescue game. Participants select a ranger to search two environments (the forest or the farmland) for lost animals. Model-based decision-makers learn which rangers are most effective at finding the maximum number of animals.	125 (25)	Mixing weight (w). Higher scores indicate more goal-directed (model-based) decision-making.
Reward valuation (risky decision-making under uncertainty)	The BrainPAC Balloon Analogue Risk Task (BART) (Lee et al., 2023)	A gamified version of the BART stretch variant (Lejuez et al., 2002) paradigm in which players inflate a series of balloons to earn hypothetical money (maximum earning of \$128 AUD p/ balloon). Each balloon has a pseudorandomized burst threshold (mean burst point at \$64 AUD), resulting in any potential earnings being lost.	30 (10)	Mean pre-committed pumps across all balloons. Higher values indicate riskier choice in the face of uncertainty.
Flexible updating	N-back Task (Ragland et al., 2002; Inquisit 5, 2018)	A letter sequencing go/no-go task. Participants respond to "M" in 0-back trials, to the previous letter in 1-back trials, and to the letter two trials back in 2-back trials, and so on.	3x60 (9)	3-back d' (parametric measure of sensitivity). Higher d' values indicate more flexible updating/ better working memory performance.
Goal selection; updating, representation and maintenance	Category Switch Task (CST) (Friedman et al., 2008; Inquisit 5, 2018)	Participants are presented with a word they must categorize in terms of A) 'living', or B) 'size'. Each trial is accompanied by a cue that indicates to the participant whether they are required to categorize the object according to conditions A or B.	65 (80)	Latency switch cost. Higher values indicate poorer task switching.
Performance monitoring	Error Awareness Task (EAT) (Hester et al., 2007; Inquisit 5, 2018)	A visual go/no-go paradigm in which participants indicate their error awareness following any commission error.	2x150 (70)	Percentage error awareness (commission errors). Higher values indicate better error awareness.
Temporal discounting	Monetary Choice Questionnaire (MCQ) (Kirby et al., 1999)	A 27-item questionnaire asks the participant to choose between two hypothetical reward options, a smaller reward now, or a larger reward at some point in the future e.g. "Would you prefer \$15 today or \$35 in 13 days".	27 (NA)	Log k . Higher values indicate a preference for sooner but smaller rewards.

P; Cyders et al., 2014) is a 20-item scale. Three scores were calculated: urgency (combining negative and positive urgency subscales), lack of perseverance and premeditation (combining lack of perseverance and premeditation subscales), and sensation seeking (SS: sensation seeking subscale). Higher values indicate greater impulsivity.

Psychological Distress – The Depression Anxiety Stress Scale (DASS-21; Szabó, 2010) is a 21-item scale used to assess current psychological distress. Items were summed into form a total score. Higher scores indicate greater distress.

2.4. Dependent variables

Participants were asked to respond reflecting on the past three months.

Problematic Alcohol Use – The Alcohol Use Disorder Identification Test (AUDIT; Saunders et al., 1993) is a 10-item scale. A total score is summed, ranging from 0 to 40.

Addictive Eating – The modified Yale Food Addiction Scale version 2 (mYFAS 2.0; Schulte & Gearhardt, 2017) is a 13-item scale. A symptom count is computed, ranging from 0 to 11.

Problematic Pornography Use (PPU) - The Problematic Pornography Consumption Scale (PPCS-6; Bóthe et al., 2021) is a 6-item scale. A total score is calculated, ranging from 6 to 42.

Problematic Use of the Internet – The abbreviated Young's Internet Addiction Test (IAT-10; Tiego et al., 2021) is a 10-item scale. A total score ranging from 10 to 50 is computed.

2.5. Data cleaning

To ensure data quality, online neurocognitive assessment requires comprehensive screening protocols and post-hoc data cleaning. Bots and fraudulent responses were identified and excluded using features embedded in the Qualtrics survey platform. Implausible responses and poor performance presumably due to lack of effort were identified and removed via a) attention check questions (e.g. "Please select the option *Piano Keys*"); b) neurocognitive task performance at less than chance levels (as per Lee et al., 2023; Albertella, Watson, et al., 2019); c) task-specific cleaning procedures (e.g. SST go trial accuracy, stop trial accuracy, [Verbruggen et al., 2019] and Independent Race Model check [Band et al., 2003]). Mann Whitney U and t-tests were used to investigate differences between individuals whose data was filtered out compared with included individuals on all variables (Table A1).

2.6. Data analysis

Individuals who reported not engaging in an addictive behavior in the past 3 months were assigned a zero for the corresponding problem behavior scale. Statistical outliers on neurocognitive measures ≥ 3 standard deviations from the mean were removed (Field et al., 2012). All analyses were conducted on complete data sets (i.e. participants who provided data for all variables of interest). Bivariate Spearman correlations, adjusting for multiple comparisons (Holm method: Holm, 1979), investigated relationships among all variables (Table 3). The

distributions of all four addictive behavior scales were positively skewed, constituting the choice of linear regression models with bootstrapping (5,000 samples) (Neal & Simons, 2007). Multicollinearity was assessed for each model independently, with VIF values less than 2.5 indicating no issue of multicollinearity (Johnston et al., 2018). Age, sex, psychological distress, trait impulsivity, and compulsivity were included as covariates in each regression model (Eisenberg et al., 2019; Sjoberg & Cole, 2018; Starcke et al., 2016). G-Power 3.1 (Faul et al., 2007) was used to calculate the minimum sample size required for multiple regression analyses with 15 predictors and an alpha error probability set to 0.05. A sample of $N = 139$ was deemed sufficient to find a medium effect ($f^2 = 0.15$, Power = 0.80).

3. Results

3.1. Participants

Nine-hundred-and-forty-four participants were enrolled. Sixty-three withdrew during the assessment sessions, and 78 were identified as fraudulent responses (i.e. spam bots/not genuine responses). Eight-hundred-and-three participants completed the assessment, with 311 removed due to missing data on one or more variables of interest (i.e. those included in the regression models), failed attention checks, or poor neurocognitive task performance. After removing outliers based on neurocognitive task performance, a final sample of 475 individuals had complete datasets (see Fig. 1). Participant demographics are displayed in Table 2.

3.2. Multiple regression models

Correlation analysis findings are presented in Table 3. The multivariate models (Tables 4-7) showed the neurocognitive variables were not significantly associated with problematic alcohol use or AE. Poorer error awareness was significantly associated with greater PPU ($\beta = -0.10$, $p = .049$) and PUI ($\beta = -0.09$, $p = .028$), and less steep delay discounting was significantly associated with higher PUI ($\beta = -0.10$, $p = .025$). Higher levels of psychological distress were significantly associated with more problematic alcohol use ($\beta = 0.22$, $p = .003$), AE ($\beta = 0.26$, $p < .001$), PPU ($\beta = 0.20$, $p < .001$), and PUI ($\beta = 0.37$, $p < .001$). Higher levels of urgency were significantly associated with more PPU (β

$= 0.14$, $p = .013$) and PUI ($\beta = 0.15$, $p = .003$). Sensation seeking was positively associated with problematic alcohol use ($\beta = 0.17$, $p = .003$) and negatively associated with AE ($\beta = -0.12$, $p = .004$) and PUI ($\beta = -0.09$, $p = .014$). Higher levels of trait compulsivity were significantly associated with more AE ($\beta = 0.18$, $p = .002$) and PUI ($\beta = 0.12$, $p = .010$). Age was positively associated with problematic alcohol use ($\beta = 0.16$, $p = .002$), and negatively associated with PUI ($\beta = -0.17$, $p < .001$). Being male was significantly associated with more PPU ($\beta = -0.46$, $p < .001$), and PUI ($\beta = -0.15$, $p < .001$) whilst being female was significantly associated with greater AE ($\beta = 0.19$, $p < .001$). None of the multivariate models showed problems with multicollinearity.

4. Discussion

This is the first study to comprehensively investigate and control for the neurocognitive correlates of non-substance addictive behaviors across AE, PPU, PUI, and problem alcohol use. We took a dimensional approach to identify correlates associated with these addictive behaviors at varying degrees of severity in the general community. Our findings indicate, in this sample, addictive behaviors are associated with a unique profile of neurocognitive functioning. Our multivariate models showed none of the neurocognitive domains were associated with AE or problematic alcohol use. Poorer performance monitoring was independently associated with more PPU and PUI, and a higher preference for delayed gratification was also independently associated with higher PUI. Importantly, these findings are identified whilst adjusting for known confounds (i.e. age, sex, psychological distress), as identified as a key methodological shortcoming of prior research (Christensen et al., 2023). Our findings suggest the need for a more nuanced approach to studying addictive behaviors that take into account the underlying neurocognitive mechanisms associated with each type of addiction.

Our performance monitoring findings are consistent with prior research (Zhou et al., 2013). However, this is the first study to show a link between poorer performance monitoring and PPU. Very little work has been done evaluating the potential neurocognitive mechanisms associated with PPU. Of the extant work, findings are mixed. For example, PPU has been associated with both improved and poorer inhibitory control (Antons & Brand, 2018; Antons & Matthias, 2020). Considerable research has investigated PUI, with studies showing PUI is associated with poorer working memory, response inhibition, and risk-




Fig. 1. Flow diagram mapping data collection, cleaning, and reasons for exclusion.

Table 2
Description of demographic characteristics of the sample.

Variables	N	
Sample	475	
Mean age (SD)	32.3 (11.8)	
Sex, N (%)		
Male	217 (45.7)	
Female	258 (54.3)	
Gender, N (%)		
Man	216 (45.5)	
Woman	254 (53.5)	
Non-binary	4 (0.8)	
Not listed/ Prefer not to say	1 (0.2)	
Race/Ethnicity, N		
Aboriginal or Torres Strait Islander	2	
African	2	
Asian	101	
Black or African American	2	
Hispanic or Latino	4	
Middle Eastern	4	
South Asian	30	
White or Caucasian	317	
Other	13	
Household income in AUD, N		
< \$10,000	15	
\$10,000 – \$20,000	17	
\$20,000 – \$40,000	48	
\$40,000 – \$60,000	74	
\$60,000 – \$80,000	58	
\$80,000 – \$100,000	66	
> \$100,000	197	
Addictive behaviors	M (SD), range	Classified as problematic: N (%)
AUDIT	5.09 (4.82), 0–30	Harmful/hazardous: 47 (10) Suspected dependence: 15 (3)
mYFAS	0.71 (1.78), 0–10	Mild: 28 (6) Moderate: 15 (3) Severe: 22 (5)
PPCS	9.61 (5.89), 6–42	31 (7)
IAT	16.31 (6.67), 10–48	163 (34)

Note: Sex was defined as biological sex. AUDIT: Alcohol Use Identification Test, no/low problem use (0–7), harmful/ hazardous use (8–14), likely alcohol dependence (≥ 15); mYFAS: modified Yale Food Addiction Scale 2.0, mild (2–3), moderate (4–5) severe (≥ 6) symptoms. PPCS: Problematic Pornography Consumption Scale, problematic use (≥ 20). IAT: an abbreviated version of Young's Internet Addiction Test, problematic use (≥ 17).

taking behaviors, as measured by the BART (for a meta-analysis: Ioannidis et al., 2019). Our study did not find these associations, potentially because the previous studies compared individuals with a PUI “diagnosis” to controls, while our sample included individuals with varying PUI severity. Further, if such deficits are related to vulnerability, they may be more strongly associated with presence vs absence of a disorder rather than its severity. Relatedly, it may be that deficits in these domains are evident in those with high levels of PUI, whereas the current study was conducted in a relatively normative sample. Performance monitoring is necessary to support higher-order functions such as cognitive control (Ferdinand & Czernochowski, 2018), and so reduced performance monitoring may be detectable before other cognitive functions, acting as an early *trans*-behavioral risk indicator for PPU and PUI. However, it is important to note the concurrence of PUI and PPU in our sample (Table 3). Given pornography is predominantly accessed via the internet, our PPU findings may instead be driven by problematic internet use to watch pornography, rather than specifically pornography-related functions.

Some of our findings diverge from that of previous work. For instance, the finding that more severe PUI was associated with less steep delay discounting, or a preference for later larger rewards, is contrary to substantial literature showing individuals with internet addiction are significantly steeper discounters compared to controls (for meta-

analysis: Cheng et al., 2021). A potential interpretation relates to the fact that we are looking at PUI dimensionally, thus including individuals at both low and high levels of problem severity. Tiego et al. (2021) have argued that PUI as measured by the IAT has a unipolar distribution in which meaningful variance is found at the higher end of the severity spectrum. Given this, including individuals at lower PUI severity may have disrupted the expected effect. Further, the finding that the severity of AE symptoms was not associated with any neurocognitive variables contrasts with previous studies (Franken et al., 2018; Rodrigue et al., 2018; VanderBroek-Stice et al., 2017). One explanation for this may be that neurocognitive functions are more likely to be implicated in more severe AE. The aforementioned studies included samples with 35–100 % of participants endorsing mild-severe food addiction, compared to 18 % currently. Similarly, contrary to expectations, the present study did not find any of the neurocognitive variables were significantly associated with problematic alcohol use. The majority of our sample had no/low alcohol use problems, making it difficult to compare with previous work which has mostly been conducted in hazardous-severe alcohol use cohorts (Stavro et al., 2013). It is possible that neurocognitive risk or sequelae may not be detectable at mild levels of severity. However, our findings may be attributed to the study sample, with PUI being the most prevalent addictive behavior endorsed. This sample may differ significantly from traditional alcohol-focused research cohorts, suggesting a unique group of individuals in this study.

Psychological distress was a *trans*-behavioral correlate across all four addictive behaviors. This is consistent with previous research showing increased psychological distress is associated with multiple addictive and compulsive behaviors (Albertella et al., 2021; Albertella, Pelley, et al., 2019; Sepas et al., 2021). This finding is likely to present a bidirectional relationship between psychological distress and addictive behaviors, a) that addictive behaviors are motivated by way of coping with psychological distress (Burnatowska et al., 2022; Rodriguez et al., 2020), and b) addictive behaviors may themselves enhance distress (Yang et al., 2022).

The present study suggests that different problematic behaviors may be associated with differing trait impulsivity and compulsivity thresholds. Higher SS was significantly associated with more problematic alcohol use, and less AE and PUI. Our alcohol findings are in keeping with the literature in that SS is consistently associated with higher levels of alcohol use, albeit with a small effect (Hittner & Swickert, 2006). Further, our AE findings replicate that of Burrows et al. (2017) who showed a negative relationship between SS and food addiction. Less SS has also been linked with more time spent online (Müller et al., 2016). We also found AE and PUI were significantly positively associated with trait compulsivity, which is in line with previous findings (Albertella et al., 2021). Taken together we see a pattern emerge according to behavior type: after controlling for key covariates, substance-related addictive behaviors (i.e. alcohol use) may be more driven by the desire for sensation, presumably the reinforcing effects of the substance; whilst non-substance addictive behaviors may be more compulsively driven. It is important to note that this does not refute the role of impulsivity in non-substance-related addictive behaviors, nor compulsivity in substance addiction both of which have been well evidenced elsewhere (Everitt & Robbins, 2016; Lee et al., 2019). Rather, the present findings may speak to the relative contributions of these constructs per addictive behavior type (Tiego et al., 2019), particularly in less severe, non-clinical general population cohorts.

4.1. Limitations and future directions

Despite the strengths of our study, namely, a large sample size, the assessment of multiple addictive behaviors, and a comprehensive evaluation of addiction-specific neurocognitive correlates for each behavior, our findings should be considered in light of several limitations. The most notable limitation is the relatively low levels of severity (see appendix Fig. A1) in the study sample preventing the generalizability of

Table 3
Spearman correlations corrected for multiple comparisons.

Variable	M	SD	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	
1. mYFAS	5.09	4.82	-																			
2. PPCS	9.61	5.89	0.00	-																		
3. IAT	16.31	6.67	0.29***	0.37***	-																	
4. AUDIT	0.71	1.78	0.07	0.10	-0.02	-																
5. SST: SSRT	0.01	0.05	-0.02	0.04	-0.17*	-0.04	-															
6. VMAC: VMAC score	57.29	16.51	-0.08	0.11	0.01	-0.06	-0.06	-														
7. BART: M pre-committed pumps	0.30	0.20	0.01	0.00	-0.03	-0.05	0.13	0.03	0.01	0.04	-											
8. CST: Switch cost latency	-4.60	1.67	0.01	0.08	-0.06	0.03	-0.04	0.03	0.01	0.04	0.04	-										
9. DDT: Log k	67.00	34.88	-0.03	-0.04	-0.02	0.04	-0.14	-0.04	0.02	-0.09	-0.15	-										
10. EAT: Error awareness	0.51	0.36	-0.07	0.10	-0.01	-0.03	-0.11	0.05	0.00	-0.03	-0.18*	0.15	-									
11. SDT: w	2.15	1.16	-0.07	0.02	-0.02	0.06	-0.04	0.03	0.02	0.05	-0.12	0.19*	0.14	-								
12. N-Back: 3-back d'	7.50	1.76	0.02	0.04	0.13	0.12	-0.10	0.02	0.06	-0.12	0.13	-0.04	-0.14	-0.04	-							
13. SUPPS-P: Lack of perseverance and premeditation	8.42	2.45	0.20***	0.23***	0.34***	0.08	-0.03	0.01	-0.04	0.01	0.12	-0.10	-0.07	-0.07	0.32***	-						
14. SUPPS-P: Urgency	9.41	2.79	-0.13	0.14	-0.01	0.25***	0.01	-0.04	0.17*	-0.04	0.13	-0.06	-0.05	0.04	0.11	0.25***	-					
15. SUPPS-P: Sensation seeking	27.23	6.19	0.27***	0.11	0.33***	-0.03	-0.05	0.03	0.02	0.02	0.02	0.02	-0.03	-0.03	-0.28***	0.34***	-0.02	-				
16. CHI-T: Trait compulsivity score	12.39	11.36	0.35***	0.15	0.53***	0.15	-0.08	-0.07	-0.08	0.00	-0.03	0.03	-0.01	-0.04	0.01	0.35***	0.34***	-0.02	-			
17. DASS: Total score	32.32	11.85	-0.06	-0.21***	-0.39***	0.00	0.34***	-0.01	0.03	0.21***	-0.06	-0.09	-0.04	-0.06	-0.17*	-0.25***	-0.09	-0.17*	-0.27***	-		
18. Age	-	-	0.25***	-0.52***	-0.08	0.00	0.09	-0.01	-0.18*	-0.03	-0.11	0.02	-0.08	-0.11	0.03	-0.08	-0.24***	0.02	0.12	0.03	-	
19. Sex (F)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Note: *p* values were adjusted for multiple comparisons using Holm method; **p* < .05, ****p* < .001, *****p* < .0001.

Table 4

Linear regression model with bootstrapping for problematic alcohol use.

	β	SE	95 % CI		<i>p</i>
			Lower	Upper	
Demographics					
Age	0.16	0.02	0.02	0.10	0.002**
Sex (F)	-0.01	0.51	-1.09	0.94	0.921
Neurocognition					
SST: SSRT	-0.06	2.87	-8.39	3.14	0.332
VMAC: VMAC score	0.01	4.10	-7.55	9.05	0.912
BART: M pre-committed pumps	0.04	0.02	-0.02	0.04	0.458
CST: Switch cost latency	-0.02	0.00	-0.00	0.00	0.666
EAT: Error awareness	0.03	0.01	-0.01	0.02	0.606
SDT: w	-0.07	0.56	-2.05	0.15	0.088
N-Back: 3-back d'	0.04	0.15	-0.15	0.45	0.291
DDT: Log k	-0.00	0.12	-0.24	0.21	0.948
Covariates					
DASS: Total score	0.22	0.03	0.03	0.15	0.003**
SUPPS-P: Lack of perseverance and premeditation	0.04	0.16	-0.20	0.40	0.475
SUPPS-P: Urgency	0.05	0.11	-0.14	0.31	0.440
SUPPS-P: Sensation seeking	0.17	0.09	0.11	0.44	0.003**
CHI-T: Trait compulsivity score	-0.05	0.05	-0.13	0.06	0.503

Note. β : Unstandardized coefficient; SE: Standard error; **p* < .05, ***p* < .01, *****p* < .0001.

Table 5

Linear regression model with bootstrapping for addictive eating.

	β	SE	95 % CI		<i>p</i>
			Lower	Upper	
Demographics					
Age	0.05	0.01	-0.01	0.02	0.310
Sex (F)	0.19	0.14	0.41	0.96	<0.001***
Neurocognition					
SST: SSRT	0.06	0.93	-0.79	2.94	0.257
VMAC: VMAC score	0.02	1.54	-2.40	3.77	0.698
BART: M pre-committed pumps	0.04	0.00	-0.00	0.01	0.350
CST: Switch cost latency	0.02	0.00	-0.00	0.00	0.714
EAT: Error awareness	0.01	0.00	-0.00	0.00	0.918
SDT: w	-0.03	0.23	-0.58	0.32	0.579
N-Back: 3-back d'	0.03	0.06	-0.06	0.16	0.391
DDT: Log k	0.07	0.05	-0.01	0.17	0.109
Covariates					
DASS: Total score	0.26	0.01	0.02	0.06	<0.001***
SUPPS-P: Lack of perseverance and premeditation	0.03	0.06	-0.08	0.14	0.621
SUPPS-P: Urgency	0.08	0.05	-0.03	0.14	0.195
SUPPS-P: Sensation seeking	-0.12	0.03	-0.14	-0.02	0.004**
CHI-T: Trait compulsivity score	0.18	0.02	0.02	0.09	0.002**

Note. β : Unstandardized coefficient; SE: Standard error; **p* < .05, ***p* < .01, *****p* < .0001.

our findings to populations with more severe addictive behaviors. Although rates of AE and PPU corresponded with what has previously been estimated in general population samples (4–15 % for AE and 7 % for PPU; Mennig et al., 2020; Meule & Gearhardt, 2019), rate of hazardous alcohol use was well below Australian population estimates (22 %; O'Brien et al., 2020). We recommend that future research in community samples focus recruitment efforts to target individuals at more severe levels of addictive behavior. Contrary to this, the rate of PUI in the study sample was much higher than global estimates (6 %; Cheng & Li, 2014), suggesting our sample experienced more PUI than what would typically be observed in the general population. An additional limitation was the unsupervised nature of data collection. While online remote-access data collection is beneficial when wanting to target demographically diverse samples at scale, it also requires a rigorous data cleaning protocol which necessitated the removal of just under 40 % of the

Table 6
Linear regression model with bootstrapping for problematic pornography use.

	β	SE	95 % CI		p
			Lower	Upper	
Demographics					
Age	-0.04	0.02	-0.06	0.03	0.450
Sex (F)	-0.46	0.53	-6.47	-4.40	<0.001***
Neurocognition					
SST: SSRT	-0.04	2.26	-6.71	0.97	0.287
VMAC: VMAC score	0.03	4.09	-3.54	12.20	0.291
BART: M pre-committed pumps	0.01	0.01	-0.03	0.03	0.883
CST: Switch cost latency	0.00	0.00	-0.00	0.00	0.924
EAT: Error awareness	-0.10	0.01	-0.03	-0.00	0.049*
SDT: w	0.03	0.67	-0.75	1.88	0.400
N-Back: 3-back d'	-0.05	0.24	-0.70	0.21	0.298
DDT: Log k	-0.05	0.16	-0.49	0.15	0.327
Covariates					
DASS: Total score	0.20	0.03	0.04	0.16	<0.001***
SUPPS-P: Lack of perseverance and premeditation	0.04	0.20	-0.26	0.53	0.581
SUPPS-P: Urgency	0.14	0.13	0.06	0.59	0.013*
SUPPS-P: Sensation seeking	-0.03	0.09	-0.24	0.13	0.542
CHI-T: Trait compulsivity score	0.05	0.06	0.06	0.16	0.414

Note. β : Unstandardized coefficient; SE: Standard error; * $p < .05$, ** $p < .01$, *** $p < .001$.

Table 7
Linear regression model with bootstrapping for problematic use of the internet.

	β	SE	95 % CI		p
			Lower	Upper	
Demographics					
Age	-0.17	0.02	-0.14	-0.05	<0.001***
Sex (F)	-0.15	0.54	-3.07	-0.98	<0.001***
Neurocognition					
SST: SSRT	0.01	2.90	-4.81	6.37	0.801
VMAC: VMAC score	-0.00	5.13	-10.66	9.10	0.886
BART: M pre-committed pumps	0.03	0.02	-0.02	0.04	0.444
CST: Switch cost latency	0.02	0.00	-0.00	0.00	0.597
EAT: Error awareness	-0.09	0.01	-0.03	-0.00	0.028*
SDT: w	-0.07	0.76	-2.81	0.22	0.093
N-Back: 3-back d'	-0.02	0.22	-0.55	0.31	0.555
DDT: Log k	-0.10	0.18	-0.75	-0.04	0.025*
Covariates					
DASS: Total score	0.37	0.03	0.15	0.29	<0.001***
SUPPS-P: Lack of perseverance and premeditation	0.08	0.19	-0.06	0.71	0.102
SUPPS-P: Urgency	0.15	0.14	0.13	0.67	0.003**
SUPPS-P: Sensation seeking	-0.09	0.09	-0.40	-0.04	0.014*
CHI-T: Trait compulsivity score	0.12	0.05	0.03	0.24	0.010*

Note. β : Standardized coefficient; SE: Standard error; * $p < .05$, ** $p < .01$, *** $p < .001$.

dataset. Post-hoc comparisons of our sample for those whose data was screened out found that the individuals who were removed from the analysis had significantly higher AE, PUI, and lack of perseverance/premeditation than included participants (see appendix). As such, our sample may be biased towards less impulsive individuals which may not be representative of the wider community. Future research should consider the trade-off associated with unsupervised data collection.

The use of gamified neurocognitive tasks to assess aspects of neurocognitive function may also have impacted our findings. Whilst gamification can enhance engagement and motivation (Lumsden et al., 2016), the addition of game-like elements did cause some of the tasks to deviate from their traditional counterparts. For instance, enhanced complexity of the visual display which may affect the salience of visual

cues, potentially impacting paradigms that rely on distractor cues such as the VMAC task. Further, the BrainPAC SST utilized a points element in which faster responses earned more points, potentially encouraging faster but less accurate responding and thus impacting SSRT calculations.

Finally, the cross-sectional nature of the present study limits our ability to determine causal relationships between the variables of interest. The next natural step would be to conduct a longitudinal evaluation of the same predictors to determine the key mechanisms that predict the development of AE, PPU, and PUI over time. This would better inform intervention and prevention targets for non-substance addictive behaviors and could shed light on the relative contribution of impulsivity and compulsivity (in terms of cognition or traits) at different stages of addiction (e.g. as relates to duration of addictive symptoms).

In conclusion, the present study revealed that different addictive behaviors may have unique neurocognitive mechanisms. There are likely partly distinct mechanisms or pathways to addiction depending on the addictive behavior in question. This has key implications for early intervention, in particular, our study supports the need for tailored treatments that focus on the specific behavior-related neurocognitive functions rather than assuming cognitive dysfunction is necessarily the same across addictions.

CRedit authorship contribution statement

Erynn Christensen: Conceptualization, Methodology, Software, Formal analysis, Investigation, Data curation, Writing – original draft, Writing – review & editing. **Lucy Albertella:** Conceptualization, Supervision, Writing – review & editing. **Samuel R. Chamberlain:** Writing – review & editing. **Maja Brydevall:** Software, Writing – review & editing. **Chao Suo:** Software. **Jon E. Grant:** Writing – review & editing. **Murat Yücel:** Funding acquisition, Supervision, Writing – review & editing. **Rico Sze Chun Lee:** Funding acquisition, Conceptualization, Supervision, Writing – review & editing.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgement

Funding for this study was provided by the National Health and Medical Research Council (NHMRC) Project Grant [APP1162031] and NHMRC Medical Research Future Fund Investigator Grant [APP1193946]. This research was also supported by an Australian Government Research Training Program (RTP) Scholarship. Author disclosures are as follows: LA and MY have received funding from the David Winston Turner Endowment Fund. MY and RSCL have received funding from the NHMRC. MY has received funding from the Australian Research Council (ARC), Australian Defence Science and Technology (DST), the Department of Industry, Innovation and Science (DIIS), the National Institutes of Health (NIH, USA), the Wilson Foundation, and sponsored Investigator-Initiated trials including Incannex Healthcare Ltd. MY has also received payments in relation to court-, expert witness-, and/or expert review-reports. MY also sits on the Advisory Boards of: Centre of The Urban Mental Health, University of Amsterdam; Monash Biomedical Imaging Centre; and Enosis Therapeutics. JEG has received research grant support from NIDA, and from Biohaven, Janssen, and Boehringer Ingelheim Pharmaceuticals. He receives yearly

compensation from Springer Publishing for acting as Editor-in-Chief of the Journal of Gambling Studies and has received royalties from Oxford University Press, American Psychiatric Publishing, Inc., Norton Press, and McGraw Hill. SRC and LA receive honoraria for editorial work at

Elsevier. The above funding sources have had no role in the present study design, collection, analysis or interpretation of the data, writing the manuscript, or the decision to submit this paper for publication.

Appendix

Table A1

Variables that significantly differed between the sample retained in the study and the sample removed during data cleaning.

Variable	Mean(SD)		Test statistic
	Sample retained	Sample removed	
mYFAS symptom count	0.72 (1.79)	2.39 (2.95)	47366***
IAT score	16.3 (6.68)	22.46 (8.86)	45768***
Lack of perseverance and premeditation	7.52 (1.77)	8.84 (2.19)	6.02***

Note: *** $p < .001$; Mann-Whitney U test was used for non-parametric comparisons (mYFAS and IAT); t -test was used for parametric comparison (lack of perseverance and premeditation).




Fig. A1. Histograms of addictive behaviors in the sample.

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.addbeh.2023.107904>.

References

- Adams, R. C., Lawrence, N. S., Verbruggen, F., & Chambers, C. D. (2017). Training response inhibition to reduce food consumption: Mechanisms, stimulus specificity and appropriate training protocols. *Appetite*, *109*, 11–23. <https://doi.org/10.1016/j.appet.2016.11.014>
- Adams, R. C., Sedgmond, J., Maizey, L., Chambers, C. D., & Lawrence, N. S. (2019). Food addiction: Implications for the diagnosis and treatment of overeating. *Nutrients*, *11*(9), 2086. <https://doi.org/10.3390/nu11092086>
- Albertella, L., Chamberlain, S. R., Le Pelley, M. E., Greenwood, L.-M., Lee, R. S., Den Ouden, L., Segrave, R. A., Grant, J. E., & Yücel, M. (2020). Compulsivity is measurable across distinct psychiatric symptom domains and is associated with familial risk and reward-related attentional capture. *CNS Spectrums*, *25*(4), 519–526. <https://doi.org/10.1017/S1092852919001330>
- Albertella, L., Pelley, M. E. L., Chamberlain, S. R., Westbrook, F., Fontenelle, L. F., Segrave, R., Lee, R., Pearson, D., & Yücel, M. (2019). Reward-related attentional capture is associated with severity of addictive and obsessive-compulsive behaviors. *Psychology of Addictive Behaviors*, *33*(5), 495–502. <https://doi.org/10.1037/adb0000484>
- Albertella, L., Rotaru, K., Christensen, E., Lowe, A., Brierley, M.-E., Richardson, K., Chamberlain, S. R., Lee, R. S. C., Kayayan, E., Grant, J. E., Schluter-Hughes, S., Ince, C., Fontenelle, L. F., Segrave, R., & Yücel, M. (2021). The influence of trait compulsivity and impulsivity on addictive and compulsive behaviors during COVID-19. *Frontiers in Psychiatry*, *12*, Article 634583. <https://doi.org/10.3389/fpsy.2021.634583>
- Albertella, L., Watson, P., Yücel, M., & Le Pelley, M. E. (2019). Persistence of value-modulated attentional capture is associated with risky alcohol use. *Addictive Behaviors Reports*, *10*, Article 100195. <https://doi.org/10.1016/j.abrep.2019.100195>
- American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders (5th ed.). Washington, DC: Author.
- Amlung, M., Vedelago, L., Acker, J., Balodis, I., & MacKillop, J. (2017). Steep delay discounting and addictive behavior: A meta-analysis of continuous associations. *Addiction (Abingdon, England)*, *112*(1), 51–62. <https://doi.org/10.1111/add.13535>
- Antons, S., & Brand, M. (2018). Trait and state impulsivity in males with tendency towards Internet-pornography-use disorder. *Addictive Behaviors*, *79*, 171–177. <https://doi.org/10.1016/j.addbeh.2017.12.029>
- Antons, S., & Matthias, B. (2020). Inhibitory control and problematic Internet-pornography use – The important balancing role of the insula. *Journal of Behavioral Addictions*, *9*(1), 58–70. <https://doi.org/10.1556/2006.2020.00010>
- Band, G. P. H., Van Der Molen, M. W., & Logan, G. D. (2003). Horse-race model simulations of the stop-signal procedure. *Acta Psychologica*, *112*(2), 105–142. [https://doi.org/10.1016/S0001-6918\(02\)00079-3](https://doi.org/10.1016/S0001-6918(02)00079-3)
- Berridge, K. C., & Robinson, T. E. (2016). Liking, wanting, and the incentive-sensitization theory of addiction. *American Psychologist*, *71*(8), 670–679. <https://doi.org/10.1037/amp0000059>

- Bóthe, B., Tóth-Király, I., Demetrovics, Z., & Orosz, G. (2021). The short version of the problematic pornography consumption scale (PPCS-6): A reliable and valid measure in general and treatment-seeking populations. *Journal of Sex Research*, 58(3), 342–352. <https://doi.org/10.1080/00224499.2020.1716205>
- Brand, M. (2022). Can internet use become addictive? *Science*, 376(6595), 798–799. <https://doi.org/10.1126/science.abn4189>
- Brand, M., Wegmann, E., Stark, R., Müller, A., Wölfling, K., Robbins, T. W., & Potenza, M. N. (2019). The Interaction of Person-Affect-Cognition-Execution (I-PACE) model for addictive behaviors: Update, generalization to addictive behaviors beyond internet-use disorders, and specification of the process character of addictive behaviors. *Neuroscience & Biobehavioral Reviews*, 104, 1–10. <https://doi.org/10.1016/j.neubiorev.2019.06.032>
- Burmeister, J. M., Hinman, N., Koball, A., Hoffmann, D. A., & Carels, R. A. (2013). Food addiction in adults seeking weight loss treatment. Implications for psychosocial health and weight loss. *Appetite*, 60, 103–110. <https://doi.org/10.1016/j.appet.2012.09.013>
- Burnatowska, E., Surma, S., & Olszanecka-Glinianowicz, M. (2022). Relationship between mental health and emotional eating during the COVID-19 pandemic: A systematic review. *Nutrients*, 14(19), 3989. <https://doi.org/10.3390/nu14193989>
- Burrows, T., Hides, L., Brown, R., Dayas, C., & Kay-Lambkin, F. (2017). Differences in dietary preferences, personality and mental health in Australian adults with and without food addiction. *Nutrients*, 9(3), 285. <https://doi.org/10.3390/nu9030285>
- Burrows, T., Kay-Lambkin, F., Pursey, K., Skinner, J., & Dayas, C. (2018). Food addiction and associations with mental health symptoms: A systematic review with meta-analysis. *Journal of Human Nutrition and Dietetics*, 31(4), 544–572. <https://doi.org/10.1111/jhn.12532>
- Camilleri, C., Perry, J. T., & Sammut, S. (2021). Compulsive internet pornography use and mental health: A cross-sectional study in a sample of university students in the United States. *Frontiers in Psychology*, 11, Article 613244. <https://doi.org/10.3389/fpsyg.2020.613244>
- Chamberlain, S. R., & Grant, J. E. (2018). Initial validation of a transdiagnostic compulsivity questionnaire: The cambridge-chicago compulsivity trait scale. *CNS Spectrums*, 23(5), 340–346. <https://doi.org/10.1017/S1092852918000810>
- Chamberlain, S. R., Lochner, C., Stein, D. J., Goudriaan, A. E., van Holst, R. J., Zohar, J., & Grant, J. E. (2016). Behavioural addiction—A rising tide? *European Neuropsychopharmacology*, 26(5), 841–855. <https://doi.org/10.1016/j.euroneuro.2015.08.013>
- Chambers, C. D., Garavan, H., & Bellgrove, M. A. (2009). Insights into the neural basis of response inhibition from cognitive and clinical neuroscience. *Neuroscience & Biobehavioral Reviews*, 33(5), 631–646. <https://doi.org/10.1016/j.neubiorev.2008.08.016>
- Cheng, C., & Li, A. Y. (2014). Internet addiction prevalence and quality of (real) life: A meta-analysis of 31 nations across seven world regions. *Cyberpsychology, Behavior, and Social Networking*, 17(12), 755–760. <https://doi.org/10.1089/cyber.2014.0317>
- Cheng, Y. S., Ko, H.-C., Sun, C.-K., & Yeh, P.-Y. (2021). The relationship between delay discounting and Internet addiction: A systematic review and meta-analysis. *Addictive Behaviors*, 114, Article 106751. <https://doi.org/10.1016/j.addbeh.2020.106751>
- Christensen, E., Brydevall, M., Albertella, L., Samarawickrama, S. K., Yücel, M., & Lee, R. S. C. (2023). Neurocognitive predictors of addiction-related outcomes: A systematic review of longitudinal studies. *Neuroscience & Biobehavioral Reviews*, 152, Article 105295. <https://doi.org/10.1016/j.neubiorev.2023.105295>
- Christiansen, P., Cole, J. C., Goudie, A. J., & Field, M. (2012). Components of behavioural impulsivity and automatic cue approach predict unique variance in hazardous drinking. *Psychopharmacology*, 219(2), 501–510. <https://doi.org/10.1007/s00213-011-2396-z>
- Criaud, M., & Boulinguez, P. (2013). Have we been asking the right questions when assessing response inhibition in go/no-go tasks with fMRI? A meta-analysis and critical review. *Neuroscience & Biobehavioral Reviews*, 37(1), 11–23. <https://doi.org/10.1016/j.neubiorev.2012.11.003>
- Cyders, M. A., Littlefield, A. K., Coffey, S., & Karyadi, K. A. (2014). Examination of a short english version of the UPPS-P impulsive behavior scale. *Addictive Behaviors*, 39(9), 1372–1376. <https://doi.org/10.1016/j.addbeh.2014.02.013>
- Dalley, J. W., Everitt, B. J., & Robbins, T. W. (2011). Impulsivity, compulsivity, and top-down cognitive control. *Neuron*, 69(4), 680–694. <https://doi.org/10.1016/j.neuron.2011.01.020>
- Duckworth, A. L., & Kern, M. L. (2011). A meta-analysis of the convergent validity of self-control measures. *Journal of Research in Personality*, 45(3), 259–268. <https://doi.org/10.1016/j.jrp.2011.02.004>
- Eisenberg, I. W., Bissett, P. G., Enkavi, A. Z., Li, J., MacKinnon, D. P., Marsch, L. A., & Poldrack, R. A. (2019). Uncovering the structure of self-regulation through data-driven ontology discovery. *Nature Communications*, 10(1). <https://doi.org/10.1038/s41467-019-10301-1>
- Ekhtiari, H., Victor, T. A., & Paulus, M. P. (2017). Aberrant decision-making and drug addiction—How strong is the evidence? *Current Opinion in Behavioral Sciences*, 13, 25–33. <https://doi.org/10.1016/j.cobeha.2016.09.002>
- Everitt, B. J., & Robbins, T. W. (2016). Drug addiction: Updating actions to habits to compulsions ten years on. *Annual Review of Psychology*, 67, 23–50. <https://doi.org/10.1146/annurev-psych-122414-033457>
- Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: A flexible statistical power analysis for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, 39, 175–191.
- Ferdinand, N. K., & Czernochowski, D. (2018). Motivational influences on performance monitoring and cognitive control across the adult lifespan. *Frontiers in Psychology*, 9, 1018. <https://doi.org/10.3389/fpsyg.2018.01018>
- Field, A., Miles, J., & Field, Z. (2012). *Discovering statistics using R*. SAGE Publications.
- Fineberg, N., Demetrovics, Z., Stein, D., Ioannidis, K., Potenza, M., Grünblatt, E., ... Chamberlain, S. (2018). Manifesto for a European research network into Problematic Usage of the Internet. *European Neuropsychopharmacology*, 28(11), 1232–1246. <https://doi.org/10.1016/j.euroneuro.2018.08.004>
- Floros, G. D., & Ioannidis, K. (2021). Editorial: The impact of online addiction on general health, well-being and associated societal costs. *Frontiers in Public Health*, 9, Article 676498. <https://doi.org/10.3389/fpubh.2021.676498>
- Forsén Mantilla, E., Clinton, D., Monell, E., Levallius, J., & Birgegård, A. (2022). Impulsivity and compulsivity as parallel mediators of emotion dysregulation in eating-related addictive-like behaviors, alcohol use, and compulsive exercise. *Brain and Behavior*, 12(1). <https://doi.org/10.1002/brb3.2458>
- Franken, I. H. A., Nijs, I. M. T., Toes, A., & van der Veen, F. M. (2018). Food addiction is associated with impaired performance monitoring. *Biological Psychology*, 131, 49–53. <https://doi.org/10.1016/j.biopsycho.2016.07.005>
- Friedman, N. P., Miyake, A., Young, S. E., DeFries, J. C., Corley, R. P., & Hewitt, J. K. (2008). Individual differences in executive functions are almost entirely genetic in origin. *Journal of Experimental Psychology: General*, 137(2), 201–225. <https://doi.org/10.1037/0096-3445.137.2.201>
- Grant, B. F., Goldstein, R. B., Saha, T. D., Chou, S. P., Jung, J., Zhang, H., Pickering, R. P., Ruan, W. J., Smith, S. M., Huang, B., & Hasin, D. S. (2015). Epidemiology of DSM-5 alcohol use disorder: Results from the national epidemiologic survey on alcohol and related conditions III. *JAMA Psychiatry*, 72(8), 757. <https://doi.org/10.1001/jamapsychiatry.2015.0584>
- Grubbs, J. B., Stauner, N., Exline, J. J., Pargament, K. I., & Lindberg, M. J. (2015). Perceived addiction to internet pornography and psychological distress: Examining relationships concurrently and over time. *Psychology of Addictive Behaviors*, 29(4), 1056–1067. <https://doi.org/10.1037/adb0000114>
- Hardee, J. E., Phaneuf, C., Cope, L., Zucker, R., Gearhardt, A., & Heitzeg, M. (2020). Neural correlates of inhibitory control in youth with symptoms of food addiction. *Appetite*, 148, Article 104578. <https://doi.org/10.1016/j.appet.2019.104578>
- Hasin, D. S., O'Brien, C. P., Auriacombe, M., Borges, G., Bucholz, K., Budney, A., Compton, W. M., Crowley, T., Ling, W., Petry, N. M., Schuckit, M., & Grant, B. F. (2013). DSM-5 criteria for substance use disorders: Recommendations and rationale. *American Journal of Psychiatry*, 170(8), 834–851. <https://doi.org/10.1176/appi.ajp.2013.12060782>
- Hester, R., Simões-Franklin, C., & Garavan, H. (2007). Post-error behavior in active cocaine users: Poor awareness of errors in the presence of intact performance adjustments. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*. <https://doi.org/10.1038/sj.npp.1301326>
- Hittner, J. B., & Swickert, R. (2006). Sensation seeking and alcohol use: A meta-analytic review. *Addictive Behaviors*, 31(8), 1383–1401. <https://doi.org/10.1016/j.addbeh.2005.11.004>
- Holm, S. (1979). A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, 65–70.
- Inquisit 5 [Computer software]. (2018). Retrieved from <https://www.millisecond.com>.
- Ioannidis, K., Grant, J. E., & Chamberlain, S. R. (2022). Problematic usage of the internet and cognition. *Current Opinion in Behavioral Sciences*, 44, Article 101104. <https://doi.org/10.1016/j.cobeha.2022.101104>
- Jeromin, F., Nyenhuis, N., & Barke, A. (2016). Attentional bias in excessive Internet gamers: Experimental investigations using an addiction Stroop and a visual probe. *Journal of Behavioral Addictions*, 5(1), 32–40. <https://doi.org/10.1556/2006.5.2016.012>
- Johnston, R., Jones, K., & Manley, D. (2018). Confounding and collinearity in regression analysis: A cautionary tale and an alternative procedure, illustrated by studies of British voting behaviour. *Quality & Quantity*, 52(4), 1957–1976. <https://doi.org/10.1007/s11355-017-0584-6>
- Kiesel, A., Steinhauser, M., Wendt, M., Falkenstein, M., Jost, K., Philipp, A. M., & Koch, I. (2010). Control and interference in task switching—A review. *Psychological Bulletin*, 136(5), 849–874. <https://doi.org/10.1037/a0019842>
- Kirby, K. N., Petry, N. M., & Bickel, W. K. (1999). Heroin addicts have higher discount rates for delayed rewards than non-drug-using controls. *Journal of Experimental Psychology: General*, 128(1), 78–87. <https://doi.org/10.1037/0096-3445.128.1.78>
- Kowalewska, E., Sadowski, J., Wordecha, M., Golec, K., Czajkowski, M., & Mateusz, G. (2017). OP-53: How impulsivity is related to problematic pornography use? Longitudinal study among participants of 12-steps sexual addiction treatment program. *Journal of Behavioral Addictions*, 6(S1), 26. Gale Academic OneFile.
- Kuss, D., Griffiths, M., Karila, L., & Billieux, J. (2014). Internet Addiction: A systematic review of epidemiological research for the last decade. *Current Pharmaceutical Design*, 20(25), 4026–4052. <https://doi.org/10.2174/13816128113199990617>
- Lee, R. S. C., Albertella, L., Christensen, E., Suo, C., Segrave, R. A., Brydevall, M., ... Yücel, M. (2023). A novel, expert-endorsed, neurocognitive digital assessment tool for addictive disorders: Development and validation study. *Journal of Medical Internet Research*, 25, Article e44414. <https://doi.org/10.2196/44414>
- Lee, R. S. C., Hoppenbrouwers, S., & Franken, I. (2019). A systematic meta-review of impulsivity and compulsivity in addictive behaviors. *Neuropsychology Review*, 29, 14–26. <https://doi.org/10.1007/s11065-019-09402-x>
- Lumsden, J., Edwards, E. A., Lawrence, N. S., Coyle, D., & Munafò, M. R. (2016). Gamification of cognitive assessment and cognitive training: A systematic review of applications and efficacy. *JMIR Serious Games*, 4(2), e11.
- Lundqvist, T. (2005). Cognitive consequences of cannabis use: Comparison with abuse of stimulants and heroin with regard to attention, memory and executive functions. *Pharmacology Biochemistry and Behavior*, 81(2), 319–330. <https://doi.org/10.1016/j.pbb.2005.02.017>
- Mechlans, D. J., Irvine, M., Banca, P., Porter, L., Mitchell, S., Mole, T. B., Lapa, T. R., Harrison, N. A., Potenza, M. N., & Voon, V. (2014). Enhanced attentional bias

- towards sexually explicit cues in individuals with and without compulsive sexual behaviours. *PLoS ONE*, 9(8), e105476.
- Mennig, M., Tennie, S., & Barke, A. (2020). A psychometric approach to assessments of problematic use of online pornography and social networking sites based on the conceptualizations of internet gaming disorder. *BMC Psychiatry*, 20(1), 318. <https://doi.org/10.1186/s12888-020-02702-0>
- Meule, A., & Gearhardt, A. N. (2019). Ten years of the Yale Food Addiction Scale: A review of version 2.0. *Current Addiction Reports*, 6(3), 218–228. <https://doi.org/10.1007/s40429-019-00261-3>
- Meule, A., Lutz, A., Vögele, C., & Kübler, A. (2012). Women with elevated food addiction symptoms show accelerated reactions, but no impaired inhibitory control, in response to pictures of high-calorie food-cues. *Eating Behaviors*, 13(4), 423–428. <https://doi.org/10.1016/j.eatbeh.2012.08.001>
- Moeller, F. G., Barratt, E. S., Dougherty, D. M., Schmitz, J. M., & Swann, A. C. (2001). Psychiatric aspects of impulsivity. *American Journal of Psychiatry*, 158(11), 1783–1793. <https://doi.org/10.1176/appi.ajp.158.11.1783>
- Müller, K. W., Dreier, M., Beutel, M. E., & Wölfling, K. (2016). Is sensation seeking a correlate of excessive behaviors and behavioral addictions? A detailed examination of patients with Gambling Disorder and Internet Addiction. *Psychiatry Research*, 242, 319–325. <https://doi.org/10.1016/j.psychres.2016.06.004>
- Murphy, C. M., Stojek, M. K., & MacKillop, J. (2014). Interrelationships among impulsive personality traits, food addiction, and Body Mass Index. *Appetite*, 73, 45–50. <https://doi.org/10.1016/j.appet.2013.10.008>
- Neal, D. J., & Simons, J. S. (2007). Inference in regression models of heavily skewed alcohol use data: A comparison of ordinary least squares, generalized linear models, and bootstrap resampling. *Psychology of Addictive Behaviors*, 21(4), 441–452. <https://doi.org/10.1037/0893-164X.21.4.441>
- Nikolaïdou, M., Fraser, D. S., & Hinvest, N. (2019). Attentional bias in Internet users with problematic use of social networking sites. *Journal of Behavioral Addictions*, 8(4), 733–742. <https://doi.org/10.1556/2006.8.2019.60>
- O'Brien, H., Callinan, S., Livingston, M., Doyle, J. S., & Dietze, P. M. (2020). Population patterns in Alcohol Use Disorders Identification Test (AUDIT) scores in the Australian population; 2007–2016. *Australian and New Zealand Journal of Public Health*, 44(6), 462–467. <https://doi.org/10.1111/1753-6405.13043>
- Odlaug, B. L., Chamberlain, S. R., Kim, S. W., Schreiber, L. R. N., & Grant, J. E. (2011). A neurocognitive comparison of cognitive flexibility and response inhibition in gamblers with varying degrees of clinical severity. *Psychological Medicine*, 41(10), 2111–2119. <https://doi.org/10.1017/S0033297111000316>
- Qualtrics software, Version 08.23. Copyright © 2023 Qualtrics. Qualtrics and all other Qualtrics product or service names are registered trademarks or trademarks of Qualtrics, Provo, UT, USA. <https://www.qualtrics.com>.
- Raj, K., Segrave, R., Tiego, J., Verdéjo-García, A., & Yücel, M. (2022). Problematic Use of the Internet among Australian university students: Prevalence and profile. *Computers in Human Behavior Reports*, 8, Article 100243. <https://doi.org/10.1016/j.chbr.2022.100243>
- Ridderinkhof, K. R., Van Den Wildenberg, W. P. M., Segalowitz, S. J., & Carter, C. S. (2004). Neurocognitive mechanisms of cognitive control: The role of prefrontal cortex in action selection, response inhibition, performance monitoring, and reward-based learning. *Brain and Cognition*, 56(2), 129–140. <https://doi.org/10.1016/j.bandc.2004.09.016>
- Rodrigue, C., Ouellette, A.-S., Lemieux, S., Tchernof, A., Biertho, L., & Bégin, C. (2018). Executive functioning and psychological symptoms in food addiction: A study among individuals with severe obesity. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity*, 23(4), 469–478. <https://doi.org/10.1007/s40519-018-0530-1>
- Rodriguez, L. M., Litt, D. M., & Stewart, S. H. (2020). Drinking to cope with the pandemic: The unique associations of COVID-19-related perceived threat and psychological distress to drinking behaviors in American men and women. *Addictive Behaviors*, 110, Article 106532. <https://doi.org/10.1016/j.addbeh.2020.106532>
- Saunders, J. B., Aasland, O. G., Babor, T. F., De La Fuente, J. R., & Grant, M. (1993). Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption-II. *Addiction*, 88(6), 791–804. <https://doi.org/10.1111/j.1360-0443.1993.tb02093.x>
- Schulte, E. M., & Gearhardt, A. N. (2017). Development of the Modified Yale Food Addiction Scale Version 2.0: Development of the mYFAS 2.0. *European Eating Disorders Review*, 25(4), 302–308. <https://doi.org/10.1002/erv.2515>
- Sepas, A., El-Hussuna, A., Atici, S. D., & Yang, W. (2021). The association between problematic Instagram use, psychological distress, and wellbeing: A systematic review and meta-analysis. *SSRN Electronic Journal*. <https://doi.org/10.2139/ssrn.3956428>
- Sjoberg, E. A., & Cole, G. G. (2018). Sex differences on the Go/No-Go Test of inhibition. *Archives of Sexual Behavior*, 47(2), 537–542. <https://doi.org/10.1007/s10508-017-1010-9>
- Smith, J. L., Mattick, R. P., Jamadar, S. D., & Iredale, J. M. (2014). Deficits in behavioural inhibition in substance abuse and addiction: A meta-analysis. *Drug and Alcohol Dependence*, 145, 1–33. <https://doi.org/10.1016/j.drugalcdep.2014.08.009>
- Starcke, K., Wiesen, C., Trotzke, P., & Brand, M. (2016). Effects of acute laboratory stress on executive functions. *Frontiers in Psychology*, 7. <https://doi.org/10.3389/fpsyg.2016.00461>
- Stavro, K., Pelletier, J., & Potvin, S. (2013). Widespread and sustained cognitive deficits in alcoholism: A meta-analysis: Alcoholism and cognition. *Addiction Biology*, 18(2), 203–213. <https://doi.org/10.1111/j.1369-1600.2011.00418.x>
- Szabó, M. (2010). The short version of the Depression Anxiety Stress Scales (DASS-21): Factor structure in a young adolescent sample. *Journal of Adolescence*, 33(1), 1–8. <https://doi.org/10.1016/j.jad.2009.05.014>
- Tiego, J., Lochner, C., Ioannidis, K., Brand, M., Stein, D. J., Yücel, M., Grant, J. E., & Chamberlain, S. R. (2021). Measurement of the problematic usage of the Internet unidimensional quasitrait continuum with item response theory. *Psychological Assessment*, 33(7), 652–671. <https://doi.org/10.1037/pas0000870>
- Tiego, J., Oostermeijer, S., Prochazkova, L., Parkes, L., Dawson, A., Youssef, G., Oldenhof, E., Carter, A., Segrave, R. A., Fontenelle, L. F., & Yücel, M. (2019). Overlapping dimensional phenotypes of impulsivity and compulsivity explain co-occurrence of addictive and related behaviors. *CNS Spectrums*, 24(4), 426–440. <https://doi.org/10.1017/S1092852918001244>
- Tiego, J., Trender, W., Hellyer, P. J., Grant, J. E., Hampshire, A., & Chamberlain, S. R. (2023). Measuring compulsivity as a self-reported multidimensional transdiagnostic construct: Large-scale (N = 182,000) validation of the Cambridge-Chicago Compulsivity Trait Scale. *Assessment*, 10731911221149083. <https://doi.org/10.1177/10731911221149083>
- VanderBroek-Stice, L., Stojek, M. K., Beach, S. R. H., vanDellen, M. R., & MacKillop, J. (2017). Multidimensional assessment of impulsivity in relation to obesity and food addiction. *Appetite*, 112, 59–68. <https://doi.org/10.1016/j.appet.2017.01.009>
- Verbruggen, F., Aron, A. R., Band, G. P., Beste, C., Bissett, P. G., Brockett, A. T., ... Boehler, C. N. (2019). A consensus guide to capturing the ability to inhibit actions and impulsive behaviors in the stop-signal task. *eLife*, 8, e46323.
- Volkow, N. D., Michaelides, M., & Baler, R. (2019). The neuroscience of drug reward and addiction. *Physiological Reviews*, 99(4), 2115–2140. <https://doi.org/10.1152/physrev.00014.2018>
- Volkow, N. D., & Morales, M. (2015). The brain on drugs: From reward to addiction. *Cell*, 162(4), 712–725. <https://doi.org/10.1016/j.cell.2015.07.046>
- Wei, L., Zhang, S., Turel, O., Bechara, A., & He, Q. (2017). A tripartite neurocognitive model of Internet Gaming Disorder. *Frontiers in Psychiatry*, 8, 285. <https://doi.org/10.3389/fpsyg.2017.00285>
- Yang, X., Guo, W., Tao, Y., Meng, Y., Wang, H., Li, X., ... Li, T. (2022). A bidirectional association between internet addiction and depression: A large-sample longitudinal study among Chinese university students. *Journal of Affective Disorders*, 299, 416–424. <https://doi.org/10.1016/j.jad.2021.12.013>
- Yücel, M., Oldenhof, E., Ahmed, S. H., Belin, D., Billieux, J., Bowden-Jones, H., ... Verdejo-García, A. (2019). A transdiagnostic dimensional approach towards a neuropsychological assessment for addiction: An international Delphi consensus study: Transdiagnostic neuropsychological approaches to addiction. *Addiction*, 114(6), 1095–1109. <https://doi.org/10.1111/add.14424>
- Zhou, Z., Li, C., & Zhu, H. (2013). An Error-Related Negativity Potential Investigation of Response Monitoring Function in Individuals with Internet Addiction Disorder. *Frontiers in Behavioral Neuroscience*, 7. <https://doi.org/10.3389/fnbeh.2013.00131>