

ARTICLE OPEN



Body composition of infants at 6 months of age using a 3-compartment model

Rebecca Kuriyan ¹✉, Andrew P. Hills ², Alexia Murphy-Alford ³, Ramya Padmanabha¹, Lukhanyo H. Nyati ⁴, Nuala M. Byrne², Anura V. Kurpad ¹, Shane Norris ^{4,5} and Multi-Center Infant Body Composition Reference Study (MIBCRS)*

© The Author(s) 2023

BACKGROUND/OBJECTIVES: Two compartment (2 C) models of body composition, including Air Displacement Plethysmography (ADP) and Deuterium Dilution (DD), assume constant composition of fat-free mass (FFM), while 3-compartment (3 C) model overcomes some of these assumptions; studies are limited in infants. The objective of the present study is to compare 3 C estimates of body composition in 6-mo. old infants from Australia, India, and South Africa, including FFM density and hydration, compare with published literature and to evaluate agreement of body composition estimates from ADP and DD.

METHODS: Body volume and water were measured in 176 healthy infants using ADP and DD. 3C-model estimates of fat mass (FM), FFM and its composition were calculated, compared between countries (age and sex adjusted) and with published literature. Agreement between estimates from ADP and DD were compared by Bland–Altman and correlation analyses.

RESULTS: South African infants had significantly higher % FM (11.5%) and density of FFM compared to Australian infants. Australian infants had significantly higher % FFM ($74.7 \pm 4.4\%$) compared to South African infants (71.4 ± 5.0) and higher FFMI ($12.7 \pm 0.8 \text{ kg/m}^2$) compared to South African ($12.3 \pm 1.2 \text{ kg/m}^2$) and Indian infants ($11.9 \pm 1.0 \text{ kg/m}^2$). FFM composition of present study differed significantly from literature. Pooled three country estimates of FM and FFM were comparable between ADP and DD; mean difference of -0.05 (95% CI: $-0.64, +0.55$) kg and $+0.05$ (95% CI: $-0.55, +0.64$) kg.

CONCLUSIONS: 3C-model estimates of body composition in infants differed between countries; future studies are needed to confirm these findings and investigate causes for the differences.

European Journal of Clinical Nutrition (2024) 78:936–942; <https://doi.org/10.1038/s41430-023-01351-2>

INTRODUCTION

Body composition measurements in infants provide a critical baseline for understanding associations between growth in early life and the risk of disease throughout later life [1]. However, the accurate assessment of body composition in infancy is challenging, with errors due to unsuitable prediction equations, movement artefacts during imaging methods, and the unsuitability of many instruments for small infants. An optimal infant body composition measurement method needs to be practical, accurate, precise with minimal assumptions. Sexual dimorphism in relation to adiposity is established in early infancy and may be of relevance for later-life diseases [2, 3].

The 2C-model divides the body into FM and FFM, and 2 C techniques such as densitometry (body volume measurements by ADP) and isotope dilution (hydrometry, DD), have commonly been used to assess body composition in infants and children [4, 5]. While these techniques are practical and relatively simple, they assume values of FFM hydration or density, primarily based on studies conducted on infants from Western countries by Fomon et al. [6] and Butte et al. [7]; these assumed age- and sex-specific FFM density or hydration may not be appropriate universally, since many external factors can impact the process of growth and

maturation [8, 9]. These measurements are limited in infants from diverse countries and ethnicities.

Multi-component models of body composition, combining measurements from different techniques, can be used to directly assess the FFM composition (water and mineral content) and thus explore the variations in FFM hydration and density. In the criterion four compartment (4 C) model [10, 11], total body water (TBW), bone mineral content (BMC) and body density (from body volume and weight), are assessed using hydrometry, dual-energy X-ray absorptiometry (DXA) and ADP, respectively. However, since DXA imaging involves exposure to a low dose of radiation and is affected by movement artefacts, it may not be appropriate for new-borns and young infants [2, 12]. On the other hand, the non-imaging 3-compartment (3 C) model [13], which combines measures of weight, body volume by ADP, and total body water by DD, avoids the assumption of water content of FFM between individuals and has been used safely in infants to yield estimates of FM, TBW and the anhydrous FFM (mineral, protein and DNA/glycogen) [14]. The 3 C model can be used in research settings to explore the assumed constants of density and hydration of FFM [6, 7] and provide important understanding and implications while applying 2 C models. Evaluation of the FFM density of Fomon et al. [6] against 3C-model estimates in a small number of Ethiopian

¹St. John's Research Institute, Bengaluru, India. ²University of Tasmania, Hobart, TAS, Australia. ³International Atomic Energy Agency, Vienna, Austria. ⁴University of Witwatersrand, Johannesburg, South Africa. ⁵University of Southampton, Southampton, UK. *A list of authors and their affiliations appears at the end of the paper. ✉email: rebecca@sjri.res.in

Received: 9 April 2023 Revised: 16 September 2023 Accepted: 21 September 2023

Published online: 13 October 2023

infants aged 1.5–6 mo. [13], found them to be similar. However, to our knowledge, no study has simultaneously estimated body composition using the 3C-model in infants from countries with varied income and environmental settings, which could affect the FFM hydration or density.

This study firstly aimed to measure and compare the anthropometry and body composition of 6-mo. old infants from Australia, India and South Africa, using the 3C-model. Secondly, the study aimed to investigate the FFM density and hydration of infants and compare the values between countries and to published values [6, 7] Thirdly, the study aimed to evaluate the agreement of FM and FFM estimates between ADP and DD.

METHODS

This study included 6-mo. old infants from three countries, who were part of the larger multi-country Multi-center Infant Body Composition Reference Study (MIBCRS), which aimed to accurately assess body composition in healthy infants from birth to 2 y. The eligibility criteria for recruitment into MIBCRS were consistent with the WHO Multicentre Growth Reference Study (MGRS) [15], and full-term healthy infants (≥ 37 to ≤ 42 wk of gestation), weighing ≥ 2500 g, from singleton pregnancies of healthy non-smoker mothers planning to breastfeed for at least up to 6 mo., were recruited into the study. Infants with significant morbidity and congenital abnormalities were excluded. This multinational study fulfilled International Ethical Guidelines for Biomedical Research involving human participants, and each country site obtained approval from their local ethics review committee. Informed consent form was obtained from the mothers. The sample size of 44 infants in each country was estimated to observe a mean difference of 3.0% in %FM between different countries, with a SD of 3.9% and 4.2% [16], at 5% level of significance and 80% power, after correcting for the multiple comparison between 3 countries.

Infant anthropometry

Infant weight on the day of measurement (at ~6 mo.) was measured using a pediatric electronic scale accurate to the nearest 0.01 kg (SECA 376, Hamburg, Germany, for Australian and South African infants; Salter 914, Delhi, India, for Indian infants). Length was measured to the nearest 0.1 cm using an infantometer (SECA 417, Hamburg, Germany). Standardized protocols were developed based on the WHO MGRS protocol [17], measurements were performed by staff who were initially trained at all sites by an international expert and then retrained every 3 months.

Assessment of body density by ADP

Body volume, and thereby body density, was measured by ADP (PEA POD, Software version 3.5.0, 201, COSMED, USA), with a mean precision of 0.07%. Precision was measured at each study site using a standard 3 L volume phantom, provided by the manufacturer. The volume of the phantom was measured 10 times for three days, and the coefficient of variation (CV%) was calculated as standard deviation (SD)/ Mean. The mean CV% of the three countries was reported. Standard protocol was used at all sites [18]. The ratio of weight (kg) and the measured body volume (L) was used to calculate total body density (kg/L), from which the proportions of FM and FFM were calculated using assumed densities [6, 19] for each. FM and FFM were expressed as a percentage of body weight (% FM, % FFM), and in relation to height, as FMI (kg/m^2) and FFMI (kg/m^2).

Assessment of TBW by DD

Infant TBW was determined by DD, where a dose of 1 g deuterium oxide (D_2O ; 99.8 atom %, Sigma-Aldrich, Canada) was orally administered to the infant using a syringe, following a baseline saliva sample, collected at least 15 min prior to their last feed. Saliva was sampled at 2.5 h and 3 h, allowing sufficient time for the isotope to equilibrate in the body water. The 3-h point was taken for the calculation [20]. Liquids were not provided to the infants between dosing and saliva sample collections. The collected saliva samples were stored in -20°C freezer until analysis. The deuterium

Table 1. Maternal and infant characteristics of the study population.

Variables	Australia (n = 46)	India (n = 86)	South Africa (n = 44)	Pooled (n = 176)	P-value
Maternal characteristics					
Maternal age (y)	30.3 \pm 4.3 ^{a,b}	25.8 \pm 6.4 ^a	25.1 \pm 5.4 ^b	26.8 \pm 6.0*	<0.01
Gestational age (wk)	39.8 \pm 1.0 ^{a,b}	38.8 \pm 0.9 ^a	39.1 \pm 1.4 ^b	39.2 \pm 1.2*	<0.01
Mode of delivery (%)					
Vaginal (spontaneous and assisted)	73.9	72.1	100.0	79.6	0.02
LSCS ^d	26.1	27.9	0.0	20.4	
Parity %					
Primiparous	35.2	57.0	56.1	53.4	0.17
Multiparous	64.8	43.0	43.9	46.6	
Infant characteristics					
Age (year)	0.49 \pm 0.02 ^{a,b}	0.51 \pm 0.02 ^{a,c}	0.50 \pm 0.01 ^{b,c}	0.50 \pm 0.02*	<0.01
Weight at birth visit (kg) ^e	3.3 \pm 0.4 ^{a,b}	2.8 \pm 0.3 ^{a,c}	3.0 \pm 0.3 ^{b,c}	3.0 \pm 0.4*	<0.01
Weight (kg)	7.3 \pm 0.7	7.2 \pm 0.8	7.2 \pm 0.7	7.2 \pm 0.8	0.75
Length (cm)	65.3 \pm 2.2	66.0 \pm 2.2	64.8 \pm 2.9	65.5 \pm 2.5	0.03
Sex %					
Female	41.3	51.2	65.9	52.3	0.06
Male	58.7	48.8	34.1	47.7	
Feeding practices at 3 mo. of infants age (%)					
Exclusively breastfed	60.9	94.9	21.4	66.9	<0.01
Exclusively formula fed	13.0	0.0	2.4	4.2	

Values are mean \pm SD; P values using ANOVA, Pearson's chi-square, or ANCOVA test as appropriate.

* $P < 0.05$;

^aWithin a row, superscripts denote significance between countries—Australia vs. India

^bWithin a row, superscripts denote significance between countries—Australia vs. South Africa

^cWithin a row, superscripts denote significance between countries—India vs. South Africa.

^dLower Segment Cesarean Section;

^e0–8 d of postnatal age.

abundance in the thawed saliva samples was analyzed in duplicate by Fourier transformed infrared spectrometry (FTIR, Agilent 4500 Series, USA) in separate laboratories. Similar SOPs were followed in all countries and efforts were made to harmonize the equipment and protocol. Quality control of TBW analysis was assured by the regular site visits by IAEA technical expert and the interlab studies, which were comparable (Supplementary Table 1). TBW was calculated as the ratio of the administered dose of deuterium to its steady state enrichment in the body water, corrected for deuterium exchange into protein using a correction factor of 1.044 [21]. FFM was calculated from TBW assuming an age- and sex-specific hydration value of FFM [6, 22]. FM was calculated as the difference in body weight and FFM.

Estimation of FM, FFM, density, and hydration of FFM, using the 3C model

The 3C-model estimates of FM and FFM were derived using body weight (BW), body volume (BV, measured by ADP) and TBW (estimated from DD), assuming the density of fat (D_{FM}) to be 0.9007 kg/L [19], and the density of water (D_W) to be 0.9937 kg/L at body temperature [19]. The density (D_{DB}) and composition of the 'dry' body (FM and the anhydrous FFM) were calculated using the Archimedes principle for binary mixtures with the proportions and assumed densities of fat (P_F and D_F) and the density of anhydrous FFM (D_{p+m}), rearranging to calculate the unknown fat proportion P_F as shown below.

$$\frac{1}{D_{DB}} = \frac{P_F}{D_F} + \frac{1 - P_F}{D_{p+m}}$$

The density of the protein and mineral (D_{p+m}) component was derived as 1.4959 kg/L, using sex-specific proportions and densities of protein (P_p , D_p) and mineral (P_m , D_m) in a 6-mo infant [19].

$$\frac{1}{D_{p+m}} = \frac{P_p}{D_p} + \frac{P_m}{D_m}$$

FFM density (DFFM) was calculated from the ratio of FFM and FFM volume as below:

$$D_{FFM}(\text{kg/L}) = \frac{\text{Body weight (kg)} - \text{FM (kg)}}{\text{Body volume (L)} - (\text{FM/FFM density}) (\text{L})}$$

FFM hydration (HF) was calculated as ratio of measured TBW, and 3C-model estimate of FFM.

Statistical analysis

Primary measurements were tested for normality assumptions using Q-Q plot and Shapiro Wilk test ($p > 0.05$). Continuous variables are presented as mean \pm SD, while categorical variables are presented as percentage (%). Maternal and infant characteristics were compared between countries using Pearson's Chi-square test and analysis of variance (ANOVA), as appropriate and multiple comparison was performed by 2×2 Chi-square (adjusted for p value) and Bonferroni correction method, respectively. The main effect of country and interaction effect of country \times sex were analysed using Analysis of covariance (ANCOVA) to compare the infant anthropometry and body composition estimates from 3C model, between the three countries adjusted for age and sex, followed by Bonferroni correction for multiple comparisons. The agreement of FM and FFM estimates obtained from ADP vs. DD were assessed using intraclass correlation (ICC) and the Bland-Altman method [23]. The one sample t -test was used to determine if the mean difference of FM estimates from ADP and DD were significantly different from zero. The estimated D_{FFM} and HF were compared with published data [6, 7] using the one sample t -test. The mean precision of the estimated FM (kg) using the 3C model, between the three countries, obtained from the propagation of error method was ± 0.03 kg (1.3% of FM), calculated using the known error in measuring TBW from DD (0.02 kg), body weight (0.01 kg) and body volume (0.002 L). SPSS statistical package version 26.0 (SPSS Inc. Chicago, Illinois) was used. $P < 0.05$ was considered statistically significant.

RESULTS

The data were normally distributed; maternal and infant characteristics are presented in Table 1. Australian mothers were significantly older than mothers from India and South Africa. Majority (95%) of the Indian infants were exclusively breastfed (EBF) at 3 mo., while 61% and 21% of infants were EBF in Australia and South Africa, respectively. About 91% of Australian infants were Caucasian, while all infants from India were of Asian ethnicity and South African infants belonged to Black African ethnicity. There were no between-country body weight and length differences in infants at 6 mo. of age.

3C-model estimates of body composition

Body composition estimates obtained from the 3C-model for infants are presented in Table 2. Australian infants had significantly higher

Table 2. Body composition estimates using the 3C-model in infants from the three countries at 6 mo. of age.

	Australia (n = 46)	India (n = 86)	South Africa (n = 44)	Pooled (n = 176)	P-value
Primary parameters					
Weight (kg)	7.3 \pm 0.7	7.2 \pm 0.8	7.2 \pm 0.7	7.2 \pm 0.8	0.50
TBW (kg)	4.3 \pm 0.4 ^{a,b}	4.1 \pm 0.5 ^a	4.0 \pm 0.4 ^b	4.2 \pm 0.4*	0.01
TBW %	59.4 \pm 4.1 ^b	57.7 \pm 4.5	56.1 \pm 4.6 ^b	57.7 \pm 4.5*	0.01
Body volume (kg/L)	7.2 \pm 0.7	7.1 \pm 0.8	7.1 \pm 0.7	7.1 \pm 0.8	0.63
Body composition estimates					
FM (kg)	1.9 \pm 0.4	2.0 \pm 0.5	2.1 \pm 0.5	2.0 \pm 0.5	0.13
FFM (kg)	5.4 \pm 0.5 ^{a,b}	5.2 \pm 0.5 ^a	5.1 \pm 0.4 ^b	5.2 \pm 0.5*	0.01
% FM	25.3 \pm 4.4 ^b	27.4 \pm 4.9	28.6 \pm 5.0 ^b	27.1 \pm 4.9*	0.02
% FFM	74.7 \pm 4.4 ^b	72.6 \pm 4.9	71.4 \pm 5.0 ^b	72.9 \pm 4.9*	0.02
FMI (kg/m ²)	4.3 \pm 1.0	4.5 \pm 1.1	5.0 \pm 1.4	4.6 \pm 1.2	0.07
FFMI (kg/m ²)	12.7 \pm 0.8 ^{a,b}	11.9 \pm 1.0 ^a	12.3 \pm 1.2 ^b	12.2 \pm 1.1*	<0.01
D_{FFM} (kg/L)	1.067 \pm 0.006 ^b	1.067 \pm 0.006	1.071 \pm 0.008 ^b	1.068 \pm 0.007*	0.03
HF	0.795 \pm 0.015	0.795 \pm 0.017	0.785 \pm 0.020	0.792 \pm 0.018	0.03

D_{FFM} Density of fat-free mass, HF Hydration factor, FMI Fat Mass Index, FFMI Fat-free mass index, TBW Total body water;

Values are mean \pm SD; P values for main effects of country \times sex using ANCOVA.

* $P < 0.05$,

^aWithin a row, superscripts denote significance between countries—Australia vs. India;

^bWithin a row, superscripts denote significance between countries—Australia vs. South Africa;

^cWithin a row, superscripts denote significance between countries—India vs. South Africa

Table 3. Sex-specific comparison of the density and hydration of fat-free mass (FFM) from the present study with published literature values [6, 7].

	Australia		India		South Africa		Pooled	
	Males n = 27	Females n = 19	Males n = 42	Females n = 44	Males n = 15	Females n = 29	Males n = 84	Females n = 92
Density of FFM (D_{FFM}) kg/L	Present data 1.067 ± 0.006	1.067 ± 0.005	1.067 ± 0.006	1.068 ± 0.007	1.069 ± 0.008	1.072 ± 0.007	1.067 ± 0.007	1.069 ± 0.007
	Fomon et al. [6] 1.066	1.067	1.066	1.067	1.066	1.067	1.066	1.067
	<i>P</i> value 0.456	0.799	0.337	0.543	0.165	0.001*	0.073	0.009*
Hydration of FFM (H_{FFM})	Butte et al. [7] 1.062 ± 0.004	1.062 ± 0.006	1.062 ± 0.004	1.062 ± 0.006	1.062 ± 0.004	1.062 ± 0.006	1.062 ± 0.004	1.062 ± 0.006
	<i>P</i> value 0.001*	<0.001*	<0.001*	<0.001*	0.004*	<0.001*	<0.001*	<0.001*
	Present data 0.796 ± 0.017	0.795 ± 0.014	0.796 ± 0.016	0.794 ± 0.017	0.790 ± 0.021	0.783 ± 0.019	0.795 ± 0.017	0.790 ± 0.018
	Fomon et al. [6] 0.796	0.794	0.796	0.794	0.796	0.794	0.796	0.794
	<i>P</i> value 0.891	0.861	0.867	0.948	0.298	0.004*	0.460	0.058
	Butte et al. [7] 0.807 ± 0.012	0.807 ± 0.016	0.807 ± 0.012	0.807 ± 0.016	0.807 ± 0.012	0.807 ± 0.016	0.807 ± 0.012	0.807 ± 0.016
	<i>P</i> value 0.002*	0.001*	<0.001*	<0.001*	0.007*	<0.001*	<0.001*	<0.001*

Values are mean ± SD; One sample *t*-test;
*depicts statistical significance $P < 0.05$.

TBW (kg) and FFM (kg) when compared to Indian and South African infants, but %TBW adjusted for body weight ($59.4 \pm 4.1\%$), was only significantly higher than infants from South Africa ($56.1 \pm 4.6\%$). The %FFM of Australian infants ($74.7 \pm 4.4\%$) was significantly higher than infants from South African ($71.4 \pm 5.0\%$), while the %FM was significantly lower in Australian infants ($25.3 \pm 4.4\%$), when compared to South African ($28.6 \pm 5.0\%$). The FM (kg) was similar among the three countries. Infants from Australia had a significantly higher FFMI ($12.7 \pm 0.8 \text{ kg/m}^2$) compared to infants from India ($11.9 \pm 1.0 \text{ kg/m}^2$) and South Africa ($12.3 \pm 1.2 \text{ kg/m}^2$). Pooled sex-specific estimates of the body composition showed that males had significantly higher TBW (kg), %TBW, FFM (kg), %FFM and FFMI, while females had significantly higher %FM. These results along with the within country sex-specific results are summarized in Supplementary Table 2.

Density and hydration of FFM

The D_{FFM} of South African infants ($1.071 \pm 0.008 \text{ kg/L}$) was significantly higher than Australian infants ($1.067 \pm 0.006 \text{ kg/L}$, P value = 0.047), but not Indian infants ($1.067 \pm 0.006 \text{ kg/L}$, $P = 0.087$). The HF of South African infants (0.785 ± 0.020) was lower compared to Australian infants (0.795 ± 0.020), however, not significant. ($P = 0.050$). These results are reported in Table 2. Sex-specific analysis between countries (Supplementary Table 1) showed that South African females were significantly different from Indian females, but not Australian females in both D_{FFM} and HF.

When comparing the sex-specific D_{FFM} and HF estimates from the present study with those of Fomon et al. [6], only females from South Africa had significantly higher D_{FFM} and lower HF, while comparison of D_{FFM} and HF data from the present study with estimates from Butte et al. [7] showed that they were significantly different in infants from all three countries (Table 3).

Comparison of ADP and DD


The ICC of pooled FM and FFM estimates obtained from ADP and DD was high (0.90, 95% CI: 0.86–0.92 and 0.91, 95% CI: 0.88–0.94, respectively). The mean difference in the pooled FM and FFM estimates obtained from DD and ADP was -0.05 ± 0.30 (95% CI: $-0.64, +0.55$) kg and $+0.05 \pm 0.30$ (95% CI: $-0.55, +0.64$) kg respectively, which were significantly different from zero ($P < 0.05$) (Fig. 1). The mean difference between the two methods for Australia, India, and South Africa were 0.00 ± 0.27 (95% CI: $-0.55, +0.54$) kg, -0.01 ± 0.28 (95% CI: $-0.57, +0.55$) kg, -0.17 ± 0.33 (95% CI: $-0.82, +0.49$) kg, respectively for FM and 0.00 ± 0.27 (95% CI: $-0.54, +0.55$) kg, $+0.01 \pm 0.28$ (95% CI: $-0.55, +0.57$) kg, 0.17 ± 0.33 (95% CI: $-0.49, +0.82$) kg, respectively for FFM. The correlation between the difference and average of FM derived from ADP and DD was inverse and significant ($r = -0.18, p = 0.02$; $y = -0.1108x + 0.171$), while it was linear and significant for FFM ($r = +0.18, p = 0.02$; $y = -0.1059x + 0.6034$).

DISCUSSION

This paper presents, for the first time, 3C-model body composition estimates of 6 mo. infants recruited from three different countries in a simultaneous multicenter study. The rank order of FFMI was Australia > South Africa > India, while for FMI, it was South Africa > India > Australia. Infants from South Africa had significantly higher density of FFM when compared to infants from Australia. Sexual dimorphism existed in South African infants, females had a significantly higher D_{FFM} and lower HF when compared to published estimates [6]. Pooled FM (kg) and FFM (kg) estimates from the three countries, measured by ADP and DD, had good reliability and relatively low differences.

Measurement of body composition at 6 mo. can provide insights on the effect of feeding patterns on body composition. A systematic review summarized the effect of infant feeding (breast feeding vs formula feeding) on infant (0–12 mo.) body composition and observed that formula-fed infants had lower FM when

A



B




Fig. 1 Bland–Altman plot comparing the body composition estimates of Air displacement plethysmography (ADP) vs. Deuterium dilution (DD) in infants from the three countries at 6 mo. of age. A Fat mass obtained from ADP vs. DD. **B** Fat-free mass obtained from ADP vs. DD. Circles denote Australia ($n = 46$), squares denote India ($n = 86$) and triangles denote South Africa ($n = 44$). The black solid line refers to zero. The black dashed line represents the mean difference (-0.05 kg) for FM and ($+0.05$ kg) for FFM, which was significantly different ($P < 0.05$), from zero. The limits of agreement are represented by dotted lines representing upper and lower limits of 95% CI: -0.64 , $+0.55$ kg for FM and -0.55 , $+0.64$ kg for FFM.

compared to breast fed infants at 3–4 mo. and 6 mo., but there was a trend toward higher FM in formula-fed infants at 12 mo. of age [24]. Similar results were observed in a large, mother-baby dyad group in USA, where breastfed babies had higher percent body fat at 6 mo. of age [25]. Breast fed infants were observed to have higher circulating leptin in comparison to formula-fed infants at ages < 4 mo., but not in later infancy [26, 27], which could be a reason for the observed higher FM in breast fed infants during early infancy. Similarly, breastfeeding was associated with lower FFM in infants between 3–6 mo. [24, 28] and at 6 mo., when compared to formula feeding [24]. The higher protein content of formula feeds compared to breast milk [25] may explain the higher accrual of FFM in early life, however, these findings need to be prospectively confirmed in longitudinal studies.

In the present study, sex-based differences in D_{FFM} and HF were only observed in South African infants, however, since males only comprised 34% ($n = 15$), results need to be considered with caution. Earlier studies [7, 29, 30] have not observed any

significant sex-specific differences and it has been suggested that attempts to identify sex-based differences in the composition of FFM in infants may not be desirable unless the methodology was perfect [31].

Most body composition reference data for infants have been published using [13, 18, 32–34] the 2C model, which assumes constant values for the composition of FFM [6, 7]. The values from Fomon et al. [6], were based on data from different published reports, modeled with many assumptions to derive reference data for D_{FFM} and HF. Butte et al. values [7] were based on longitudinal data from infants of US aged birth to 2 y. The D_{FFM} of infants from all three countries in the present study were significantly higher, while HF estimates were significantly lower than those from Butte et al. [7]. Similarly, lower body FM estimates at 12 wk of age [35], was observed when using the D_{FFM} value from Butte et al. [7] in comparison to Fomon et al. [6], being particularly evident at 1 wk of age, which could have implications for estimates of FM and FM accretion in early life. However, the biological importance of these

findings may be more relevant rather than just the statistical significance and hence population specific D_{FFM} and HF, which may be best applicable for each country should be used.

Estimates of body composition measured by ADP and DD in the present study showed good reliability, with a low mean difference, concurring to results observed in infants aged 0.5 to 6 mo., where no significant differences were observed between percent FM estimates obtained from both methods [36]. While DD is accurate for estimating body composition in infancy, it is not practical in infants under 3 mo. due to difficulties in dosing or collecting saliva samples. ADP is a safe and rapid measurement, validated using the 4C-model in term infants [37, 38], but being practical only for infants weighing up to 8 kg. The results from the two methods should, however, not be used interchangeably while measuring body composition in infants, and may be reliable only for population estimations rather than individual measurements.

Limitations of the present study were the single DD measurement and the lack of feeding data at 6 mo. Factors such as socio-economic status, feeding patterns, ethnicity, and climate, which may have contributed, were not considered in the analysis. In conclusion, accurate estimates of body composition in infants from three countries, using 3C model suggests some significant between-country variability, however, in view of the significant demographic, social, and environmental differences among the countries, larger studies are needed to investigate the true causes of the differences. Estimates using ADP and DD correlated well with relatively low bias, however, the wide 95% CI suggests that these techniques may be more reliable for population estimates. The D_{FFM} of South African infants differed from Australian infants and from estimates from Fomon et al. [6], while all countries differed from estimates of Butte et al. [7]. This highlights the need for further investigation of the assumed FFM variables in infancy, which may have important implications for the different 2C models used in infant research.

DATA AVAILABILITY

Data described in the manuscript, code book, and analytic code will be made publicly and freely available without restriction at, <https://osf.io/r5pbn>, OSF | Body composition of infants at 6 months of age using a 3-compartment model.

REFERENCES

- Wells JC. Body composition in infants: evidence for developmental programming and techniques for measurement. *Rev Endocr Metab Disord.* 2012;13:93–101.
- Davis SM, Kaar JL, Ringham BM, Hockett CW, Glueck DH, Dabelea D. Sex differences in infant body composition emerge in the first 5 months of life. *J Pediatr Endocrinol Metab.* 2019;32:1235–9.
- Gale C, Logan KM, Jeffries S, Parkinson JR, Santhakumaran S, Uthaya S, et al. Sexual dimorphism in relation to adipose tissue and intrahepatocellular lipid deposition in early infancy. *Int J Obes (Lond).* 2015;39:629–32.
- Demerath EW, Fields DA. Body composition assessment in the infant. *Am J Hum Biol.* 2014;26:291–304.
- Gallagher D, Andres A, Fields DA, Evans WJ, Kuczmarski R, Lowe WL Jr, et al. Body composition measurements from birth through 5 years: challenges, gaps, and existing & emerging technologies—a National Institutes of Health Workshop. *Obes Rev.* 2020;21:e13033.
- Fomon SJ, Haschke F, Ziegler EE, Nelson SE. Body composition of reference children from birth to age 10 years. *Am J Clin Nutr.* 1982;35:1169–75.
- Butte NF, Hopkinson JM, Wong WW, Smith EO, Ellis KJ. Body composition during the first 2 years of life: an updated reference. *Pediatr Res.* 2000;47:578–85.
- Wells JCK. Body composition of children with moderate and severe under-nutrition and after treatment: a narrative review. *BMC Med.* 2019;17:215.
- Wang Z, Deurenberg P, Wang W, Pietrobelli A, Baumgartner RN, Heymsfield SB. Hydration of fat-free body mass: review and critique of a classic body-composition constant. *Am J Clin Nutr.* 1999;69:833–41.
- Wells JC, Fuller NJ, Dewit O, Fewtrell MS, Elia M, Cole TJ. Four-component model of body composition in children: density and hydration of fat-free mass and comparison with simpler models. *Am J Clin Nutr.* 1999;69:904–12.
- Heyward V. ASEP methods recommendation: body composition assessment. *J Exerc Physiol Online.* 2001;4:1–12.
- Toro-Ramos T, Paley C, Pi-Sunyer FX, Gallagher D. Body composition during fetal development and infancy through the age of 5 years. *Eur J Clin Nutr.* 2015;69:1279–89.
- Andersen GS, Girma T, Wells JC, Kästel P, Leventi M, Hother AL, et al. Body composition from birth to 6 mo of age in Ethiopian infants: reference data obtained by air-displacement plethysmography. *Am J Clin Nutr.* 2013;98:885–94.
- Siri WE. Body composition from fluid spaces and density: analysis of methods. 1961. *Nutrition.* 1993;9:480–91.
- de Onis M, Garza C, Victora CG, Onyango AW, Frongillo EA, Martines J. The WHO Multicentre Growth Reference Study: planning, study design, and methodology. *Food Nutr Bull.* 2004;25:515–26.
- Paley C, Hull H, Ji Y, Toro-Ramos T, Thornton J, Bauer J, et al. Body fat differences by self-reported race/ethnicity in healthy term newborns. *Pediatr Obes.* 2016;11:361–8.
- de Onis M, Onyango AW, Van den Broeck J, Chumlea WC, Martorell R. Measurement and standardization protocols for anthropometry used in the construction of a new international growth reference. *Food Nutr Bull.* 2004;25:527–36.
- Urlando A, Dempster P, Aitkens S. A new air displacement plethysmograph for the measurement of body composition in infants. *Pediatr Res.* 2003;53:486–92.
- Brozek J. Densitometric analysis of body composition: revision of some quantitative assumptions. *Ann NY Acad Sci.* 1963;110:113–40.
- Murphy-Alford AJ, Johnson W, Nyati LH, Santos IS, Hills AP, Ariff S, et al. Body composition reference charts for infants from birth to 24 months: multi-center infant body composition reference study. *Am J Clin Nutr.* 2023;117:1262–9.
- Racette SB, Schoeller DA, Luke AH, Shay K, Hnilicka J, Kushner RF. Relative dilution spaces of 2H- and 18O-labeled water in humans. *Am J Physiol.* 1994;267:E585–90.
- Schoeller DA. Hydrometry. In: Heymsfield S, Lohman T, Wang Z-MGS editors. *Human body composition 918, human kinetics: champaign.* 2nd edn. IL, USA; Am J Clin Nutr. 2005. p. 536.
- Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet.* 1986;1:307–10.
- Gale C, Logan KM, Santhakumaran S, Parkinson JR, Hyde MJ, Modi N. Effect of breastfeeding compared with formula feeding on infant body composition: a systematic review and meta-analysis. *Am J Clin Nutr.* 2012;95:656–69.
- Tahir MJ, Ejima K, Li P, Demerath EW, Allison DB, Fields DA. Associations of breastfeeding or formula feeding with infant anthropometry and body composition at 6 months. *Matern Child Nutr.* 2021;17:e13105.
- Heinig MJ, Nommsen LA, Peerson JM, Lonnerdal B, Dewey KG. Energy and protein intakes of breast-fed and formula-fed infants during the first year of life and their association with growth velocity: the DARLING Study. *Am J Clin Nutr.* 1993;58:152–61.
- Savino F, Costamagna M, Prino A, Oggero R, Silvestro L. Leptin levels in breast-fed and formula-fed infants. *Acta Paediatr.* 2002;91:897–902.
- Sauder KA, Kaar JL, Starling AP, Ringham BM, Glueck DH, Dabelea D. Predictors of infant body composition at 5 months of age: the Healthy Start Study. *J Pediatr.* 2017;183:94–99.e1.
- Fuller NJ, Jebb SA, Laskey MA, Coward WA, Elia M. Four-component model for the assessment of body composition in humans: comparison with alternative methods, and evaluation of the density and hydration of fat-free mass. *Clin Sci (Lond).* 1992;82:687–93.
- Wells JC, Fuller NJ, Wright A, Fewtrell MS, Cole TJ. Evaluation of air-displacement plethysmography in children aged 5–7 years using a three-component model of body composition. *Br J Nutr.* 2003;90:699–707.
- Fomon SJ, Nelson SE. Body composition of the male and female reference infants. *Annu Rev Nutr.* 2002;22:1–17.
- Rigo J, Nyamugabo K, Picaud JC, Gerard P, Pieltain C, De Curtis M. Reference values of body composition obtained by dual energy X-ray absorptiometry in preterm and term neonates. *J Pediatr Gastroenterol Nutr.* 1998;27:184–90.
- Fields DA, Gilchrist JM, Catalano PM, Gianni ML, Roggero PM, Mosca F. Longitudinal body composition data in exclusively breast-fed infants: a multicenter study. *Obesity (Silver Spring).* 2011;19:1887–91.
- Wells JCK, Davies PSW, Fewtrell MS, Cole TJ. Body composition reference charts for UK infants and children aged 6 weeks to 5 years based on measurement of total body water by isotope dilution. *Eur J Clin Nutr.* 2020;74:141–8.
- Eriksson B, Löf M, Eriksson O, Hannestad U, Forsum E. Fat-free mass hydration in newborns: assessment and implications for body composition studies. *Acta Paediatr.* 2011;100:680–6.
- Ma G, Yao M, Liu Y, Lin A, Zou H, Urlando A, et al. Validation of a new pediatric air-displacement plethysmograph for assessing body composition in infants. *Am J Clin Nutr.* 2004;79:653–60.
- Ellis KJ, Yao M, Shypailo RJ, Urlando A, Wong WW, Heird WC. Body-composition assessment in infancy: air-displacement plethysmography compared with a reference 4-compartment model. *Am J Clin Nutr.* 2007;85:90–5.

38. Heard-Lipsmeyer ME, Hull H, Sims CR, Cleves MA, Andres A. Evaluating body composition in infancy and childhood: a comparison between 4C, QMR, DXA, and ADP. *Pediatr Obes.* 2020;15:e12617.

ACKNOWLEDGEMENTS

The authors acknowledge the study team at the different sites and are grateful to all the mothers and their infants who participated in the study. We thank Jonathan CK Wells for his guidance with the 3 C model.

AUTHOR CONTRIBUTIONS

RK, APH, NMB, AJMA, LN, and SN designed the research (project conception, development of overall research plan, and study oversight); RK, RP, and AVK performed the statistical analyses and wrote the drafts of the paper. RK has primary responsibility for the final content. All authors have read and approved the final manuscript. The authors declare no competing financial interests.

FUNDING

This work was supported, in part, by the International Atomic Energy Agency (CRP E43028) and the Bill & Melinda Gates Foundation [OPP1143641]. Under the grant conditions of the Foundation, a Creative Commons Attribution 4.0 Generic License has already been assigned to the Author Accepted Manuscript version that might arise from this submission. Bill & Melinda Gates Foundation did not participate in the design, management, analysis, interpretation, or preparation of the manuscript.

COMPETING INTERESTS

R.K. is an Associate Editor of *European Journal of Clinical Nutrition*. The other authors have no conflicts or competing financial interests to declare.

MULTI-CENTER INFANT BODY COMPOSITION REFERENCE STUDY (MIBCRS)

Shabina Ariff⁶, Andrew P. Hills², Rebecca Kuriyan¹✉, Shane Norris^{4,5}, Ina S. Santos⁷, V. Pujitha Wickramasinghe⁸, Alexia J. Murphy-Alford³, Lukhanyo Nyati⁴, Caroline S. Costa⁷, Nishani Lucas⁸, Tanvir Ahmad⁹, Kiran D. K. Ahuja², Jeffrey M. Beckett², Renata M. Bielemann⁷, Nuala M. Byrne², Laila Charania⁶, Michele P. Christian¹, Priscilla J. Divya¹, Anne Hanley², Manoj P. Herath², Leila C. Ismail^{10,11}, Sisitha Jayasinghe², Anura V. Kurpad¹, Pulani Lanerolle⁸, Cornelia Loechl³, Najat Moktar³, Upul Senerath⁸, Christine Slater³, Sajid Soofi⁶, Steven J. Street², Neiva C. J. Valle⁷ and Ayesha Yameen⁹

⁶The Aga Khan University, Karachi, Pakistan. ⁷Federal University of Pelotas, Pelotas, Brazil. ⁸University of Colombo, Colombo, Sri Lanka. ⁹Isotope Application Division, Pakistan, Institute of Nuclear Science and Technology (PINSTECH), Islamabad, Pakistan. ¹⁰University of Sharjah, Sharjah, United Arab Emirates. ¹¹University of Oxford, Oxford, UK.

ETHICAL APPROVAL

In each country, the study protocol was cleared by the Ethics Research Committee of the institution where the study was conducted. All women signed an informed consent form agreeing to participate in the study.

ADDITIONAL INFORMATION

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41430-023-01351-2>.

Correspondence and requests for materials should be addressed to Rebecca Kuriyan.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023