

Archives of Disease in Childhood

Magnet Ingestion in Children in the United Kingdom: a national prospective observational surveillance study.

Journal:	<i>Archives of Disease in Childhood</i>
Manuscript ID	archdischild-2024-328195.R1
Article Type:	Original research
Date Submitted by the Author:	20-Dec-2024
Complete List of Authors:	Neville, Jonathan; University of Southampton Faculty of Medicine, University Surgery Unit Lyttle, Mark; Bristol Royal Hospital for Children, Emergency Department; University of the West of England, Academic Department of Emergency Care Messahel, Shrouk; Alder Hey Children's Hospital, Emergency Department Parkar, Shabnam; St George's University Hospitals NHS Foundation Trust Mytton, Julie; University of the West of England Hall, Nigel; University of Southampton Faculty of Medicine, University Surgery Unit
Keywords:	Child Health, Paediatrics, Epidemiology, Emergency Service, Hospital

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Magnet Ingestion in Children in the United Kingdom: a national prospective observational surveillance study.

Jonathan J Neville¹, Mark D Lyttle^{2, 3}, Shrouk Messahel⁴, Shabnam Parkar⁵, Julie Mytton⁶, Nigel J Hall^{1*}, Magnet Ingestion in Children (MAGNETIC) Study collaborators, Paediatric Emergency Research in the UK and Ireland (PERUKI) Group

1. University Surgery Unit, University of Southampton Faculty of Medicine, Southampton UK
2. Paediatric Emergency Department, Bristol Royal Hospital for Children, Bristol, UK
3. Research in Emergency Care Avon Collaborative Hub (REACH), University of the West of England, Bristol, UK
4. Paediatric Emergency Department, Alder Hey Children's NHS Foundation Trust, Liverpool, UK
5. Department of Paediatric Surgery, St George's University Hospitals NHS Foundation Trust, London, UK
6. Centre for Academic Child Health, University of the West of England, Bristol, UK

*Corresponding author: Professor Nigel J Hall, n.j.hall@soton.ac.uk, University Surgery Unit, University of Southampton Faculty of Medicine, Southampton, SO16 6YD

Competing interests: None declared.

Funding statement: This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Contribution statement: Study conception and design: JJN, MDL, SM, SP, JM, NJH. Data collection: MAGNETIC study collaborators. Data analysis and interpretation of results: JJN, MDL, SM, JM, NJH. Draft manuscript preparation: JJN. All authors reviewed the results and approved the final version of the manuscript. JJN and NJH are responsible for the overall content as guarantors.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Objective Magnet ingestion in children and young people (CYP) is associated with significant harm. We aimed to describe the incidence, circumstances and outcomes of magnet ingestion in CYP in the United Kingdom (UK).

Design Prospective multicentre observational surveillance study.

Setting UK secondary and tertiary level hospitals in urban and rural settings.

Patients CYP ≤16 years of age who ingested ≥1 magnet.

Interventions Data were collected regarding demographics, circumstances surrounding ingestion, clinical features, and management. The primary outcome was the incidence of magnet ingestion in the UK.

Results Between 01/05/2022-30/04/2023, 366 cases of magnet ingestion were recorded, of which 314 met eligibility (median age 8.7 years [IQR 5.1–12.0]). The incidence of magnet ingestion in the UK was at least 2.4/100,000 (95% CI 2.2 – 2.7) CYP per year. CYP sourced magnets from toys (38%), and magnet products were predominantly purchased by parents or caregivers (19%). Magnet-related injuries occurred in 23 (7%) cases and surgery was undertaken in 32 (10%). Single magnet ingestions did not cause magnet-related injury. Swallowing greater numbers of magnets associated with an increased risk of injury (OR 1.1 [95% CI 1.0–1.2], p=0.002). CYP were asymptomatic in 75% of cases, but clinical features on presentation associated with an increased risk of injury (OR 3.8 [95% CI 1.4–10.3], p=0.008).

Conclusions Whilst magnet ingestion in children is uncommon, ingestion of multiple magnets can cause injuries requiring surgery. Greater public and clinician awareness of the associated risks is warranted. This study can inform public health interventions and evidence-based guidelines.

KEY MESSAGES

Evidence before this study

Magnets are dangerous when ingested by children and young people (CYP). Up to 50% of CYP who ingest multiple magnets require endoscopic or surgical intervention. Significant morbidity, including bowel resection and stoma formation, occurs in up to 10%. Multiple studies in Europe and North America have observed an apparent increase in magnet ingestions, but no study has investigated the incidence and circumstances surrounding magnet ingestion in the United Kingdom (UK).

What this study adds

The incidence of magnet ingestions in this study was 2.4/100,000 CYP per year. This is likely to be a conservative estimate as not all UK centres contributed data. Magnets were typically sourced from children's toys and products containing magnets are predominantly purchased by parents or caregivers. Social media was directly implicated in 6% of ingestions. Most CYP were asymptomatic, but the presence of clinical features on presentation was associated with a four-fold higher risk of magnet-related injury. Initial radiological imaging rarely showed complications and in a minority of CYP the reported number of magnets ingested was different to the number identified on imaging. No single magnet ingestions resulted in a magnet-related injury. Half of all cases involved the ingestion of multiple magnets, which caused magnet-related injuries in 14%.

How this study might affect research, practice or policy

Greater public awareness of the dangers of magnet ingestion is necessary and warnings at the point of sale appear an insufficient measure to prevent CYP accessing and ingesting magnets. A co-ordinated effort by healthcare professionals, government agencies and relevant charity groups to influence regulations on the sale of high-power magnet products is warranted. Data from the MAGNETIC study can be used to inform evidence-based management guidelines for ingested magnetic foreign bodies. Single magnet ingestions may be treated differently to ingestions of multiple magnets, with fewer radiological investigations and less intensive follow-up. With multiple magnet ingestions, healthcare professionals must be aware of the potential for complications at any time until magnets have passed and should not be reassured by an absence of clinical features or complications on initial imaging. CYP who ingest a greater number of magnets and those with clinical features on presentation are at a higher risk of magnet-related injury.

INTRODUCTION

Foreign body ingestion by children and young people (CYP) is common. Young children place items in their mouth when exploring their environment and older children may accidentally or intentionally swallow objects¹. Inert items typically pass through the gastrointestinal tract without complications, unless large or sharp. Certain objects interact with gastrointestinal tract tissues to cause harm. Magnets are particularly dangerous when more than one is ingested, as they can attract one another from different locations within the gastrointestinal tract². This can cause bowel obstruction, pressure necrosis, fistulation, or perforation, requiring endoscopic or surgical treatment in up to 60% of children^{3,4}.

Identifying and treating CYP with complications of magnet ingestion is challenging, as they may not report the event to caregivers and can present late to healthcare services^{5,6}. Patients are often asymptomatic, and the decision to intervene is based on identifying complications on radiographs⁷. Non-progression of magnets on serial radiographs is usually interpreted as signifying entrapment of bowel between magnets or fistulation^{3,7}. However, assessment of magnet progression is subjective and may be unreliable⁸. Current guidelines suggest repeating radiographs every 6–12 hours, or 2–3 days if the child is asymptomatic^{9,10}. However, there remains variability in management, which results in a high burden of investigations, radiation exposure and resource use^{3,4}. Early identification of CYP at risk of complications may limit morbidity, prevent unnecessary investigations, and reduce the need for intervention.

Several studies have observed an apparent increase in presentations to healthcare services following magnet ingestion^{2,11–13}. This may be due to the influence of social media, increasing availability of high-power magnets, or other factors¹⁴. However, the incidence of magnet ingestions, sources of these magnets, circumstances surrounding ingestions, and variation in management in the United Kingdom (UK) are unknown. These data are important to inform public health policy actions to limit harm from magnets and enable the development of evidence-based guidelines.

In this prospective observational surveillance study, our primary aim was to determine the incidence of magnet ingestion in CYP in the UK. Secondary objectives included examining the circumstances surrounding magnet ingestion, the investigations and management strategies utilised, and patient outcomes.

METHODS

This prospective observational surveillance study was undertaken in Paediatric Emergency Research in the UK and Ireland (PERUKI) sites, representing secondary and tertiary level hospitals in urban and rural settings¹⁵. CYP ≤ 16 years of age, who presented between 01/05/2022–30/04/2023, reporting the ingestion of ≥ 1 magnetic foreign body, were eligible for inclusion. Patients were excluded if they had ingested only non-magnetic objects or had magnetic foreign bodies only outside the gastrointestinal tract.

Due to the rarity of magnet ingestion and the necessity to report all cases, we sought approval from the Confidentiality Advisory Group to collect data without patient consent (22CAG0031). Patients and the public were involved in the study design and were consulted on the appropriateness of opt-out consent. Approvals were obtained from the UK Health Research Authority (22/NW0050). This study is reported in accordance with STROBE¹⁶.

Data were collected and managed in a secure online Research Electronic Data Capture tools (REDCap) database¹⁷. Participants were identified and screened by the clinical teams, who uploaded prospective data. Participation did not change management decisions made by the local clinical team.

On first presentation to hospital, treating healthcare professionals (HCP) completed a case report form (CRF) detailing demographic data, circumstances of ingestion, clinical features, initial investigations, and management. Further investigations, management and outcome data were collected at seven and 28-days post-presentation. Reminders to complete CRF were sent to the initial treating HCP. If a CYP was transferred between HCP at a single hospital, or between different hospitals, all teams were asked to complete a CRF to capture complete incident data. The patient's unique NHS number was used to consolidate duplicate entries.

Socioeconomic status was defined using nation-specific index of multiple deprivation (IMD) data from 2017 (Northern Ireland), 2019 (England and Wales) and 2020 (Scotland). Successful endoscopic or surgical treatment was defined as removal of all magnetic foreign bodies, so that the patient required no further interventions. Magnet-related injury was defined as harm occurring at any time as a direct result of magnet ingestion.

The primary outcome was the minimum incidence of magnet ingestions in CYP in the UK during the one-year surveillance period. Secondary outcomes included determining whether sociodemographic and clinical factors were associated with the ingestion of magnets and the development of magnet-related injuries.

Prior to analysis, all entries were cleaned and validated. Duplicates were consolidated. Records of patients not meeting eligibility and records with insufficient data to inform the study primary outcome were removed.

Incidence was calculated using mid-2021 UK population census data for CYP aged 0–16 years and 95% confidence intervals (CI) were calculated assuming a Poisson distribution. Skewed data are reported as median and interquartile range (IQR) and were compared using the Mann-Whitney U test. Categorical data were compared using Chi-squared or Fisher exact tests, as appropriate. Correlation was assessed using Spearman’s rank correlation coefficient. Logistic regression analysis was undertaken to identify factors associated with magnet-related injury. Data are described as odds ratios (OR) with 95% CI. All tests were two-sided. A p-value <0.05 was considered significant. Analysis was conducted in SPSS (v29.0.0) and GraphPad (v10.2.3).

RESULTS

Patient cohort

A total of 366 records were uploaded to the database. Thirty-eight duplicate entries were consolidated. Eleven records not meeting eligibility and three records with insufficient data were excluded. The final dataset for analysis included 314 unique episodes of magnet ingestion, representing data from 66 hospitals (Table 1).

The incidence of magnet ingestion in our study was 2.4/100,000 CYP per year (CI 2.2–2.7). Of the 314 presentations, 262/314 (83%) cases were reported from England, 41/314 (13%) from Scotland, 6/314 (2%) from Wales and 5/314 (2%) from Northern Ireland.

CYP reported how many magnets they had ingested in 301/314 cases (96%). The median number of magnets ingested was 2 (IQR 1–2, range 1–52). Single magnet ingestions were reported by 137/301 (46%) and multiple magnet ingestions by 164/301 (54%).

Socio-demographics and circumstances

Median age was 8.7 years (IQR 5.1–12.0) and 164/314 (52%) cases were male (Table 1, Supplementary Figure 1). The highest proportion of ingestion cases was in children aged 12 years. The median IMD decile was five (IQR 2–8) and the highest proportion of ingestions was decile one (16%) (Supplementary Figure 2). A non-significant negative correlation was observed between increasing IMD decile and frequency of magnet ingestions

(Spearman's $\rho = -0.50$ [CI -0.86–0.21], $p = 0.143$). Of the 294/314 (94%) cases for which information on previous involvement with social services was available, 36 children (12%) had prior involvement.

Fourteen CYP (4%) had an existing diagnosis of anxiety or depression, whilst 28/314 (9%) had autism spectrum disorder (ASD), and 7/314 (2%) had attention deficit hyperactivity disorder (ADHD). Six had previously ingested magnetic foreign bodies, of whom two had a diagnosis of depression and three had a diagnosis of ASD. All six had previously committed acts of self-harm, and four had previously ingested non-magnetic foreign bodies.

Magnets were sourced from toys or games in 118/314 (38%) cases (Table 2). Magnet containing products had been purchased by parents or caregivers in 60/314 (19%). Social media was directly implicated in 20/314 (6%) cases: 17/20 (85%) were female and 19/20 (95%) ingested multiple magnets. All 20 referred to mimicking a tongue piercing with magnets. Twelve specifically mentioned the social media site "TikTok".

Presentation and investigations

In 174/314 (55%) cases, the primary point of healthcare contact was attending ED. Secondary presentation to ED after discussion with emergency services or formal phone/online advice (NHS 111) occurred in 44/314 (14%) and 39/314 (12%) cases respectively. Two (1%) attended ED after seeing their General Practitioner. Fourteen (8%) cases were transfers from another hospital and 10 (3%) were planned reviews (either from the same or a different hospital). All inter-hospital transfers were for paediatric surgery services.

Approximate date and time of magnet ingestion was reported in 306/314 cases (97%). The median time from magnet ingestion to presentation was 2.4 hours (IQR 1–8). The longest time from ingestion to presentation was 22 days.

Most CYP had no clinical features; only 80/314 (25%) had at least one symptom or sign (Supplementary Table 1). The commonest clinical features were abdominal pain (48/314, 15%) and tenderness (20/314, 6%). Children who had ingested multiple magnets (confirmed radiologically) were not more likely to have clinical features on presentation (11% for single *versus* 14% for multiple, $p = 0.404$).

The median number of magnets identified on imaging was 2 (IQR 1–2). Single magnets were identified in 123/314 (39%) cases and multiple magnets in 170/314 (54%). Eight (3%) received no diagnostic imaging to confirm magnet ingestion; all of these were suspected single magnet ingestions (Supplementary Table 2).

Five who reported multiple magnet ingestions had a single magnet or no magnets on imaging. Conversely, 11 who had reported single magnet ingestions had multiple magnets identified on imaging. Initial imaging identified complications in one child who developed small bowel obstruction after ingesting multiple magnets (performed 17 days after ingestion). No other complications were identified on presentation imaging.

Management of magnet ingestion

After ED management (Figure 1), 141/314 (45%) cases underwent further imaging, of which 139/141 (99%) had plain radiographs (Supplementary Table 2). In 30/146 (21%) cases the magnets had not progressed from the initial imaging within seven days of ingestion, and two children had developed new complications: one perforation and one gastrointestinal fistula. Overall, the median number of plain radiographs performed per patient was 2 (IQR 0–3). Five or more radiographs were performed in 24 CYP, and four received ≥10.

Of 247/314 (79%) cases which did not require intervention, 112/314 (45%) were multiple magnet ingestions. Conservative management was more common in single magnet ingestions (94% single *versus* 67% multiple, $p<0.001$).

Endoscopy was performed in 21/314 (7%) cases, with a median time to procedure of 1 day (IQR 0 – 3). Magnets were successfully removed endoscopically in 9/21 (43%) children and median time to endoscopy was not associated with success (1 day [IQR 0–1] success *versus* 1 day [IQR 0–4] failure, $p=0.803$). Of the 12 unsuccessful cases, seven underwent surgery. There were no procedural complications.

Thirty-two (10%) CYP underwent surgery with a median time to surgery of 3 days (IQR 2–7, range 0–32) (Table 3). The most common indication was failure of magnets to progress on serial radiographs (21/32, 66%). Most underwent laparotomy (14/32, 44%), nine children underwent laparoscopic magnet removal, and nine children were converted from laparoscopy to open surgery. Injuries caused by magnet ingestion were identified in 23/314 (7%) cases intra-operatively (Table 3). Bowel resection was performed in five patients and four required parenteral nutrition. One patient developed a post-operative wound infection. Date of discharge post-surgery was recorded in 20 cases, and median length of stay post-procedure was 1 day (IQR 1–4).

Factors related to magnet-related injury

A greater number of magnets were ingested by children who developed a magnet-related injury (4 [IQR 3–12] in the injury group *versus* 2 [IQR 1–2] in the non-injury group, $p<0.001$) (Table 4). Similarly, children who developed a magnet-related injury presented later following ingestion (5 hours [IQR 2–36] in the injury group *versus* 2 [IQR 1–7] in the non-injury group, $p=0.046$). A significantly greater proportion of children with clinical features on presentation suffered a magnet-related injury compared to those with no clinical features (52% *versus* 23%, $p=0.002$). Patient age, sex and ethnicity were not significant.

On multivariable logistic regression, an association was observed between increasing numbers of ingested magnets and the risk of magnet-related injury (OR 1.1 (CI 1.0–1.2), $p=0.002$). Similarly, the presence of clinical features on presentation was associated with an increased risk of injury (OR 3.8 (CI 1.4–10.3), $p=0.008$) (Supplementary Table 3).

DISCUSSION

The incidence of magnet ingestion in the UK during the study period was at least 2.4/100,000 CYP per year. Toys and games, which are predominantly purchased by parents or caregivers, were the most common source of magnets. Delayed presentation was associated with a higher incidence of magnet-related injury, and while CYP are often asymptomatic, presence of symptoms on presentation was associated with a four-fold increased risk of injury. Magnet-related injuries occurred in 14% of multiple magnet ingestions. No magnet-related injuries were identified in single magnet ingestions but increasing numbers of ingested magnets were significantly associated with an increased risk of injury.

Low socioeconomic status is recognised as being associated with childhood injury morbidity and mortality¹⁸. We observed that the frequency of magnet ingestions was highest amongst children in the lowest IMD decile. Few studies have investigated the relationship between socioeconomic status and paediatric foreign body ingestion. Chen *et al.* observed that residential instability and maternal deprivation were independently associated with foreign body ingestion in a cohort of 680 children in Canada¹⁹. Further research is required to understand the relationship between deprivation and foreign body ingestion to identify potential interventions.

There have been several reports of magnet ingestion inspired by social media in the UK news, particularly the “viral trend” of placing magnets in the mouth to mimic tongue piercings¹³. In our study, 6% of cases implicated social media, all replicating a tongue piercing. Most were female and 95% ingested multiple magnets. Future

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

efforts to reduce magnet ingestion could include placing greater responsibility on social media companies to ensure that posted content does not promote self-injurious behaviours.

CYP that may be at a higher risk of magnet-related injuries are those with behavioural diagnoses, such as ASD and ADHD²⁰. A diagnosis of ASD was over-represented in the CYP included in this study, which corroborates previous observations³. Similarly, a cohort of CYP with mental health diagnoses who ingest magnets as a form of self-harm has been identified²¹. Caregivers to CYP in these high-risk groups should be particularly aware of the dangers of magnet ingestion. Manufacturers of products targeted to these groups should also explore methods to minimise risk.

This study confirms several key findings regarding the management of magnet ingestion in CYP. When presenting to healthcare services following magnet ingestion, patients are typically asymptomatic²². The presence of symptoms does not reliably distinguish between single and multiple magnet ingestions. Our findings corroborate research that demonstrates an association between symptoms and signs, and an increased risk of magnet-related injury²³. Plain radiographs rarely identify complications on presentation and normal imaging findings should not reassure HCP. No cases of single magnet ingestion resulted in magnet-related injuries, confirming that single magnet ingestions can be managed differently to multiple^{5,9,10}. Although, single magnets may still require intervention if causing obstruction or the child is symptomatic³.

The strengths of the MAGNETIC study are its large sample and prospective data collection by HCP. However, the incidence is likely to be an underestimation. Despite our best efforts not all hospitals in the UK contributed data and therefore all cases of magnet ingestion may not have been captured in this dataset. No internal audits were performed at any participating sites to confirm complete data collection. Similarly, due to geographic variability in reporting, IMD data may not be nationally representative. Limitations of the study also include the presence of unknown or missing datapoints for secondary outcomes.

This study has confirmed previous reports that multiple magnets are dangerous when ingested by children. Single magnets do not appear to cause injuries when ingested, and therefore HCP should have a high index of suspicion for multiple magnet ingestions. Despite current legislation mandating warnings at the point of sale, magnet ingestions continue to occur. Further work must focus on raising public awareness of this issue and

standardising guidelines based on published evidence. A co-ordinated effort by HCP, government agencies and relevant charity groups to limit the sale of high-power magnets is warranted.

REFERENCES

- 1 Speidel AJ, Wölfle L, Mayer B, Posovszky C. Increase in foreign body and harmful substance ingestion and associated complications in children: a retrospective study of 1199 cases from 2005 to 2017. *BMC Pediatr* 2020; **20**: 560.
- 2 Middelberg LK, Funk AR, Hays HL, McKenzie LB, Rudolph B, Spiller HA. Magnet Injuries in Children: An Analysis of the National Poison Data System from 2008 to 2019. *J Pediatr* 2021; **232**: 251-256.e2.
- 3 Paediatric Surgery Trainee Research Network. Magnet and button battery ingestion in children: multicentre observational study of management and outcomes. *BJS Open* 2022; **6**: zrac056.
- 4 De Roo AC, Thompson MC, Chounthirath T, *et al.* Rare-Earth Magnet Ingestion–Related Injuries Among Children, 2000-2012. *Clin Pediatr (Phila)* 2013; **52**: 1006–13.
- 5 Miyamoto R, Okuda M, Kikuchi S, Iwayama H, Hataya H, Okumura A. A nationwide questionnaire survey on accidental magnet ingestion in children in Japan. *Acta Paediatr* 2021; **110**: 314–25.
- 6 Maksimyan S, Bernshteyn M, Ravi SJK, Srivatana U. Clinical course and management of an unknown multiple-magnet ingestion in a teenage male. *BMJ Case Rep* 2023; **16**: e256418.
- 7 Otjen JP, Rohrmann CA, Iyer RS. Imaging pediatric magnet ingestion with surgical-pathological correlation. *Pediatr Radiol* 2013; **43**: 851–9.
- 8 Paediatric Surgery Trainee Research Network. Inter-rater agreement of serial radiographs to identify non-progression of ingested multiple magnets in children. *Arch Dis Child* 2024; **109**: 172.
- 9 Nugud AA, Tzivinikos C, Assa A, *et al.* Pediatric Magnet Ingestion, Diagnosis, Management, and Prevention. *J Pediatr Gastroenterol Nutr* 2023; **76**: 523–32.
- 10 Stibbards S, Kenney S, Emsden S, *et al.* Royal College of Emergency Medicine Best Practice Guidelines: Ingestion of Super Strong Magnets in Children. 2021.
- 11 Flaherty MR, Buchmiller T, Vangel M, Lee LK. Pediatric Magnet Ingestions After Federal Rule Changes, 2009-2019. *JAMA* 2020; **324**: 2102–4.
- 12 Thakkar H, Burnand KM, Healy C, *et al.* Foreign body ingestion in children: a magnet epidemic within a pandemic. *Arch Dis Child* 2021; **106**: 1240–1.
- 13 McLeish S, Harwood R, Decker E, Almond S, Hall NJ, Durand C. Managing magnets: An audit of introduction of the Royal College of Emergency Medicine Best Practice Guideline. *Acta Paediatr* 2024; **113**: 127–34.
- 14 Middelberg LK, Leonard JC, Shi J, *et al.* Warning Labels and High-Powered Magnet Exposures. *Pediatrics* 2022; **150**: e2022056325.

15 Lyttle MD, O’Sullivan R, Hartshorn S, *et al.* Pediatric Emergency Research in the UK and Ireland (PERUKI): developing a collaborative for multicentre research. *Arch Dis Child* 2014; **99**: 602–3.

16 Elm E von, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. Strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ* 2007; **335**: 806.

17 Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009; **42**: 377–81.

18 Birken CS, MacArthur C. Socioeconomic status and injury risk in children. *Paediatr Child Health* 2004; **9**: 323–5.

19 Chen T, M Siu J, Madan Y, *et al.* Pediatric Esophageal Foreign Bodies: The Role of Socioeconomic Status in Ingestion Patterns. *Laryngoscope* 2024; **n/a**. DOI:<https://doi.org/10.1002/lary.31274>.

20 Rashid F, Davies L, Iftikhar SY. Magnetised intragastric foreign body collection and autism: An advice for carers and literature review. *Autism* 2010; **14**: 139–45.

21 Low Kapalu C, Lantos J, Booser A, Thomson M, Attard T. Preventing Self-Harm From Repeat Foreign-Body Ingestion. *Pediatrics* 2020; **145**: e20191515.

22 Price J, Malakounides G, Stibbards S, Agrawal S. Ball magnet ingestion in children: a stronger and more dangerous attraction? *Emergency Medicine Journal* 2022; **39**: 467.

23 Zhang RY, Cai P, Zhang TT, *et al.* Clinical predictors of surgical intervention for gastrointestinal magnetic foreign bodies in children. *BMC Pediatr* 2023; **23**: 323.

DATA SHARING

Anonymised data from MAGNETIC will be made available upon reasonable request to the authors.

FIGURES

Figure 1: Summary of emergency department (ED) management (n = 304). Of the 115 patients discharged home without follow-up, 18 (16%) were radiologically confirmed multiple magnet ingestions. No patients discharged from the ED without follow-up underwent subsequent surgical intervention.

TABLES

Table 1: Characteristics of patients on presentation to a healthcare professional. IQR – interquartile range.

All patients (n = 314)	
Median age at presentation in years (IQR)	8.7 (5.1 – 12.0)
Sex	
Female	150 (48%)
Male	164 (52%)
Ethnicity	
Asian or Asian British	31 (10%)
Black, Black British, Caribbean or African	4 (1%)
Mixed or Multiple Ethnic Groups	4 (1%)
Other Ethnic Group	34 (11%)
White	241 (77%)
History of social services involvement	
Yes	36 (12%)
No	258 (82%)
Unknown/unspecified	20 (6%)
Siblings (n = 141)	
Yes	111 (79%)
No	30 (21%)

Table 2: Summary of circumstances of magnet ingestion.

All patients (n = 314)	
Reason ingestion was identified	
Witnessed by parent/caregiver	50 (16%)
Witnessed by another person	21 (7%)
Reported by patient	188 (60%)
Unwitnessed and unreported	12 (4%)
Patient symptomatic	13 (4%)
Other	13 (4%)
Social media implicated	
Yes	20 (6%)
No	285 (91%)
Unknown/unspecified	6 (2%)
Ingestion intentional	
Yes	62 (20%)
No	248 (79%)
Unknown/unspecified	4 (1%)
Source of magnets	
Toys or games	118 (38%)
Home devices	4 (1%)
Jewellery	24 (8%)
Executive toys	0 (0%)
Other	32 (10%)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Unknown/unspecified		134 (43%)
Purchaser of magnets		
	Patient	6 (2%)
	Parent/caregiver	60 (19%)
	Siblings	2 (1%)
	Friends	9 (3%)
	Other	15 (5%)
	Unknown/unspecified	222 (71%)

Confidential: For Review Only

Table 3: Characteristics of 32 children managed surgically. XR – plain film radiographs, PN – parenteral nutrition via a central venous catheter.

Age (yrs)	Sex	Number Magnets Ingested	Indication for Surgery	Procedure	Surgical Findings and Magnet-Related Injury	Type of Magnets Removed	Surgical Complication
1.7	Male	3	Clinical and radiological suspicion of complications	Laparoscopy converted to open	Ileal and caecal obstruction and perforation	High-power ball magnets	None
1.8	Female	2	Failure to progress on XR	Laparotomy	Small bowel perforation	High-power ball magnets	None
1.8	Male	9	Clinical suspicion of complications	Laparotomy	Small and large bowel perforations	High-power ball magnets	None
2	Male	3	Failure to progress on XR	Laparoscopy converted to open	Small and large bowel mucosal damage	High-power ball magnets	Wound infection
2.7	Female	3	Failure to progress on XR	Laparoscopy	None – removed from stomach	High-power ball magnets	None
2.8	Male	3	Failure to progress on XR	Laparotomy	Fistulation between posterior stomach and splenic flexure of large bowel	High-power disc magnets	None
2.8	Male	18	Failure to progress on XR	Laparotomy	None – removed from small bowel	High-power ball magnets	None
3.1	Female	1	Clinical suspicion of complications	Laparoscopy converted to open	None – removed from large bowel	High-power disc magnet	None
3.9	Male	4	Failure to progress on XR	Laparoscopy	None – removed from caecum via appendix	High-power ball magnets	None
4.7	Female	11	Clinical and radiological suspicion of complications and failure to progress on XR	Laparoscopy converted to open	Fistulation between duodenum and terminal ileum, received 7 days of PN	High-power bar magnets	None
4.8	Male	22	Clinical suspicion of complications	Laparotomy	Stomach obstruction	Fridge magnets	None
4.9	Male	52	Clinical and radiological suspicion of complications and failure to progress on XR	Laparotomy	Small bowel perforation with intra-abdominal abscess, fistulation between jejunum and ileum, and ileum and caecum, bowel resection, received 7 days of PN	High-power ball magnets	None
5.5	Male	20	Clinical suspicion of complications	Laparoscopy	Fistulation jejunum to jejunum, bowel resection	High-power ball magnets	None
6.7	Male	13	Failure to progress on XR	Laparoscopy	Fistulation small bowel to small bowel, bowel resection	High-power ball magnets	None
8.1	Male	4	Clinical suspicion of complications and failure to progress on XR	Laparoscopy converted to open	Small bowel obstruction, bowel resection	Disc magnets	None
8.3	Female	2	Failure to progress on XR	Laparoscopy	Ileo-caecal valve obstruction	High-power ball magnets	None
8.8	Male	2	Failure to progress on XR	Laparoscopy	None – removed from caecum	High-power ball magnets	None
9.2	Female	2	Failure to progress on XR	Laparoscopy	None – removed from caecum	High-power ball magnets	None

9.6	Female	18	Clinical and radiological suspicion of complications	Laparotomy	Small bowel mucosal damage, received 7 days of PN	High-power ball magnets	None
9.6	Male	1	Clinical suspicion of complications	Laparotomy	None – removed from stomach	Disc magnet	None
11	Male	4	Clinical suspicion of complications and failure to progress on XR	Laparoscopy converted to open	Fistulation between posterior stomach and jejunum, received 3 days of PN	High-power ball magnets	None
11.1	Female	9	Clinical suspicion of complications	Laparoscopy	None – magnets removed from large bowel	High-power disk magnets	None
12	Male	2	Failure to progress on XR	Laparotomy	Large bowel mucosal damage	High-power ball magnets	None
12	Female	4	Failure to progress on XR	Laparotomy	Small bowel perforation	High-power ball magnets	None
12.3	Female	4	Clinical suspicion of complications and failure to progress on XR	Laparotomy	Fistulation between duodenum and transverse colon	High-power ball and disc magnets	None
12.3	Female	3	Radiological suspicion of complications	Laparoscopy converted to open	Small bowel mucosal damage	High-power ball magnets	None
12.3	Female	3	Failure to progress on XR	Laparoscopy converted to open	Small bowel perforation	High-power disc magnets	None
12.8	Male	4	Failure to progress on XR	Laparoscopy	None – magnets had passed	Unspecified	None
13.6	Male	5	Clinical suspicion of complications and failure to progress on XR	Laparoscopy converted to open	Stomach obstruction	High-power disc magnets	None
13.9	Male	15	Radiological suspicion of complications	Laparoscopy converted to open	Small and large bowel mucosal damage	High-power ball magnets	None
16.3	Male	7	Failure to progress on XR	Laparotomy	Fistulation between stomach and small bowel, and small bowel to rectum	High-power disc magnets	None
16.8	Female	7	Clinical suspicion of complications	Laparotomy	Fistulation between ileum and ileum, bowel resection	High-power disc magnets	None

Confidential: For Review Only

Table 4: Factors associated with magnet-related injury. IQR – interquartile range.

	Injury (n = 23)	No Injury (n = 291)	p-value
Median age in years (IQR)	8.3 (4.8 – 12.3)	8.7 (5.3 – 11.9)	0.975
Male:female	14:9	150:141	0.389
Ethnicity			0.550
White	18 (78%)	223 (77%)	
Asian	1 (4%)	30 (10%)	
Black	0 (0%)	4 (1%)	
Mixed	1 (4%)	3 (1%)	
Other	3 (13%)	31 (11%)	
Social services involvement	3 (13%)	33 (11%)	0.805
Median IMD decile (IQR) (n = 291)	6 (3 – 8)	5 (2 – 8)	0.320
Past medical history			
Mental health diagnosis	1 (4%)	9 (3%)	0.741
Behavioural diagnosis	5 (22%)	28 (10%)	0.068
Median number of magnets ingested (IQR)	4 (3 – 12)	2 (1 – 2)	<0.001
Median hours to presentation (IQR) (n = 306)	5 (2 – 36)	2 (1 – 7)	0.046
Presence of clinical features	12 (52%)	68 (23%)	0.002
Location of magnets on initial imaging (n = 307)			0.872
Oesophagus	0 (0%)	6 (2%)	
Stomach	6 (26%)	105 (36%)	
Small bowel	9 (39%)	96 (33%)	
Large bowel	5 (22%)	39 (13%)	
Oesophagus and stomach	0 (0%)	3 (1%)	
Oesophagus and large bowel	0 (0%)	1 (0.3%)	
Stomach and small bowel	0 (0%)	17 (6%)	
Small bowel and large bowel	1 (4%)	15 (5%)	
Stomach, small and large bowel	0 (0%)	2 (1%)	
All	0 (0%)	2 (1%)	
Median days to surgery (IQR)	3 (2 – 5)	6 (2 – 7)	0.694
Intervention			
Endoscopy	5 (22%)	16 (5%)	0.003
Surgery	23 (100%)	9 (3%)	<0.001




Figure 1: Summary of emergency department (ED) management (n = 304). Of the 115 patients discharged home without follow-up, 18 (16%) were radiologically confirmed multiple magnet ingestions. No patients discharged from the ED without follow-up underwent subsequent surgical intervention.

159x98mm (220 x 220 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

MAGNETIC STUDY COLLABORATORS


Collaborator Name (affiliation)

Holly Brooker (Torbay and South Devon NHS Foundation Trust), Rachel Varnam (Torbay and South Devon NHS Foundation Trust), Daniel Putt-Willis (Queen Alexandra Hospital, Portsmouth), Matthew Smith (Queen Alexandra Hospital, Portsmouth), Lotty Smith (Queen Alexandra Hospital, Portsmouth), Ibrahim Yusuf (Queen Alexandra Hospital, Portsmouth), Natasha Dean (Queen Alexandra Hospital, Portsmouth), David Patel (Queen Alexandra Hospital, Portsmouth), Mahbuba Rahman (Queen Alexandra Hospital, Portsmouth), Elesha Vooght (Queen Alexandra Hospital, Portsmouth), Refat Parveen (Queen Alexandra Hospital, Portsmouth), Luke Shirley-Mansell (Queen Alexandra Hospital, Portsmouth), Rosie Cresner (Southampton Children’s Hospital), Thomas Cromarty (Southampton Children’s Hospital), Rebecca Broomfield (Southampton Children’s Hospital), Jane Bayreuther (Southampton Children’s Hospital), George Bethell (Southampton Children’s Hospital), Christina Major (Southampton Children’s Hospital), Jason Barling (Southampton Children’s Hospital), Victoria Wilson (Royal Belfast Hospital for Sick Children), Julianne Maney (Royal Belfast Hospital for Sick Children), Kathryn Wilson (Royal Belfast Hospital for Sick Children), Daniel Ratnaraj (Whipp’s Cross Hospital), Divya Divakaran (Whipp’s Cross Hospital), Jenny Hickey (Newham University Hospital), Darren Ranasinghe (Epsom and St. Helier Hospitals), Andrew Foster (Epsom and St. Helier Hospitals), Benjamin Martin (Bristol Royal Hospital for Children), Robert Walker (Bristol Royal Hospital for Children), Catherine Jones (Bristol Royal Hospital for Children), Edward Soans (Bristol Royal Hospital for Children), Alice Monk (Bristol Royal Hospital for Children), Alma Rahman (Bristol Royal Hospital for Children), Erin Jones (Bristol Royal Hospital for Children), Katherine Tambudze (Bristol Royal Hospital for Children), David Hopgood (Bristol Royal Hospital for Children), Alice Downes (Leeds General Infirmary), Tahmina Nasreen (Leeds General Infirmary), Sasha Preskey (Leeds General Infirmary), Jack Long (Leeds General Infirmary), Joshua Adamson (Leeds General Infirmary), Roger Henderson (Leeds General Infirmary), Hannah Andreassen (Leeds General Infirmary), Hannah Lee Chadwick (Leeds General Infirmary), Gina Towart (South Tyneside and Sunderland NHS Foundation Trust), Khalid Abdelhafiz (South Tyneside and Sunderland NHS Foundation Trust), Elizabeth O’Connor (Great North Children’s Hospital, Newcastle-upon-Tyne), Darren Carlyle (South Tyneside and Sunderland NHS Foundation Trust), L Tubman (South Tyneside and Sunderland NHS Foundation Trust), Katy Wallace (South Tyneside and Sunderland NHS Foundation Trust), Ahmed Mohamed (Alder Hey Children’s Hospital, Liverpool), Sarah Siner (Alder Hey Children’s Hospital,

Liverpool), Sarah Fissler (Alder Hey Children's Hospital, Liverpool), Sally Mcleish (Alder Hey Children's Hospital, Liverpool), Meriel Tolhurst-Cleaver (Alder Hey Children's Hospital, Liverpool), Stephen Fletcher (Alder Hey Children's Hospital, Liverpool), Michelle Russell (Alder Hey Children's Hospital, Liverpool), Katherine Winrow (Alder Hey Children's Hospital, Liverpool), Josef Taylor (Alder Hey Children's Hospital, Liverpool), Angela Armitage (Alder Hey Children's Hospital, Liverpool), Karen Geoghegan (Alder Hey Children's Hospital, Liverpool), Rheanan Buckle (Alder Hey Children's Hospital, Liverpool), Sarah Wood (Alder Hey Children's Hospital, Liverpool), Laura Tremarco (Alder Hey Children's Hospital, Liverpool), Vicky Collins (North Tees and Hartlepool Foundation Trust), Dawn Egginton (North Tees and Hartlepool Foundation Trust), George Simpson (North Tees and Hartlepool Foundation Trust), Sophie Dowsett (St George's University Hospitals NHS Foundation Trust), Florin Djendow (St George's University Hospitals NHS Foundation Trust), Heather Jarman (St George's University Hospitals NHS Foundation Trust), Klata Edyta (Colchester Hospital), Michelle Dotchin (Colchester Hospital), Sarah Potter (Royal United Hospital, Bath), Kanya Kamaraj (Wexham Park Hospital), Anas Fagelnor (John Radcliffe Hospital, Oxford), Christopher Dadnam (Leicester Royal Infirmary), Ayeesha Shafiq (Leicester Royal Infirmary), Sophie Lewis (Royal Manchester Children's Hospital), Iyad Zarifa (Royal Manchester Children's Hospital), Ross Craigie (Royal Manchester Children's Hospital), Patrick Aldridge (Frimley Park Hospital), Nirubhan Veeraragavan (Frimley Park Hospital), Zena Haslam (Ormskirk and District General Hospital), Alex Carney (Hull Royal Infirmary), Grace Rimmer (Hull Royal Infirmary), Sarah Jones (Hull Royal Infirmary), Simon Richardson (Hull Royal Infirmary), Laura Riddick (Hull Royal Infirmary), Ellen McCourt (Hull Royal Infirmary), A Azad-Karim (Hull Royal Infirmary), Kate Quigley (Hull Royal Infirmary), Sarah Yassin (Hull Royal Infirmary), Vanessa Merrick (St Mary's Hospital, Imperial College Healthcare NHS Trust), Rebecca Salter (St Mary's Hospital, Imperial College Healthcare NHS Trust), Rie Yoshida (St Mary's Hospital, Imperial College Healthcare NHS Trust), John Bass (Royal Alexandra Children's Hospital, Brighton), Elizabeth Vincent (Royal Alexandra Children's Hospital, Brighton), Costa Healy (Royal Alexandra Children's Hospital, Brighton), Emily Jones (University Hospital Wales, Cardiff), Emily Ball (University Hospital Wales, Cardiff), Ahmed Azam (University Hospital Wales, Cardiff), Elizabeth Ryan (University Hospital Wales, Cardiff), Sofia Bedoya (University Hospital Wales, Cardiff), Sophie Keers (University Hospital Lewisham, London), Eoin Blaney (University Hospital Lewisham, London), Phil Peacock (Great Western Hospital, Swindon), Stuart Hartshorn (Birmingham Children's Hospital), Victoria Cash (Birmingham Children's Hospital), Rachel Varnam (Torbay Hospital), Edward Snelson (Norfolk and Norwich University Hospital), Vanessa Coles (Norfolk and Norwich University Hospital), Amelia Stacey (Airedale NHS Foundation Trust), S Zuhairy (Great

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


North Children's Hospital, Newcastle Upon Tyne), Lauren Chandler, Joshua Pinedo (Northumbria Healthcare NHS Foundation Trust), Andrew Bradley (Salford Royal Hospital), Victoria Gate (Blackpool Victoria Hospital), Natalie Sanlon, Uday Juttiga, Adrian Marsh (Shrewsbury & Telford NHS Trust), Chinedu Okeke (Shrewsbury and Telford NHS Trust), Nasra Ali (Shrewsbury and Telford NHS Trust), Shammi Ramlakhan (Sheffield Children's NHS Foundation Trust), Thejasvi Subramanian (Sheffield Children's NHS Foundation Trust), Verity Haffenden, Joana Obire, David Hartin (Ipswich Hospital), Nadine Darlow (Ipswich Hospital), Deborah Beeby (Ipswich Hospital), Rebecca Francis (Ipswich Hospital), Suvradeep Basu (Bradford Royal Infirmary), Amulya Saxena (Chelsea and Westminster NHS Foundation Trust), Renos Jeropoulos (Chelsea and Westminster NHS Foundation Trust), Ashleigh Hegan (Royal Hospital for Children & Young People, Edinburgh), Jen Browning (Royal Hospital for Children & Young People, Edinburgh), Elliot Craven (Royal Hospital for Children, Glasgow), Steven Foster (Royal Hospital for Children, Glasgow)



Supplementary Figure 1: Histogram showing the frequency of magnet ingestions by patient age at presentation.

108x78mm (144 x 144 DPI)

Frequency of Magnet Ingestions by IMD Decile



Supplementary Figure 2: Histogram showing the frequency of magnet ingestions at each Index of Multiple Deprivation (IMD) decile. Data from 291 cases; median decile 5 (interquartile range 2 – 8).

120x82mm (144 x 144 DPI)

SUPPLEMENTAL FIGURES

Supplementary Figure 1: Histogram showing the frequency of magnet ingestions by patient age at presentation.

Supplementary Figure 2: Histogram showing the frequency of magnet ingestions at each Index of Multiple Deprivation (IMD) decile. Data from 291 cases; median decile 5 (interquartile range 2 – 8).

SUPPLEMENTAL TABLES

Supplementary Table 1: Clinical features.

		All patients (n = 314)
Symptoms	Abdominal pain	48 (15%)
	Vomiting	14 (5%)
	Chest pain	4 (1%)
	Globus sensation	4 (1%)
	Cough/choking	6 (2%)
	Throat pain	3 (1%)
	Other	7 (2%)
	None	243 (77%)
Signs	Abdominal tenderness	20 (6%)
	Tachycardia (age defined)	10 (3%)
	Abdominal distention	3 (1%)
	Abdominal guarding/peritonism	3 (1%)
	Pyrexia (>37.5°C)	2 (1%)
	Other	2 (1%)
	None	281 (90%)

Supplemental Table 2: Imaging modalities used and magnet locations. CXR – chest plain film radiograph, AXR – abdominal plain film radiograph, CT - Computerised tomography. *Patients may receive multiple diagnostic imaging tests. **Magnets may be identified in more than one anatomical location.

	All patients (n = 314)
Diagnostic imaging modality used*	
Metal detector	41 (13%)
Anteroposterior CXR	142 (45%)
Lateral CXR	21 (7%)
Anteroposterior AXR	233 (74%)
Lateral AXR	83 (26%)
Ultrasound scan	1 (0.3%)
CT scan	1 (0.3%)
Endoscopy	2 (1%)
None	8 (3%)
Location of magnets on presentation**	
Oesophagus	13 (4%)
Stomach	135 (43%)
Small bowel	141 (45%)
Large bowel	66 (21%)
Further imaging studies within 7 days (n = 141)*	
XR	139
Contrast study	5
Ultrasound scan	1
CT scan	2

Supplemental Table 3: Multivariate analysis of factors associated with magnet-related injury.

	Odds Ratio	95% Confidence Interval	p-value
Time from ingestion to presentation	1.01	0.99 – 1.01	0.112
Number of magnets ingested	1.10	1.04 – 1.17	0.002
Clinical features on presentation	3.81	1.41 – 10.27	0.008

Reviewer: 1

Many thanks for this paper. This is an excellent study identifying the current burden of disease in the UK caused by magnets. The association with social deprivation identified is particularly interesting and I would encourage the authors to now look at ways in which this can be further addressed to look at more targeted intervention in lower socioeconomic groups.

Thank you for your kind comments on our study. We intend to conduct further work investigating parent and caregiver awareness of the dangers of magnet ingestion in the UK, with a focus on social deprivation.

Reviewer: 2

I would like to congratulate the authors on the completion of this prospective national multi-centre study focussing on a very important area of paediatric care. The overall premise and study set-up is excellent, and the overall message of the paper is important. I think that by addressing the following points the paper can be strengthened.

Thank you for your comments and feedback on our manuscript.

I am a little dubious about the reporting of incidence in the paper. It is reported that data were submitted from 66 hospitals. How does this compare to the number of EDs that children can present to in the UK? Was any internal audit requested from participating sites to make sure that all patients meeting the inclusion criteria were identified? If this hasn't been done I would make that clear and also state that this is a conservative estimate of incidence. The IMD data may also be influenced by the location of reporting units if this was not undertaken universally.

We have updated the manuscript to make it clear that the reported incidence is likely to be an underestimate. We have presented it in the results section as the 'incidence identified by our study', but discussed in the limitations that it is likely to under-represent the true incidence because of incomplete case reporting at participating centres, and because some centres did not participate. No internal audits were performed at any centres, and this has been added to the limitations section. The IMD result may be skewed by incomplete geographic reporting, and this has also been made clear in the limitations.

Reporting of incidence should also give a unit of time – so in this paper cases/100,000/year.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

We have given the incidence a unit of time throughout.

I would be interested to know a bit more about the age distribution – was this a parametric spread or were there different peaks (toddler vs adolescent) that may highlight where public messaging needs to be directed?

We have added a supplementary figure describing the age distribution.

The paper is reassuring for single magnet ingestion suggesting that no intervention is require. It would be useful to reference to PSTRN magnet paper where some single magnets were removed because they did not progress beyond the stomach and were very large – need to temper the ‘no intervention necessary’ message in this context, particularly in a symptomatic child.

We have highlighted the PSTRN paper that includes children with single magnet ingestions that required intervention.

Distinction between rare earth magnets and all magnets – the “Evidence before this study” section discusses rare earth magnets but you describe harm with other types of magnets too (fridge magnets!) so I think it would be helpful to be less specific about the type of magnet within the paper. Within the ‘Management of Magnet Ingestion’ section I would change ‘required surgery’ to ‘underwent surgery’ as a few patients did not have a magnet related injury found at operation.

We have updated the text in the “Evidence before this study” and “Management of magnet ingestion” sections.

In the final conclusions I think it would be helpful to highlight the need for evidence based clinical guidelines (rather than just guidelines). Potentially focussing on the particular areas of need – timing/frequency of repeated radiographs, maximum period of time that magnets should be left prior to intervening, drawbacks of our current approach to ingested magnets.

Our conclusion comments that “Further work must focus on ... standardising guidelines based on published evidence” which we feel appropriately highlights the need for evidence-based guidance. We decided to focus our discussion and conclusion on the public health messages of the study rather than the clinical management.

Reviewer: 3

This is an interesting study which describes the incidence of magnet ingestion in children in the UK and explores the relationship between factors associated with ingestion and harm. The authors outline the objectives and methodology clearly and appropriate statistical methods have been conducted. I have a few minor comments:

1. Definition of injury could be introduced more formally including details of how and when it was defined.

2. The term non-parametric refers to the statistical method and not to the data, it is preferable to refer to skewed data.

We have updated the methods section to formally define magnet-related injury and clarify the data as skewed.

Many thanks for your review and your comments.