



Closed Genome Sequence Obtained Using Hybrid Nanopore/Illumina Assembly of a *Bacillus anthracis* Isolate from an Animal-Skin-Drum-Associated Anthrax Case in the United Kingdom

Steven T. Pullan,^a Rory W. Miles,^a Kuiama Lewandowski,^a Richard Vipond^a

^aPublic Health England, National Infection Service, Salisbury, Wiltshire, United Kingdom

ABSTRACT Hybrid *de novo* assembly of Illumina/Nanopore reads produced a complete closed genome sequence of the chromosome and two virulence plasmids of a *Bacillus anthracis* isolate from a fatal anthrax case in the United Kingdom linked to imported animal skins/drums; this provides a high-quality representative sequence for this lineage.

Bacillus anthracis is the causative agent of the zoonotic disease anthrax. Despite its relatively rare occurrence in humans, it has gained notoriety due to its potential use as a bioweapon (1). In the United Kingdom, the most recent human cases have been associated with the intravenous use of contaminated heroin (2, 3) or the importation of animal hides for drum making (4). We previously characterized multiple isolates from two animal-skin-drum-associated anthrax cases in the United Kingdom, utilizing Illumina sequencing and a reference mapping-based approach to identify core genome single-nucleotide polymorphisms (SNPs). Phylogenetic analysis showed that these strains (along with another collected from a similar U.S. case [5]) formed a novel branch on the global *B. anthracis* phylogeny that was distinct from previously characterized West African strains, despite this being the likely origin of the contaminated animal skins (6). In order to produce a more complete reference sequence for this novel lineage, we used the Oxford Nanopore MinION genomic DNA sequencing kit (SQK-MAP006) on a MinION Mk1 device, with 1 µg of DNA (isolated as described previously) as input to produce 11,231 2D reads with a mean length of 7.5 kb; the predicted genome coverage depth was ~16×. Fasta sequences were extracted from FAST5 files using poRe (7) and used along with the existing Illumina data (~50× read depth [6]) in a hybrid assembly using Unicycler (8). Assembly of the Illumina data alone using SPAdes (9) produced an assembly of 87 scaffolds, including a complete pX02 plasmid sequence but multicontig chromosomal and pX01 sequences (comprised of 82 and 4 contigs, respectively). The addition of the Nanopore data and use of the Unicycler assembly pipeline produced complete circular sequences for the chromosome (5,288,599 bp) and the pX01 and pX02 plasmids (180,666 bp and 94,967 bp, respectively).

Data availability. These data are available under NCBI BioProject number [PRJNA287512](#), BioSample number [SAMN03790807](#), and GenBank accession numbers [CP029805](#) (London_499 chromosome), [CP029806](#) (London_499 pX01), and [CP029807](#) (London_499 pX02).

ACKNOWLEDGMENTS

We thank all those involved in sample collection, Cath Arnold, and the Public Health England Genomic Services and Development Unit sequencing team.

Received 15 June 2018 **Accepted** 29 August 2018 **Published** 4 October 2018

Citation Pullan ST, Miles RW, Lewandowski K, Vipond R. 2018. Closed genome sequence obtained using hybrid Nanopore/Illumina assembly of a *Bacillus anthracis* isolate from an animal-skin-drum-associated anthrax case in the United Kingdom. Microbiol Resour Announc 7:e00802-18. <https://doi.org/10.1128/MRA.00802-18>.

Editor Catherine Putonti, Loyola University Chicago

© Crown copyright 2018. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#).

Address correspondence to Steven T. Pullan, steven.pullan@phe.gov.uk.

The views expressed here are those of the authors and not necessarily those of Public Health England or the Department of Health.

REFERENCES

1. Sahl JW, Pearson T, Okinaka R, Schupp JM, Gillette JD, Heaton H, Birdsell D, Hepp C, Fofanov V, Noseda R, Fasanella A, Hoffmaster A, Wagner DM, Keim P. 2016. A *Bacillus anthracis* genome sequence from the Sverdlovsk 1979 autopsy specimens. *mBio* 7:e01501-16. <https://doi.org/10.1128/mBio.01501-16>.
2. Keim P, Grunow R, Vipond R, Grass G, Hoffmaster A, Birdsell DN, Klee SR, Pullan S, Antwerpen M, Bayer BN, Latham J, Wiggins K, Hepp C, Pearson T, Brooks T, Sahl J, Wagner DM. 2015. Whole genome analysis of injectional anthrax identifies two disease clusters spanning more than 13 years. *EBioMedicine* 2:1613–1618. <https://doi.org/10.1016/j.ebiom.2015.10.004>.
3. Price EP, Seymour ML, Sarovich DS, Latham J, Wolken SR, Mason J, Vincent G, Drees KP, Beckstrom-Sternberg SM, Phillippy AM, Koren S, Okinaka RT, Chung W-K, Schupp JM, Wagner DM, Vipond R, Foster JT, Bergman NH, Burans J, Pearson T, Brooks T, Keim P. 2012. Molecular epidemiologic investigation of an anthrax outbreak among heroin users, Europe. *Emerg Infect Dis* 18:1307–1313. <https://doi.org/10.3201/eid1808.111343>.
4. Anaraki S, Addiman S, Nixon G, Krahé D, Ghosh R, Brooks T, Lloyd G, Spencer R, Walsh A, McCloskey B, Lightfoot N. 2008. Investigations and control measures following a case of inhalation anthrax in east London in a drum maker and drummer, October 2008. *Euro Surveill* 13:pii=19076. <https://www.eurosurveillance.org/content/10.2807/es.13.51.19076-en>.
5. Nguyen TQ, Clark N, 2006 NYC Anthrax Working Group, Karpati A, Goldberg A, Paykin A, Tucker A, Baker A, Almiroudis A, Fine A, Tsoi B, Aston C, Berg D, Weiss D, Connelly E, Beaudry G, Weisfuse I, Durrah JC, Prudhomme J, Leighton J, Ackelsberg J, Mahoney K, Van Vynck L, Lee L, Moskin L, Layton M, Wong M, Raphael M, Robinson M, Phillips M, Jones M, Jeffery N, Nieves R, Slavinski S, Mullin S, Beatrice ST, Balter S, Blank S, Frieden T, Keifer M, Rosenstein N, Diaz P, Clark T, Compton H, Daloia J, Cardarelli J, Norrell N, Horn E, Jackling S, Bacon C, Glasgow E, Gomez T, Baltzersen RA, Kammerdener C, Margo-Zavazky D, Colgan J, Pulaski P. 2010. Public health and environmental response to the first case of naturally acquired inhalational anthrax in the United States in 30 years: infection of a New York City resident who worked with dried animal hides. *J Public Health Manag Pract* 16:189–200. <https://doi.org/10.1097/PHH.0b013e3181ca64f2>.
6. Pullan ST, Sahl JW, Vipond R, Atkinson B, Pearson TR, Hoffmaster AR, Latham J, Birdsell D, Silman NJ, Keim P, Mason J, Marston CK. 2015. Whole-genome sequencing investigation of animal-skin-drum-associated UK anthrax cases reveals evidence of mixed populations and relatedness to a US case. *Microb Genom* 1:e000039. <https://doi.org/10.1099/mgen.0.000039>.
7. Watson M, Thomson M, Risse J, Talbot R, Santoyo-Lopez J, Gharbi K, Blaxter M. 2015. poRe: an R package for the visualization and analysis of nanopore sequencing data. *Bioinformatics* 31:114–115. <https://doi.org/10.1093/bioinformatics/btu590>.
8. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
9. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.