Tidal heating as a direct probe of strangeness inside neutron stars

Suprovo Ghosh,* Bikram Keshari Pradhan,[†] and Debarati Chatterjee[‡]

Inter-University Centre for Astronomy and Astrophysics,

Pune University Campus, Pune 411007, India

It has been discussed whether viscous processes in neutron star matter during a binary inspiral can damp out the tidal energy induced by the companion and heat up the star. Earlier investigations concluded that this tidal heating is negligible for normal neutron star matter. In this work, we suggest a novel effect of tidal heating involving strange matter in the neutron star interior, that can significantly heat up the star, and is potentially observable by current and future gravitational wave detectors. We propose that this could serve as a direct probe of strangeness in neutron stars.

I. INTRODUCTION

Neutron stars (NS) are unique astrophysical dense objects to study cold and dense matter under extreme conditions. In the high density environment of the NS interior, strangeness containing matter e.g. hyperons, kaons or deconfined quarks, can become stable components and have important consequences for the equation of state (EOS), structure and evolution of these compact objects [1–3, 76, 77]. Various astrophysical observations e.g., the measurement of high mass NS [4, 5], radius measurement from low-mass X-ray binaries [6, 7], thermal evolution [8] and gravitational wave (GW) emission from hot and rapidly rotating newly born neutron stars due to unstable r-modes [9–11] as well as binary neutron star merger event GW170817 [13–15] by the LIGO-Virgo detectors [16–18] have been used to constrain the hyperon content inside their core [12, 70, 71]. The measurement of tidal deformability from GW170817 [13] along with recent NICER estimates of mass-radius posteriors from two pulsars, PSR J0030 + 0451 [20, 21] and PSR J0740 + 6620 [22, 23] have significantly constrained the unknown neutron star EOS [24–29].

During binary neutron star inspiral, tidal interaction due to the companion transfers mechanical energy and angular momentum to the star at the expense of the orbital energy and the damping mechanisms inside the star convert this energy to heat ("tidal heating"). This tidal heating could potentially cause mass ejection in a radiation-driven outflow prior to the merger [30] or spin the stars up to corotation before merger [31, 32]. However this heating is found to be negligible for viscosities arising from nucleonic or leptonic weak processes, because of the long viscous timescale. Lai (1994) [33] found that shear and bulk viscous damping of mode oscillations of normal NS matter could only heat the stellar core to $\sim 3 \times 10^7 {\rm K}$. Recent work by Arras et al. (2019) [34] considers heating due to direct Urca

reactions during inspiral and concluded that they are heated upto $T \sim 10^8 \mathrm{K}$, which is one order of magnitude higher than the estimates by Lai (1994) [33].

Bulk viscosity originating from non-leptonic weak interaction involving strange matter in the neutron star core have been studied extensively in the context of damping of unstable r-mode oscillations [9–11, 35–40]. Hyperon bulk viscosity is found to be several orders higher than viscosity due to nucleonic Urca ($\approx 10^8 - 10^{10}$ times the canonical shear viscosity from e - e scattering in the temperature range $10^6 - 10^8 \text{K}$) and strongly suppress the r-mode instability below 10¹⁰K [9, 38]. In this article, for the first time we propose that this tidal heating due to bulk viscosity from strange matter in the neutron star core can sufficiently heat up the star during the inspiral to temperatures much higher than estimated from earlier studies. This effect is potentially detectable by current and future generation GW detectors as a deviation of the orbital decay rate from the general relativistic point-mass result.

The structure of the article is as follows: in Sec. II, we compute the hyperon bulk viscosity for several choice of EOS models and compare the strength with canonical shear and bulk viscosity sources inside neutron star cores. In Sec. III, we describe the formalism to calculate tidal heating in binary neutron star inspiral and estimate the temperatures reached during inspiral due to dissipation from hyperon bulk viscosity. Then in Sec.IV, we estimate the phase difference introduced in the gravitational wave signal from binary NS inspiral due to this viscous dissipation and argue that it will be detectable in future generation GW detectors. In Sec. V we discuss the main implications of this work and also future directions.

II. HYPERONIC BULK VISCOSITY

In this work, we explore the effect by calculating the bulk viscosity from hyperons within a simple model. During binary inspiral, bulk viscosity arises due to the pressure and density variations associated with the dynamical tidal excitation of NS oscillation modes (e.g. f or

^{*} suprovo@iucaa.in

[†] bikramp@iucaa.in

[‡] debarati@iucaa.in

g-modes) that drive the system out of chemical equilibrium [35]. Weak reactions among the core constituents, having timescales comparable to the oscillation modes, bring the system back to equilibrium. Among the possible hyperons that may appear in the core, Λ hyperons contribute dominantly to the bulk viscosity [9, 35]. Nuclear experiments indicate a repulsive Σ -nucleon potential [45–47] that result in suppression of Σ -hyperons, while the Ξ-hyperons may appear at large densities [4] beyond the maximum densities reached even in very massive NSs. However, one must note that for certain parametrizations e.g. FSU2H [72] and density dependent RMF parametrizations [73, 74], the onset densities of Λ , Σ and Ξ hyperons are quite close. As a proof-of-principle, in this study we consider the dominant non-leptonic weak process involving only the Λ hyperon

$$n + p \longleftrightarrow p + \Lambda.$$
 (1)

Though there may be other processes involving the Λ hyperon and also other hyperon species (Σ and Ξ hyperons) in NS core [44], this estimate provides a reasonable lower limit on the rates and hence an upper limit on the hyperon bulk viscosity [35, 44]. To investigate the effect of the EOS, we consider a diverse set of parametrizations within the Relativistic Mean Field (RMF) model [41–43] consistent with recent multi-messenger observations. The potential depth of Λ hyperons in normal nuclear matter is set to $U_{\Lambda}^{\dot{N}}=-30$ MeV [5] for all the parametrizations. In Fig. 1, we plot the particle fraction of all the constituents $(npe\mu \& \Lambda)$ for chosen set of RMF parametrizations and also mention the density for appearance of Λ hyperon in Table I. We see that for all the parametrizations, Λ -hyperon appear first around twice the nuclear saturation density ($\sim 2n_0$) and their fraction increase with increasing density. For the chosen EOSs, we also plot the mass-radius in Fig. 2 along with the astrophysical constraints from binary NS merger event GW170817 [13] and NICER estimates of PSR J0030+0451 [20, 21] and PSR J0740+6620 [22, 23]. The maximum mass reached by all the EOSs are $\geq 2M_{\odot}$ as also mentioned in Table I and consistent with the observed massive pulsar PSR J0740 + 6620 [78]. our study, to consider significant hyperon fraction we choose a couple of stiff EOSs (e.g. NL3, TM1) for which radius just extends($\sim 1 \text{ km}$) beyond the 90% confidence interval of GW170817 estimates but they are still well within the limits of NICER estimates of PSR J0030 + 0451.

Now, we calculate the hyperon bulk viscosity analytically for these chosen EOSs, whose details are provided in Table I along with hyperon potentials and onset densities. Given an EOS, the real part of the bulk viscosity coefficient (ζ) can be calculated in terms of relaxation times (τ) of weak processes [9, 10]

$$\zeta = n_B \frac{\partial P}{\partial n_n} \frac{dn_n}{dn_B} \frac{\tau}{1 + (\omega \tau)^2},\tag{2}$$




FIG. 1. Particle fraction as a function of normalised density(n_0 is the nuclear saturation density) for the chosen EOS parametrizations given in Table I. The compositions considered here are : $n(\text{dashed lines}), p(\text{dashed dotted line}), e(\text{dotted lines}), \mu(\text{dense dotted lines})$ and Λ -hyperon(solid lines).




FIG. 2. Mass radius diagram for the different EOSs given in Table I along with the Mass-Radius posterior from GW170817 [13] and $2-\sigma$ NICER posterior of PSR J0030+0451 [20, 21] and PSR J0740+6620 [22, 23]. The $1-\sigma$ mass estimate of massive pulsar PSR J0740+6620 [78] is shown in light green band with the median value of $2.08M_{\odot}$ shown in the deep blue line.

where P is the pressure, n_B the total baryon number density, n_n the neutron density and ω the frequency of the oscillation mode. The rates $1/\tau$ of these reactions are calculated from tree-level Feynman diagrams involving exchange of a W-boson [9, 35, 39]. In Fig. 3, we display the bulk viscosity due to the process 1 as a function of temperature for the equations of state in Table I, for typical characteristic values of mode frequency: 1 &




FIG. 3. Relative strengths of various sources of viscosities inside NS matter as a function of temperature for $n=2.5n_0$ ($n_0=$ the nuclear saturation density). The bands corresponds to hyperon bulk viscosity coming from different EOSs given in Table I with the black solid line corresponding to HZTCS EOS. The standard neutron matter shear viscosity coming from e-e scattering and bulk viscosity from m-Urca reactions at $f=1 \rm kHz$ are also plotted along with their temperature dependence.

2 kHz for f-mode [48] and ~ 100 Hz for g-mode oscillations [49]. Depending on the mode oscillation frequency, we see that the resonances occurs in the temperature range of $10^8 - 10^9$ K and the resonant maximum value of ζ can reach values of $10^{31} - 10^{32}$ gm cm⁻¹s⁻¹, which are several orders of magnitude higher than the canonical shear and bulk viscosity [50] in this temperature regime.

III. TIDAL HEATING IN BINARY NS

To investigate the effect of this hyperon viscous damping on heating of the NS, we first analyse the timescale for mode damping. The mode damping rate is given by

$$\gamma_{visc,\alpha} = \frac{1}{2E_{\alpha}} \dot{E}_{visc,\alpha} \tag{3}$$

where α denotes the particular eigenmode, E_{α} is the energy of the mode and $\dot{E}_{visc,\alpha}$ is the energy dissipation rate. In the adiabatic approximation, the effect of the tidal potential due to the companion star is measured in terms of the Lagrangian fluid displacement vector $\boldsymbol{\xi}(r,t)$ from its equilibrium position. This displacement can be analysed in terms of the normal modes of the NS. In this work, we only consider the dominant l=m=2 fundamental (f) mode contribution to the tidal heating. We determine the f-mode frequency via the Cowling approximation [48] and the normalised mode eigenfunctions are used to calculate the tidal coupling [55]. Although general relativistic effects can modify f-mode frequencies by

up to 30% [69], we restrict ourselves to Cowling approximation to be consistent with the Newtonian description of tidal heating used in this work. The bulk viscous dissipation rate is expressed as [33]

$$\gamma_{bulk} = 12 \int_0^R r^2 dr \zeta \left(\frac{\partial \xi^r}{\partial r} + \frac{2}{r} \xi^r - l(l+1) \frac{\xi^\perp}{r} \right)^2. \quad (4)$$

where ζ is the bulk viscous coefficient, R is the radius of the star, ξ^r and ξ^{\perp} are radial and tangential component of the displacement vector respectively.

We compare this viscous dissipation timescale given by inverse of the rate $\tau_{bulk} = 1/\gamma_{bulk}$, with the inspiral timescale to confirm whether the hyperon bulk viscous dissipation can effectively drain energy from the decaying orbit during the inspiral. We consider leading order gravitational radiation where the GW emission takes energy from the orbit at a rate of

$$\dot{E}_{gw} = \frac{-32\mathcal{M}\Omega}{5c^5} (G\mathcal{M}\Omega)^{7/3} . \tag{5}$$

Here the chirp mass is given by $\mathcal{M}=M\left(\frac{q^3}{1+q}\right)^{1/5}$ with the primary star mass is M and the companion mass is qM. Ω denotes the orbital frequency which is given by $\Omega^2=\frac{GM(1+q)}{D^3}$ where D is the separation between the masses. In this evolution dynamics, the orbital timescale is given as [55]

$$\frac{1}{t_D} \equiv \frac{\dot{\Omega}}{\Omega} = \frac{96}{5c^5} (GM\Omega)^{5/3} \Omega \ . \tag{6}$$

In Fig. 4, we plot this inspiral timescale for the LIGO-Virgo-KAGRA (LVK) frequency band of 20-500 Hz [16–19] with the viscous dissipation and tidal heating timescale for the case of equal binary mass of $1.8 \rm M_{\odot}$ for all the EOSs given in Table. I. We see that both the hyperon bulk viscous dissipation and tidal heating timescales are smaller than the inspiral timescale, confirming that unlike shear viscous dissipation, hyperon bulk viscous dissipation and heating happens faster than the orbital evolution and can efficiently damp out the tidal energy to heat up the star during the inspiral.

Since we only consider the dominant mode l=m=2 f-mode contribution, the energy dissipation rate can be estimated from the mode amplitude [33]

$$\dot{E}_{visc} = \frac{12\pi}{5} \frac{GM^2}{R} q^2 (1+q) \omega_0^{-4} Q_0^2 \left(\frac{R}{D}\right)^9 2\gamma_{bulk}$$
 (7)

where Q_0 tidal coupling strength of the f-mode and ω_0 the normalised frequency of the f-mode. The heat content of the star due to the degenerate fermionic gas in the core can be given as $U \approx 4.5 \times 10^{22} T^2 J$ [33]. During the inspiral, the timescales for cooling due to neutrino emission and surface photon luminosity are very

EOS	Max. mass	A onset	Mass (M_{\odot})	Central	Radius	Hyperon core	f-mode	Max. temp (K)	$\Delta \Phi = 2\pi \Delta N$
	(M_{\odot})	density	111000 (1110)	density	(km)	radius(km)	Freq.(Hz)	at $D/R = 3$	(rad)
	()			1 22 2 2 2 2 3	()	()	1.()		(====)
NL3	2.10	$1.90n_0$	1.6	$2.07n_0$	14.7	3.42	1847	9.7×10^{8}	0.001
[42]			1.8	$2.48n_0$	14.6	6.19	1909	3.3×10^{9}	0.08
			2.0	$3.35n_0$	14.2	8.10	2009	6.2×10^9	0.3
TM1	2.06	$2.02n_0$	1.6	$2.24n_0$	14.55	3.16	1873	8.7×10^{8}	0.0008
[51]			1.8	$2.77n_0$	14.37	6.18	1947	3.4×10^9	0.09
			2.0	$4.06n_0$	13.6	8.15	2092	6.7×10^9	0.34
TMA	2.12	$2.09n_0$	1.8	$2.54n_0$	14.2	5.13	1948	2.3×10^{9}	0.02
[51]			2.0	$3.35n_0$	13.89	7.36	1909	5.1×10^9	0.16
HZTCS	2.00	$2.28n_0$	1.6	$2.67n_0$	13.2	4.89	2108	2.3×10^{9}	0.02
[52]			1.8	$3.32n_0$	13.1	6.82	2171	4.7×10^9	0.16
			2.0	$5.32n_0$	12.25	8.17	2305	7.9×10^9	0.44
FSU2	2.03	$1.92n_0$	1.6	$2.22n_0$	14.4	4.98	1898	2.1×10^9	0.03
[42]			1.8	$2.72n_0$	14.2	7.07	1968	4.5×10^9	0.19
			2.0	$3.82n_0$	13.6	8.54	2099	7.4×10^9	0.47
Stiffest EOS	2.01	$2.31n_0$	1.6	$2.71n_0$	13.5	3.88	2047	1.4×10^9	0.004
from Ghosh et al.			1.8	$3.39n_0$	13.4	6.34	2119	4.1×10^9	0.11
(2022) [53]			2.0	$5.5n_0$	12.5	8.05	2256	7.2×10^9	0.37

TABLE I. Detailed list of NS properties corresponding to different EOS parametrizations considered in this work. The potential depth of Λ hyperons in normal nuclear matter is set to $U_{\Lambda}^{N}=-30~{\rm MeV}$ [5] for all the parametrizations. Densities are given in terms of n_0 , the nuclear saturation density ($\sim 0.15~{\rm fm}^{-3}$ [54]). The phase difference $\Delta\Phi$ in the last column denotes the total phase difference accumulated at the end of inspiral around $f\sim 500{\rm Hz}$.




FIG. 4. Estimated timescale for the different processes as a function of GW frequency compared against the inspiral timescale (t_D) for a NS binary of equal mass $1.8M_{\odot}$. Shear viscous and hyperon bulk viscosity dissipation corresponds to the dominant f-mode dissipation by shear viscosity from e-e scattering [33] and bulk viscosity from hyperons respectively. The tidal heating corresponds to the heating timescale from the hyperon bulk viscous dissipation. The different bands indicates uncertainties due to the choice of different EOSs given in Table I.

high compared to the binary inspiral timescale as shown in Fig. 4, and therefore negligible [30]. After integrating the thermal evolution equation $\frac{dU}{dt}=\dot{E}_{visc}+\dot{E}_{cool}$ from $D\to\infty$ when the stars were far apart and at a very low

temperature $(10^5 - 10^6 \text{K})$, we can get an estimate of the temperature reached as a function of their separation D

$$\begin{split} \left[\frac{T^4}{4} + B ln(T)\right] &= \frac{\pi}{21870} \omega_0^{-4} Q_0^2 q \frac{A}{10^{22}} \\ &\times \left(\frac{c^2 R}{GM}\right) \left(\frac{c^3 R^2}{G}\right) \left(\frac{3R}{D}\right)^5, \end{split} \tag{8}$$

where A and B are parameters fitted to the functional dependence of γ_{bulk} on the temperature (T), $\gamma_{bulk} = \frac{AT^2}{B+T^4}$ coming from the temperature dependence of timescale for hyperon bulk viscosity. In Table I, we provide the estimates of the temperature reached at a separation of D=3R when the stars are about to merge. We find temperatures $\sim 10^9-10^{10}{\rm K}$, which are two orders of magnitude higher than the earlier estimates [33, 34].

IV. PHASE ERROR ESTIMATION AND DETECTABILITY

The energy loss due to tidal heating during the binary inspiral will lead to a change in the number of wave cycles $(\Delta \mathcal{N})$ or equivalently, a phase shift $\Delta \Phi = 2\pi \Delta \mathcal{N}$ in the observed frequency range of the GW detectors. This is crucial to accurately guess the phase of the signal, otherwise the GW template can destroy a possible detection using matched filter technique [56]. This additional torque to the viscous dissipation of energy will

lead to a total change in the number of cycles given by

$$\Delta \mathcal{N} = \frac{15\pi}{384} \left(\frac{c^2 R}{GM} \right)^6 \omega_0^{-4} Q_0^2 \frac{1}{q(1+q)} \left(\frac{R}{c} \right)^2 \int_{f_a}^{f_b} \gamma_{bulk} df.$$
(9)

where f_a is the frequency when the signal first enters the detector band, and f_b , when it dives into the noise again. Since, f-modes are not resonantly excited, we need to do the integration over the whole frequency range unlike the cases for resonantly excited g-modes or r-modes where additional energy loss is associated with the particular mode frequency [55, 57]. In Table I, we demonstrate the net phase difference accumulated at GW frequency of 500 Hz for all the different EOSs and different values of equal mass binaries. In Fig. 5, we display how this phase difference grows as a function of GW frequency, taking into consideration uncertainties due to the EOSs. For equal $1.6M_{\odot}$ binaries, we see that the net phase difference is of the order of $10^{-3} - 10^{-2}$ rad and for higher masses of $1.8M_{\odot}$ or $2M_{\odot}$, we get a net phase difference in the order of 0.1 - 0.5 rad.

To be able to measure this phase difference using the current or future generation GW detectors, the phase uncertainty of detected GWs must be smaller than this extra phase shift. Recent analysis [58] have shown the phase error to be around $\Delta \phi \sim \pm 0.1$ rad for $f_{GW} \leq 300$ Hz inclusive of calibration uncertainties for the GW170817 signal analysed using GW waveform model IMRPhenomPv2_NRTidal [59]. More recently, Read (2023) [60] compares a number of GW waveform models and shows that the uncertainty due to waveform differences is \sim ± 0.02 rad for A+ [61] and $\pm 10^{-3}$ rad for Cosmic Explorer (CE) [62]. So, from these estimates, we see that a binary neutron star event with signal-to-noise ratio(SNR) like GW170817 would produce enough tidal heating to be detectable using the current LVK detectors if it has a heavier component mass $\geq 1.8 M_{\odot}$. In the 3G detectors, we can measure evidence of tidal heating due to hyperons even for much lower mass NS components.

V. DISCUSSION

In this work, we propose a novel effect of tidal heating in NSs during binary inspiral due to bulk viscosity originating from non-leptonic weak interaction processes involving hyperons in the NS core. We demonstrate the effect by calculating the bulk viscous dissipation of the dominant f-mode oscillations excited due to the tidal interaction during the inspiral. We consider several state-of-the-art EOSs including the hyperonic degrees of freedom consistent with multi-messenger observations. This dissipated energy can be effectively converted to thermal energy during the inspiral timescale and can heat up the stars upto 0.1-1 MeV during the last orbits before coalescing. Although these estimates are higher by orders of magnitudes than earlier estimates by




FIG. 5. Estimated phase shift accumulated in the GW signal for equal mass binaries as a function of GW frequency. The different bands indicate uncertainties due to the choice of different EOSs given in Table I corresponding to different masses of equal mass binaries considered and the solid lines define their boundaries.

Lai (1994) [33] and Arras et al. (2019) [34], they are not sufficiently high to demand inclusion of thermal effects which are very relevant for post-merger studies [75], in the EOS during inspiral [63]. The induced phase shift in the GW signal due to this tidal heating is ~ 0.1 rad for component masses $\geq 1.8 M_{\odot}$ which is potentially detectable by current and future generation GW detectors.

This work suggests for the first time that tidal heating can be used as a probe of strange matter inside NS core, and opens up the possibility of exploring the outof-equilibrium effects of NS matter on the GW signals. These findings pave the way for more sophisticated analyses as followup of this work, including a consistent general relativistic formulation of tidal heating and f-mode calculations, other EOS models including the contribution from other species of hyperons, other forms of strange matter (e.g. quarks) in order to get a more quantitatively accurate estimate of the observable effects. There are various other directions in which our work can be further developed. First, a detailed investigation of the Post-Newtonian(PN) order at which the tidal heating becomes relevant will have to be performed to confirm the conclusions related to the qualitative Newtonian inspiral discussed in this work. resonant g-mode contribution to tidal heating can also be as significant [33] since the hyperon bulk visocity value is one order of magnitude order higher than those of f-modes (see Fig. 3). Third, bulk viscosity from the dominant channel in unpaired strange quark matter is $\sim 10^{29} - 10^{30} \text{ gm cm}^{-1} \text{ s}^{-1} \text{ around } 10^8 - 10^9 \text{K } [67, 68]$ and could also contribute to the tidal heating. Finally,

the effect of superfluidity on the hyperon bulk viscosity should be investigated since they are known to reduce the rates of the weak interactions and also effective below the critical temperature of 10⁹K [38]. Several of these followup studies are currently in progress and will be reported in forthcoming publications.

We are currently developing a detailed Bayesian analysis using post-Newtonian waveform including the effect of viscous dissipation to accurately determine the bulk viscosity as well as estimate any possible degeneracy with other intrinsic parameters of binary NS inspiral. Considering the recent order-of-magnitude post-Newtonian estimate [66] of the direct effect of bulk viscosity on the orbital motion at f = 100Hz, when the bulk viscosity is $\sim 10^{32} \text{ gm cm}^{-1} \text{ s}^{-1} \text{(from Fig. 3), one}$ would require SNR of 100 to get a 10% measurement on the bulk viscosity, which is achievable using the third generation GW detectors. Recent observations of binary neutron star merger GW190425 [64] or neutron starblack hole merger GW200105 [65] having component NS masses $m_1^{90\%} = [1.6, 1.87] M_{\odot}$ and $m_2 = 1.9^{+0.3}_{-0.2}$ respectively, suggest that there might be other formation

channels for higher mass NS in binary systems which can have sufficient hyperons in their core to contribute significantly to the tidal heating that can be detectable. A future detection of this high bulk viscous effect originating from non-leptonic weak interaction processes involving hyperons either in terms of deviation in the orbital motion or tidal heating would strongly indicate the presence of strange hyperons in the NS interior, having a great significance in nuclear astrophysics. Even a non-detection of this effect can place an upper limit on the bulk viscosity and the hyperon fraction in the NS core, with important implications for understanding of dense nuclear matter.

ACKNOWLEDGMENTS

The authors thank Alexander Haber for his useful comments regarding the paper. The authors would like to acknowledge the usage of the high-performance supercomputer Pegasus at Inter-University Centre for Astronomy and Astrophysics for the numerical calculations.

- [1] N. Glendenning, Compact Stars: Nuclear Physics, Particle Physics and General Relativity, Astronomy and Astrophysics Library (Springer New York, 2012).
- [2] J. M. Lattimer and M. Prakash, The physics of neutron stars, Science **304**, 536 (2004).
- [3] D. Blaschke and N. Chamel, Phases of dense matter in compact stars, in *The Physics and Astrophysics of Neu*tron Stars, edited by L. Rezzolla, P. Pizzochero, D. I. Jones, N. Rea, and I. Vidaña (Springer International Publishing, Cham, 2018) pp. 337–400.
- [4] S. Weissenborn, D. Chatterjee, and J. Schaffner-Bielich, Hyperons and massive neutron stars: Vector repulsion and su(3) symmetry, Phys. Rev. C 85, 065802 (2012).
- [5] S. Weissenborn, D. Chatterjee, and J. Schaffner-Bielich, Hyperons and massive neutron stars: The role of hyperon potentials, Nuclear Physics A 881, 62 (2012).
- [6] C. m. c. Providência and A. Rabhi, Interplay between the symmetry energy and the strangeness content of neutron stars, Phys. Rev. C 87, 055801 (2013).
- [7] Fortin, M., Zdunik, J. L., Haensel, P., and Bejger, M., Neutron stars with hyperon cores: stellar radii and equation of state near nuclear density, A&A 576, A68 (2015).
- [8] F. Anzuini, A. Melatos, C. Dehman, D. Viganò, and J. A. Pons, Fast cooling and internal heating in hyperon stars, Monthly Notices of the Royal Astronomical Society 10.1093/mnras/stab3126 (2021).
- [9] L. Lindblom and B. J. Owen, Effect of hyperon bulk viscosity on neutron-star r-modes, Phys. Rev. D 65, 063006 (2002).
- [10] D. Chatterjee and D. Bandyopadhyay, Effect of hyperonhyperon interaction on bulk viscosity and r-mode instability in neutron stars, Phys. Rev. D 74, 023003 (2006).
- [11] O. P. Jyothilakshmi, P. E. S. Krishnan, P. Thakur, V. Sreekanth, and T. K. Jha, Hyperon bulk vis-

- cosity and r-modes of neutron stars, Monthly Notices of the Royal Astronomical Society **516**, 3381 (2022), https://academic.oup.com/mnras/article-pdf/516/3/3381/45882806/stac2360.pdf.
- [12] D. Chatterjee and I. Vidaña, Do hyperons exist in the interior of neutron stars?, The European Physical Journal A 52, 10.1140/epja/i2016-16029-x (2016).
- [13] B. Abbott, R. Abbott, T. Abbott, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. Adhikari, V. Adya, and et al., Gw170817: Observation of gravitational waves from a binary neutron star inspiral, Physical Review Letters 119, 10.1103/physrevlett.119.161101 (2017).
- [14] B. Abbott, R. Abbott, T. Abbott, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. Adhikari, V. Adya, and et al., Gw170817: Measurements of neutron star radii and equation of state, Physical Review Letters 121, 10.1103/physrevlett.121.161101 (2018).
- [15] B. Abbott, R. Abbott, T. Abbott, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. Adhikari, V. Adya, and et al. (LIGO Scientific Collaboration and Virgo Collaboration), Properties of the binary neutron star merger gw170817, Phys. Rev. X 9, 011001 (2019).
- [16] J. Aasi, B. P. Abbott, R. Abbott, T. Abbott, M. R. Abernathy, K. Ackley, C. Adams, T. Adams, P. Addesso, and et al., Advanced ligo, Classical and Quantum Gravity 32, 074001 (2015).
- [17] B. P. Abbott, R. Abbott, T. D. Abbott, M. R. Abernathy, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, and R. X. e. a. Adhikari (LIGO Scientific Collaboration and Virgo Collaboration), Gw150914: The advanced ligo detectors in the era of first discoveries, Phys. Rev. Lett. 116, 131103 (2016).
- [18] F. Acernese, M. Agathos, K. Agatsuma, D. Aisa, N. Alle-

- mandou, A. Allocca, J. Amarni, P. Astone, G. Balestri, G. Ballardin, and et al., Advanced virgo: a second-generation interferometric gravitational wave detector, Classical and Quantum Gravity **32**, 024001 (2014).
- [19] T. Akutsu, M. Ando, K. Arai, Y. Arai, S. Araki, A. Araya, N. Aritomi, Y. Aso, S. W. Bae, and Y. B. B. et al., Overview of KAGRA: Detector design and construction history, Progress of Theoretical and Experimental Physics 2021, 05A101 (2021), arXiv:2005.05574 [physics.ins-det].
- [20] M. C. Miller, F. K. Lamb, A. J. Dittmann, S. Bog-danov, Z. Arzoumanian, K. C. Gendreau, S. Guillot, A. K. Harding, W. C. G. Ho, J. M. Lattimer, and et al., Psr j0030+0451 mass and radius from nicer data and implications for the properties of neutron star matter, The Astrophysical Journal 887, L24 (2019).
- [21] T. E. Riley, A. L. Watts, S. Bogdanov, P. S. Ray, R. M. Ludlam, S. Guillot, Z. Arzoumanian, C. L. Baker, A. V. Bilous, D. Chakrabarty, and et al., A nicer view of psr j0030+0451: Millisecond pulsar parameter estimation, The Astrophysical Journal 887, L21 (2019).
- [22] M. C. Miller, F. K. Lamb, A. J. Dittmann, S. Bogdanov, Z. Arzoumanian, K. C. Gendreau, S. Guillot, W. C. G. Ho, J. M. Lattimer, M. Loewenstein, and et al., The radius of psr j0740+6620 from nicer and xmm-newton data, The Astrophysical Journal Letters 918, L28 (2021).
- [23] T. E. Riley, A. L. Watts, P. S. Ray, S. Bogdanov, S. Guillot, S. M. Morsink, A. V. Bilous, Z. Arzoumanian, D. Choudhury, J. S. Deneva, and et al., A nicer view of the massive pulsar psr j0740+6620 informed by radio timing and xmm-newton spectroscopy, The Astrophysical Journal Letters 918, L27 (2021).
- [24] E. Annala, T. Gorda, A. Kurkela, and A. Vuorinen, Gravitational-wave constraints on the neutron-star-matter equation of state, Phys. Rev. Lett. 120, 172703 (2018).
- [25] S. De, D. Finstad, J. M. Lattimer, D. A. Brown, E. Berger, and C. M. Biwer, Tidal deformabilities and radii of neutron stars from the observation of gw170817, Phys. Rev. Lett. 121, 091102 (2018).
- [26] E. R. Most, L. R. Weih, L. Rezzolla, and J. Schaffner-Bielich, New constraints on radii and tidal deformabilities of neutron stars from gw170817, Physical Review Letters 120, 10.1103/physrevlett.120.261103 (2018).
- [27] T. Dietrich, M. W. Coughlin, P. T. H. Pang, M. Bulla, J. Heinzel, L. Issa, I. Tews, and S. Antier, Multimessenger constraints on the neutron-star equation of state and the hubble constant, Science 370, 1450–1453 (2020).
- [28] P. T. H. Pang, I. Tews, M. W. Coughlin, M. Bulla, C. V. D. Broeck, and T. Dietrich, Nuclear physics multimessenger astrophysics constraints on the neutron star equation of state: Adding nicer's psr j0740 + 6620 measurement, The Astrophysical Journal 922, 14 (2021).
- [29] I. Legred, K. Chatziioannou, R. Essick, S. Han, and P. Landry, Impact of the psr J0740 + 6620 radius constraint on the properties of high-density matter, Phys. Rev. D 104, 063003 (2021).
- [30] P. Meszaros and M. J. Rees, Tidal Heating and Mass Loss in Neutron Star Binaries: Implications for Gamma-Ray Burst Models, The Astrophysical Journal 397, 570 (1992).
- [31] C. S. Kochanek, Coalescing Binary Neutron Stars, The Astrophysical Journal 398, 234 (1992).
- [32] L. Bildsten and C. Cutler, Tidal Interactions of Inspiral-

- ing Compact Binaries, The Astrophysical Journal **400**, 175 (1992).
- [33] D. Lai, Resonant oscillations and tidal heating in coalescing binary neutron stars, Monthly Notices of the Royal Astronomical Society 270, 611 (1994).
- [34] P. Arras and N. N. Weinberg, Urca reactions during neutron star inspiral, Monthly Notices of the Royal Astronomical Society 486, 1424 (2019), https://academic.oup.com/mnras/articlepdf/486/1/1424/28449547/stz880.pdf.
- [35] P. B. Jones, Bulk viscosity of neutron-star matter, Phys. Rev. D 64, 084003 (2001).
- [36] E. N. E. van Dalen and A. E. L. Dieperink, Bulk viscosity in neutron stars from hyperons, Phys. Rev. C 69, 025802 (2004).
- [37] T. K. Jha, H. Mishra, and V. Sreekanth, Bulk viscosity in a hyperonic star and r-mode instability, Phys. Rev. C 82, 025803 (2010).
- [38] B. Haskell and N. Andersson, Superfluid hyperon bulk viscosity and the r-mode instability of rotating neutron stars, Monthly Notices Royal Astronomical Society of the **408**. 1897 (2010),https://academic.oup.com/mnras/articlepdf/408/3/1897/18589963/mnras0408-1897.pdf.
- [39] D. D. Ofengeim, M. E. Gusakov, P. Haensel, and M. Fortin, Bulk viscosity in neutron stars with hyperon cores, Phys. Rev. D 100, 103017 (2019).
- [40] M. G. Alford and A. Haber, Strangeness-changing rates and hyperonic bulk viscosity in neutron star mergers, Phys. Rev. C 103, 045810 (2021).
- [41] N. K. Glendenning and S. A. Moszkowski, Reconciliation of neutron-star masses and binding of the λ in hypernuclei, Phys. Rev. Lett. **67**, 2414 (1991).
- [42] W.-C. Chen and J. Piekarewicz, Building relativistic mean field models for finite nuclei and neutron stars, Phys. Rev. C 90, 044305 (2014).
- [43] S. Ghosh, D. Chatterjee, and J. Schaffner-Bielich, Imposing multi-physics constraints at different densities on the neutron Star Equation of State, Eur. Phys. J. A 58, 37 (2022), arXiv:2107.09371 [astro-ph.HE].
- [44] M. E. Gusakov and E. M. Kantor, Bulk viscosity of superfluid hyperon stars, Physical Review D 78, 10.1103/physrevd.78.083006 (2008).
- [45] E. Friedman and A. Gal, In-medium nuclear interactions of low-energy hadrons, Physics Reports 452, 89–153 (2007).
- [46] J. Schaffner-Bielich and A. Gal, Properties of strange hadronic matter in bulk and in finite systems, Phys. Rev. C 62, 034311 (2000).
- [47] J. Mareš, E. Friedman, A. Gal, and B. Jenning, Constraints on σ -nucleus dynamics from dirac phenomenology of σ^- atoms, Nuclear Physics A **594**, 311 (1995).
- [48] B. K. Pradhan and D. Chatterjee, Effect of hyperons on f-mode oscillations in neutron stars, Physical Review C 103, 10.1103/physrevc.103.035810 (2021).
- [49] V. Tran, S. Ghosh, N. Lozano, D. Chatterjee, and P. Jaikumar, g-mode oscillations in neutron stars with hyperons (2022), arXiv:2212.09875 [nucl-th].
- [50] R. F. Sawyer, Bulk viscosity of hot neutron-star matter and the maximum rotation rates of neutron stars, Phys. Rev. D 39, 3804 (1989).
- [51] A. W. Steiner, M. Hempel, and T. Fischer, Core-collapse supernova equations of state based on neutron star observations, The Astrophysical Journal 774, 17 (2013).

- [52] N. Hornick, L. Tolos, A. Zacchi, J.-E. Christian, and J. Schaffner-Bielich, Relativistic parameterizations of neutron matter and implications for neutron stars, Phys. Rev. C 98, 065804 (2018).
- [53] S. Ghosh, B. K. Pradhan, D. Chatterjee, and J. Schaffner-Bielich, Multi-Physics Constraints at Different Densities to Probe Nuclear Symmetry Energy in Hyperonic Neutron Stars, Front. Astron. Space Sci. 9, 864294 (2022), arXiv:2203.03156 [astro-ph.HE].
- [54] M. Oertel, M. Hempel, T. Klähn, and S. Typel, Equations of state for supernovae and compact stars, Rev. Mod. Phys. 89, 015007 (2017).
- [55] N. Andersson and W. C. Ho, Using gravitational-wave data to constrain dynamical tides in neutron star binaries, Physical Review D 97, 10.1103/physrevd.97.023016 (2018).
- [56] L. Lindblom, B. J. Owen, and D. A. Brown, Model waveform accuracy standards for gravitational wave data analysis, Phys. Rev. D 78, 124020 (2008).
- [57] S. Ma, H. Yu, and Y. Chen, Detecting resonant tidal excitations of rossby modes in coalescing neutron-star binaries with third-generation gravitational-wave detectors, Phys. Rev. D 103, 063020 (2021).
- [58] B. Edelman, F. J. Rivera-Paleo, J. D. Merritt, B. Farr, Z. Doctor, J. Brink, W. M. Farr, J. Gair, J. S. Key, J. McIver, and A. B. Nielsen, Constraining unmodeled physics with compact binary mergers from gwtc-1, Phys. Rev. D 103, 042004 (2021).
- [59] T. Dietrich, S. Khan, R. Dudi, S. J. Kapadia, P. Kumar, A. Nagar, F. Ohme, F. Pannarale, A. Samajdar, S. Bernuzzi, G. Carullo, W. Del Pozzo, M. Haney, C. Markakis, M. Pürrer, G. Riemenschneider, Y. E. Setyawati, K. W. Tsang, and C. Van Den Broeck, Matter imprints in waveform models for neutron star binaries: Tidal and self-spin effects, Phys. Rev. D 99, 024029 (2019).
- [60] J. Read, Waveform uncertainty quantification and interpretation for gravitational-wave astronomy, Classical and Quantum Gravity 40, 135002 (2023).
- [61] R. Abbott, T. D. Abbott, S. Abraham, F. Acernese, K. Ackley, A. Adams, C. Adams, R. X. Adhikari, V. B. Adya, C. Affeldt, D. Agarwal, M. Agathos, K. Agatsuma, and et al., Prospects for observing and localizing gravitational-wave transients with advanced LIGO, advanced virgo and KAGRA, Living Reviews in Relativity 23, 10.1007/s41114-020-00026-9 (2020).
- [62] B. P. Abbott, R. Abbott, T. D. Abbott, M. R. Abernathy, K. Ackley, C. Adams, P. Addesso, R. X. Adhikari, V. B. Adya, C. Affeldt, N. Aggarwal, and et al., Exploring the sensitivity of next generation gravitational wave detectors, Classical and Quantum Gravity 34, 044001 (2017).
- [63] A. Kanakis-Pegios, P. Koliogiannis, and C. Moustakidis, Thermal effects on tidal deformability in the last orbits of an inspiraling binary neutron star system, Physics Letters B 832, 137267 (2022).
- [64] R. Abbott, T. D. Abbott, S. Abraham, F. Acernese, K. Ackley, A. Adams, C. Adams, R. X. Adhikari, V. B. Adya, C. Affeldt, D. Agarwal, M. Agathos, K. Agatsuma, and et al., The Astrophysical Journal Letters 892, L3 (2020).
- [65] R. Abbott, T. D. Abbott, S. Abraham, F. Acernese, K. Ackley, A. Adams, C. Adams, R. X. Adhikari, V. B. Adya, C. Affeldt, D. Agarwal, M. Agathos, K. Agatsuma,

- N. Aggarwal, and et al., Observation of gravitational waves from two neutron star-black hole coalescences, The Astrophysical Journal Letters **915**, L5 (2021).
- [66] E. R. Most, S. P. Harris, C. Plumberg, M. G. Alford, J. Noronha, J. Noronha-Hostler, F. Pretorius, H. Witek, and N. Yunes, Projecting the likely importance of weakinteraction-driven bulk viscosity in neutron star mergers, Monthly Notices of the Royal Astronomical Society 509, 1096 (2021).
- [67] M. G. Alford, S. Mahmoodifar, and K. Schwenzer, Large amplitude behavior of the bulk viscosity of dense matter, Journal of Physics G: Nuclear and Particle Physics 37, 125202 (2010).
- [68] M. G. Alford and K. Schwenzer, What the timing of millisecond pulsars can teach us about their interior, Phys. Rev. Lett. 113, 251102 (2014).
- [69] B. K. Pradhan, D. Chatterjee, M. Lanoye, and P. Jaikumar, General relativistic treatment of f-mode oscillations of hyperonic stars, Phys. Rev. C 106, 015805 (2022).
- [70] R. O. Gomes, P. Char, and S. Schramm, Constraining strangeness in dense matter with GW170817, The Astrophysical Journal 877, 139 (2019).
- [71] S. Blacker, H. Kochankovski, A. Bauswein, A. Ramos, and L. Tolos, Thermal behavior as indicator for hyperons in binary neutron star merger remnants (2023), arXiv:2307.03710 [astro-ph.HE].
- [72] L. Tolos, M. Centelles, and A. Ramos, Equation of state for nucleonic and hyperonic neutron stars with mass and radius constraints, The Astrophysical Journal 834, 3 (2016).
- [73] A. R. Raduta, M. Oertel, and A. Sedrakian, Protoneutron stars with heavy baryons and universal relations, Monthly Notices of the Royal Astronomical Society 499, 914 (2020), https://academic.oup.com/mnras/article-pdf/499/1/914/33873654/staa2491.pdf.
- [74] Z.-H. Tu and S.-G. Zhou, Effects of the ϕ meson on the properties of hyperon stars in the density-dependent relativistic mean field model, The Astrophysical Journal **925**, 16 (2022).
- [75] J. Fields, A. Prakash, M. Breschi, D. Radice, S. Bernuzzi, and A. da Silva Schneider, Thermal effects in binary neutron star mergers, The Astrophysical Journal Letters 952, L36 (2023).
- [76] E. Annala, T. Gorda, J. Hirvonen, O. Komoltsev, A. Kurkela, J. Nättilä, and A. Vuorinen, Strongly interacting matter exhibits deconfined behavior in massive neutron stars, Nature Communications 14, 10.1038/s41467-023-44051-y (2023).
- [77] L. Tolos and L. Fabbietti, Strangeness in nuclei and neutron stars, Progress in Particle and Nuclear Physics 112, 103770 (2020).
- [78] E. Fonseca, H. T. Cromartie, T. T. Pennucci, P. S. Ray, A. Y. Kirichenko, S. M. Ransom, P. B. Demorest, I. H. Stairs, Z. Arzoumanian, L. Guillemot, A. Parthasarathy, M. Kerr, I. Cognard, P. T. Baker, H. Blumer, P. R. Brook, M. DeCesar, T. Dolch, F. A. Dong, E. C. Ferrara, W. Fiore, N. Garver-Daniels, D. C. Good, R. Jennings, M. L. Jones, V. M. Kaspi, M. T. Lam, D. R. Lorimer, J. Luo, A. McEwen, J. W. McKee, M. A. McLaughlin, N. McMann, B. W. Meyers, A. Naidu, C. Ng, D. J. Nice, N. Pol, H. A. Radovan, B. Shapiro-Albert, C. M. Tan, S. P. Tendulkar, J. K. Swiggum, H. M. Wahl, and W. W. Zhu, Refined mass and geometric measurements of the high-mass psr j0740+6620, The Astrophysical Journal

Letters **915**, L12 (2021).