

The first law of binary black hole scattering

Riccardo Gonzo*

Higgs Centre for Theoretical Physics, School of Physics and Astronomy, University of Edinburgh, EH9 3FD, UK

Jack Lewis[†] and Adam Pound[‡]

*School of Mathematical Sciences and STAG Research Centre,
University of Southampton, Southampton SO17 1BJ, United Kingdom*

(Dated: August 27, 2025)

In the last decade, the first law of binary black hole mechanics played an important unifying role in the gravitational two-body problem. More recently, binary black hole scattering and the application of high-energy physics methods have provided a new avenue into this classical problem. In this Letter, we connect these two themes by extending the first law to the case of scattering orbits. We present derivations based on classical S-matrix, Hamiltonian, and pseudo-Hamiltonian methods, the last of which allows us to include dissipative effects for the first time. Finally, a “boundary to bound” map links this first law to the traditional bound-orbit version. Through this map a little-known observable for scatter orbits, the elapsed proper time, is mapped to the Detweiler redshift for bound orbits, which is an invariant building block in gravitational waveform models.

Introduction.—The discovery of gravitational waves (GWs) from compact binary systems opened a new chapter in astronomy. Given the enhanced sensitivity and expanded frequency range of future GW detectors, we expect a dramatic increase in the number and variety of detectable compact binary sources [1–5]. Increasingly accurate waveform models will be needed to detect and analyse these sources [6, 7], calling for the development of new tools to study the classical two-body problem.

Motivated by GW modeling, a host of techniques have been developed to solve the two-body problem in general relativity, including numerical relativity, which numerically solves the fully nonlinear Einstein equations [8]; gravitational self-force (GSF) theory, a perturbative method that applies when one body is much smaller than the other [9]; and Post-Newtonian (PN) and Post-Minkowskian (PM) theory, weak-field expansions that apply when the two bodies are widely separated [10, 11]. Historically, focus has been on the bound, inspiralling systems that are the dominant sources for GW detectors. However, the case of hyperbolic, scattering encounters is now of great interest: it is now known that data for scattering orbits can inform bound-orbit models using the effective one-body framework [11–17] or through an analytic continuation from scattering to bound observables [18–23], spurring the development of new particle physics tools [24–28] that have enabled analytical computations of the two-body scattering Hamiltonian and related observables at high PM order [29–50].

In the bound case, synergies between different methods have consistently helped drive progress [51, 52]. An important tool in those synergies has been the first law of binary black hole (BH) mechanics [53–60], which describes how a binary system responds to variations of its parameters (see also [61]). This law has played an important role in the most accurate GSF waveform model [62, 63] and in utilizing GSF results within PN, effective one body, and

numerical relativity calculations [64, 65]; see [66] for a review. For spinless particles, the binary’s response to variations is determined by a basis of observables $\mathcal{B}^<$ consisting of the periastron advance $\Delta\Phi$, the radial frequency Ω_r , and the averaged Detweiler redshift $\langle z \rangle$ [53, 57].

To date, a first law for scattering scenarios has not been derived. In this Letter, we establish such a law and find the corresponding basis of scattering observables $\mathcal{B}^>$. Two of these observables are well studied: the deflection angle χ and the time delay. We complete the basis with a third observable: the elapsed proper time $\Delta\tau$. Our approach is based on a pseudo-Hamiltonian formulation of GSF theory [58, 67–69]. This allows us to include dissipative contributions in the first law, unlike all previous formulations for bound orbits. By comparing to bound-orbit formulations, we also establish a novel analytic continuation between the elements of the scattering $\mathcal{B}^>$ and bound $\mathcal{B}^<$ bases of observables.

Finally, we link our calculations to high-energy physics methods, proving that the exponential representation of the classical S-matrix [37, 38] provides a generating functional for the basis of scattering observables and deriving a first law from a PM Hamiltonian.

Conventions We use geometric units with $G = c = 1$ and the $(-+++)$ metric signature.

First law in the probe limit.—In the GSF approach, the smaller body (of mass m_1) is treated as a point-particle perturbing the spacetime of the large body, which we take to be a Schwarzschild BH of mass m_2 . We first consider the probe limit (0SF order), in which the particle moves on a geodesic of the Schwarzschild metric $g_{\alpha\beta}^{\text{Schw}}$. The particle’s motion is governed by the geodesic Hamiltonian $H_0 = (1/2)g_{\text{Schw}}^{\mu\nu}p_\mu p_\nu$, where $p_\alpha = m_1 g_{\alpha\beta}^{\text{Schw}} dx^\beta/d\tau$ is the particle’s 4-momentum and τ is its proper time. Assuming, without loss of generality, that the motion lies on the equatorial plane $\theta = \pi/2$, we label the position of the particle with $x^\alpha(\tau) = (t(\tau), r(\tau), \pi/2, \varphi(\tau))$. Because

of Schwarzschild's Killing symmetries, the particle's energy and angular momentum $E = -p_{t,0}$ and $L = p_{\phi,0}$ are conserved (here and below, a subscript 0 indicates the on-shell geodesic value). We now consider unbound geodesic orbits that begin and end at $r = \infty$; such orbits have $E > m_1$ and $L > L_{\text{crit}}(E)$, where $L_{\text{crit}}(E)$ is a critical value of the angular momentum [70].

Following Carter's application of Hamilton-Jacobi theory [71], we use the constants of motion $P_i = (m_1, E, L)$ as canonical momenta and transform to canonical coordinates (X^i, P_i) using the type-2 generating function

$$W(t, r, \varphi; P_i) = -Et + L\varphi + I_{r,0}(r; P_i), \quad (1)$$

$$I_{r,0}(r; P_i) = \int_{r_m}^r dr p_{r,0}(r; P_i),$$

where $r_m(P_i)$ is the geodesic's minimum radius (i.e., closest approach to the BH). $g_{\text{Schw}}^{\mu\nu} p_{\mu,0} p_{\nu,0} = -m_1^2$ implies

$$p_{r,0}(r; P_i) = \frac{\sqrt{E^2 r^4 - r(r - 2m_2)(L^2 + m_1^2 r^2)}}{r(r - 2m_2)}. \quad (2)$$

In the coordinates (X^i, P_i) , where $X^i = \partial W / \partial P_i$, the Hamiltonian is simply its on-shell value, $H_0 = -m_1^2/2$, meaning that Hamilton's equations become [72]

$$m_1 \frac{dX^i}{d\tau} = \frac{\partial H_0}{\partial P_i} = -m_1 \delta_1^i. \quad (3)$$

Therefore X^2 and X^3 are constants, while X^1 is linear in τ . Since $X^i = \partial W / \partial P_i$, this implies

$$\left. \frac{\partial W}{\partial E} \right|_{\text{out}} = \left. \frac{\partial W}{\partial E} \right|_{\text{in}}, \quad \left. \frac{\partial W}{\partial L} \right|_{\text{out}} = \left. \frac{\partial W}{\partial L} \right|_{\text{in}},$$

$$\tau_{\text{out}} - \tau_{\text{in}} = - \left[\left. \frac{\partial W}{\partial m_1} \right|_{\text{out}} - \left. \frac{\partial W}{\partial m_1} \right|_{\text{in}} \right], \quad (4)$$

where “in” and “out” denote the initial, incoming state and final, outgoing state.

Equations (1) and (4) imply that the total changes in coordinate time, azimuthal angle, and proper time between initial and final states are

$$\Delta t_0 = t_{\text{out}} - t_{\text{in}} = \frac{\partial}{\partial E} [I_{r,0}(r_{\text{out}}; P_i) - I_{r,0}(r_{\text{in}}; P_i)], \quad (5)$$

$$\Delta \varphi_0 = \varphi_{\text{out}} - \varphi_{\text{in}} = - \frac{\partial}{\partial L} [I_{r,0}(r_{\text{out}}; P_i) - I_{r,0}(r_{\text{in}}; P_i)],$$

$$\Delta \tau_0 = \tau_{\text{out}} - \tau_{\text{in}} = - \frac{\partial}{\partial m_1} [I_{r,0}(r_{\text{out}}; P_i) - I_{r,0}(r_{\text{in}}; P_i)].$$

We are interested in the limit where the initial and final states are defined at past and future timelike infinity, with $r_{\text{in}} = \infty = r_{\text{out}}$, passing through the single radial turning point r_m . In this limit, $\Delta \varphi_0$ remains finite, but $I_{r,0}(r_{\text{in/out}}; P_i)$, Δt_0 , and $\Delta \tau_0$ all diverge. However, we

can define regularized versions. Using a convenient dimensionless regulator ϵ [22], we first define the scattering radial action

$$I_{r,0}^{>\epsilon}(P_i) = 2 \int_{r_m}^{+\infty} dr r^\epsilon p_{r,0}(r; P_i). \quad (6)$$

Intermediate results depend on the finite value of ϵ , but we obtain ϵ -independent observables in the limit $\epsilon \rightarrow 0$; when necessary, functions are first defined in regions of the complex- ϵ plane where integrals converge [e.g., $\text{Re}(\epsilon) < -1$ in Eq. (6)] and are then analytically continued to $\epsilon = 0$ [73]. In terms of $I_{r,0}^{>\epsilon}$ we can write the regularized $r_{\text{in/out}} \rightarrow \infty$ limit of Eq. (5) for the full scattering path as

$$\Delta \varphi_0^\epsilon = \pi + \chi_0^\epsilon = - \frac{\partial}{\partial L} I_{r,0}^{>\epsilon}(P_i), \quad (7)$$

where $\chi_0 = \lim_{\epsilon \rightarrow 0} \chi_0^\epsilon$ is the physical scattering angle, and

$$\Delta t_0^\epsilon = \frac{\partial}{\partial E} I_{r,0}^{>\epsilon}(P_i), \quad \Delta \tau_0^\epsilon = - \frac{\partial}{\partial m_1} I_{r,0}^{>\epsilon}(P_i). \quad (8)$$

Unlike χ_0^ϵ , the elapsed times Δt_0^ϵ and $\Delta \tau_0^\epsilon$ diverge as $\epsilon \rightarrow 0$. The associated physical observables, which are well defined when $\epsilon \rightarrow 0$, are *relative* measurements: the difference between $\Delta t_0(P_i)$ along the geodesic and $\Delta t_0(P_{i,\text{ref}})$ along some reference orbit; and these relative quantities will take the same values as if we had worked consistently with finite $r_{\text{in/out}}$ and only taken the limit $r_{\text{in/out}} \rightarrow \infty$ at the end of the calculation. In the Supplemental Material [74], we provide exact expressions for the geodesic scattering observables as well as the first few terms in their PM expansions (corresponding to $m_1 m_2 / L \ll 1$).

Finally, Eqs. (7) and (8) can be immediately combined into a single equation,

$$\delta I_{r,0}^{>\epsilon} = -(\pi + \chi_0) \delta L + \Delta t_0^\epsilon \delta E - \Delta \tau_0^\epsilon \delta m_1. \quad (9)$$

This is our first law for scattering geodesics. Here and below, we discard terms that vanish at $\epsilon = 0$, and equalities should be understood in this sense.

First law at all SF orders.—Beyond leading order in the mass ratio m_1/m_2 , the particle generates a metric perturbation $h_{\alpha\beta}$ on the Schwarzschild background. m_1 then moves on a geodesic of a certain *effective* metric $\tilde{g}_{\alpha\beta} = g_{\alpha\beta}^{\text{Schw}} + h_{\alpha\beta}^{\text{R}}$ [75], where $h_{\alpha\beta}^{\text{R}}$ is a certain regular piece of $h_{\alpha\beta}$. At 1SF order, we can write $h_{\alpha\beta}^{\text{R}}$ in terms of the Detweiler-Whiting Green's function $G_{\text{R}}^{\alpha\beta\alpha'\beta'}$ [76]:

$$h_{\text{R}}^{\alpha\beta}(x^\mu; \Gamma) = \frac{1}{m_1} \int_{\Gamma} G_{\text{R}}^{\alpha\beta\alpha'\beta'}(x^\mu, x'^\mu(\tilde{\tau}')) \tilde{p}_{\alpha'} \tilde{p}_{\beta'} d\tilde{\tau}'. \quad (10)$$

Here $\tilde{p}_\alpha := \tilde{g}_{\alpha\beta} dx^\beta / d\tilde{\tau}$, $\tilde{\tau}$ is proper time in $\tilde{g}_{\alpha\beta}$, and Γ denotes the particle's phase-space trajectory. Due to curvature-induced tail effects, $G_{\text{R}}^{\alpha\beta\alpha'\beta'}$ is nonzero for all points x'^μ in the past of x^μ , implying $h_{\text{R}}^{\alpha\beta}$ at a point on

Γ depends on the entire prior history of Γ . At higher SF orders, there is no known Green's-function form analogous to (10), but an appropriate $h_{\alpha\beta}^R(x^\mu; \Gamma)$ exists at all SF orders [77].

In this setting, we again consider scattering orbits with initial parameters $P_i = (m_1, E, L)$. The particle's energy and angular momentum evolve due to dissipation, but the orbit remains planar ($\theta = \pi/2$, $\tilde{p}_\theta = 0$). For L above a critical threshold, the orbit remains close to a Schwarzschild geodesic with the same initial P_i [70, 78].

Since the motion is geodesic in $\tilde{g}_{\mu\nu}$, it obeys Hamilton's equations with the test-mass pseudo-Hamiltonian $H = (1/2)\tilde{g}^{\mu\nu}\tilde{p}_\mu\tilde{p}_\nu$ [58, 67, 68]. However, we deviate from [58, 67] by restricting to the 4D phase space $(x^A, \tilde{p}_A)$ satisfying the on-shell condition $H = -m_1^2/2$, with $x^A = (r, \varphi)$. Solving the on-shell condition for $\tilde{p}_t = -\mathcal{H}(t, x^A, \tilde{p}_A; \Gamma)$ gives the new pseudo-Hamiltonian

$$\begin{aligned} \mathcal{H} &= \frac{1}{\tilde{g}^{tt}} \left[\tilde{g}^{tA} \tilde{p}_A - \sqrt{(\tilde{g}^{tA} \tilde{p}_A)^2 - \tilde{g}^{tt}(\tilde{g}^{AB} \tilde{p}_A \tilde{p}_B + m_1^2)} \right] \\ &= -p_t(r, \tilde{p}_A) - \frac{h_{\text{R}}^{\mu\nu}(t, x^A; \Gamma) \tilde{p}_\mu \tilde{p}_\nu}{2g_{\text{Schw}}^{tt} p_t} + \mathcal{O}\left(\frac{m_1^2}{m_2^2}\right), \end{aligned} \quad (11)$$

with t now the parameter along the trajectory. $\mathcal{H}$ is referred to as a pseudo-Hamiltonian because it depends on the trajectory $\Gamma = \{(x^A(t), \tilde{p}_A(t)) \mid t \in \mathbb{R}\}$. Hamilton's equations in this context read

$$\frac{dx^A}{dt} = \left[\frac{\partial \mathcal{H}}{\partial \tilde{p}_A} \right] \quad \text{and} \quad \frac{d\tilde{p}_A}{dt} = - \left[\frac{\partial \mathcal{H}}{\partial x^A} \right], \quad (12)$$

where $[\cdot]$ indicates specification of Γ as the self-consistent trajectory [79] passing through $(x^A, \tilde{p}_A)$; prior to that specification, Γ is treated as independent, and derivatives do not act on it. We go further by replacing m_1 by m'_1 in Eq. (10), setting $m'_1 = m_1$ only when $[\cdot]$ is applied. Importantly, Eq. (12) captures the full dynamics, including dissipation, unlike an ordinary Hamiltonian description.

We derive the first law from $\mathcal{H}$. Doing so will require its relationship to the redshift z :

$$z := \frac{d\tilde{\tau}}{dt} = \left[\frac{\partial \mathcal{H}}{\partial m_1} \right]. \quad (13)$$

To establish this relationship, we consider the normalization condition

$$-m_1 = \tilde{p}_\mu \frac{dx^\mu}{d\tilde{\tau}} = \left(-\mathcal{H} + \tilde{p}_A \left[\frac{\partial \mathcal{H}}{\partial \tilde{p}_A} \right] \right) z^{-1}, \quad (14)$$

where we used $\tilde{p}_t = -\mathcal{H}$ together with (12). Next we note the first line of (11) shows that, at fixed m'_1 , $\mathcal{H}$ is a homogeneous function of $(m_1, \tilde{p}_A)$ of order 1. Euler's homogeneous function theorem hence implies $\mathcal{H} = m_1 \frac{\partial \mathcal{H}}{\partial m_1} + \tilde{p}_A \frac{\partial \mathcal{H}}{\partial \tilde{p}_A}$. Comparing this with (14), we obtain (13).

Now, to derive the first law, we loosely follow [57] by considering how $\mathcal{H}$ changes under variations δP_i of the

initial data, with $\delta x_{\text{in}}^A = 0$. Given (12) and (13), we find

$$\begin{aligned} [\delta \mathcal{H}] &= \left[\frac{\partial \mathcal{H}}{\partial x^A} \delta x^A \right] + \left[\frac{\partial \mathcal{H}}{\partial \tilde{p}_A} \delta \tilde{p}_A \right] + \left[\frac{\partial \mathcal{H}}{\partial m_1} \delta m_1 \right] \\ &= -\frac{d\tilde{p}_A}{dt} \delta x^A + \frac{dx^A}{dt} \delta \tilde{p}_A + \frac{d\tilde{\tau}}{dt} \delta m_1. \end{aligned} \quad (15)$$

Since $h_{\mu\nu}^R$ vanishes for an inertial particle in Minkowski [75], its contribution to $\tilde{p}_A$ and $\mathcal{H}$ vanishes in the initial state, such that $\tilde{p}_\varphi^{\text{in}} = p_{\varphi,0}^{\text{in}} = L$ and $\mathcal{H}_{\text{in}} = -p_{t,0}^{\text{in}} = E$. We isolate δL and δE in (15) by defining 'interaction' quantities $\tilde{p}_\varphi := \tilde{p}_\varphi - L$, $\tilde{p}_r := \tilde{p}_r$, and $\mathcal{H} := \mathcal{H} - E$, such that

$$\delta E + [\delta \mathcal{H}] = -\frac{d\tilde{p}_A}{dt} \delta x^A + \frac{dx^A}{dt} \delta \tilde{p}_A + \frac{d\varphi}{dt} \delta L + \frac{d\tilde{\tau}}{dt} \delta m_1. \quad (16)$$

Next, we integrate (16) along the physical scattering trajectory from $t = -\infty$ to $t = +\infty$, introducing the regularized integral $\langle f \rangle_\Gamma := \int_\Gamma dt r^\epsilon f$ as in the 0SF case. Integrating the term $\frac{d\tilde{p}_A}{dt} \delta x^A$ by parts, we obtain

$$\begin{aligned} \Delta t^\epsilon \delta E + \langle [\delta \mathcal{H}] \rangle_\Gamma &= \delta \left\langle \tilde{p}_A \frac{dx^A}{dt} \right\rangle_\Gamma + \Delta \varphi^\epsilon \delta L + \Delta \tilde{\tau}^\epsilon \delta m_1, \end{aligned} \quad (17)$$

where $\Delta t^\epsilon := \langle 1 \rangle_\Gamma$, $\Delta \varphi^\epsilon := \langle \frac{d\varphi}{dt} \rangle_\Gamma$, and $\Delta \tilde{\tau}^\epsilon := \langle \frac{d\tilde{\tau}}{dt} \rangle_\Gamma$. We have discarded boundary terms by choosing $\text{Re}(\epsilon)$ sufficiently negative and discarded terms that arise from derivatives acting on r^ϵ because they vanish when analytically continued to $\epsilon = 0$. Defining also the regularized 'interaction' action

$$I^{>,\epsilon} := \int_\Gamma dt r^\epsilon \tilde{p}_A \frac{dx^A}{dt} = I_r^{>,\epsilon} + \int_\Gamma d\varphi \tilde{p}_\varphi, \quad (18)$$

with $I_r^{>,\epsilon} := \int_\Gamma dr r^\epsilon \tilde{p}_r$, we rewrite (17) as

$$\Delta t^\epsilon \delta E + \langle [\delta \mathcal{H}] \rangle_\Gamma = \delta I^{>,\epsilon} + \Delta \varphi^\epsilon \delta L + \Delta \tilde{\tau}^\epsilon \delta m_1. \quad (19)$$

We can rewrite Eq. (19) in an alternative form by absorbing $\langle [\delta \mathcal{H}] \rangle_\Gamma$ into 'renormalized' variables $\{I_{\text{ren}}^{>,\epsilon}, E_{\text{ren}}, L_{\text{ren}}\}$. Following [58]'s treatment of the bound case, we define renormalized variables

$$E_{\text{ren}} = \lambda E, \quad L_{\text{ren}} = \lambda L, \quad I_{\text{ren}}^{>,\epsilon} = \lambda I^{>,\epsilon}. \quad (20)$$

Choosing $\lambda(P_i)$ appropriately to eliminate $\langle [\delta \mathcal{H}] \rangle$ from Eq. (19), we are left with

$$\delta I_{\text{ren}}^{>,\epsilon} = -(\pi + \chi^\epsilon) \delta L_{\text{ren}} + \Delta t^\epsilon \delta E_{\text{ren}} - \Delta \tilde{\tau}^\epsilon \delta m_1; \quad (21)$$

see the Supplemental Material [74] for more details.

Equation (21) is the first law for scattering orbits, valid at all SF orders and including all dissipative effects. To help understand the renormalization of the variables, we observe that the first law defines a sense of conjugacy between variables and observables, just as in the first

law of thermodynamics. In that sense, the renormalized variables are the ones conjugate to the physical observables. In Ref. [80], we show that in the conservative sector, this sense of conjugacy reduces to the usual sense in Hamiltonian mechanics: the renormalized variables are the true, invariant action variables that are canonically conjugate to the system's action-angles. The need for this renormalization stems from the fact that, as highlighted in Ref. [68], if a system can be equivalently described by both a pseudo-Hamiltonian and a Hamiltonian, then variables which are conjugate in one description are not generally conjugate in the other. As a consequence, the momenta $\tilde{p}_\mu$, from which E , L , and $I^{>,\epsilon}$ are built, are *not* the canonical momenta in a Hamiltonian description of the conservative sector (i.e., they are not the momenta one would define from a Lagrangian for the conservative sector). We refer to Refs. [69, 80] for details.

From scattering to bound.—There is a well-known analytic continuation between the deflection angle χ for unbound orbits and the periastron advance $\Delta\Phi$ for bound orbits, as well as between the scattering and bound radial actions [18, 19, 22]. Here, using the first laws for unbound and bound motion, we extend these analytic continuations to include all the observables in the scattering and bound bases, $\mathcal{B}^> = (\chi, \Delta t^\epsilon, \Delta\tau^\epsilon)$ and $\mathcal{B}^< = (\Delta\Phi, \Omega_r, \langle z \rangle)$. We limit our analysis to 0SF order, as the analytic continuation for the radial action is not known to be valid beyond 0SF order due to nonlocal-in-time tail effects [21, 43].

We write the first law for bound geodesics in terms of the bound radial action,

$$I_{r,0}^<(P_i) = 2 \int_{r_-(P_i)}^{r_+(P_i)} dr p_{r,0}(r; P_i), \quad (22)$$

where $r_\mp$ are the orbit's minimum and maximum radius (i.e., the radii at periastris and apoapsis). Following the same arguments as for unbound orbits, one can write the accumulated φ , t , and τ over a single radial period ($T_{r,0} = 2\pi/\Omega_{r,0}$) as derivatives of $I_{r,0}^<$, leading to the first law for bound orbits [56, 57]:

$$\delta I_{r,0}^< = -(2\pi + \Delta\Phi_0)\delta L + \frac{2\pi}{\Omega_{r,0}}\delta E - \frac{2\pi}{\Omega_{r,0}}\langle z \rangle_0 \delta m_1, \quad (23)$$

where $\langle z \rangle_0 := (1/T_{r,0}) \int_0^{T_{r,0}} dt (d\tau/dt)_0$.

Knowing the scatter-to-bound map for the radial action [19, 22, 39],

$$I_{r,0}^<(P_i) = \lim_{\epsilon \rightarrow 0} [I_{r,0}^{>,\epsilon}(E, L, m_1) - I_{r,0}^{>,\epsilon}(E, -L, m_1)], \quad (24)$$

and comparing Eq. (23) to Eq. (9), we immediately conclude that there is an analytic continuation between the

full set of scattering and bound observables:

$$\begin{aligned} \Delta\Phi_0 &= \chi_0(E, L, m_1) + \chi_0(E, -L, m_1), \\ \frac{2\pi}{\Omega_{r,0}} &= \lim_{\epsilon \rightarrow 0} [\Delta t_0^\epsilon(E, L, m_1) - \Delta t_0^\epsilon(E, -L, m_1)], \\ \frac{2\pi\langle z \rangle_0}{\Omega_{r,0}} &= \lim_{\epsilon \rightarrow 0} [\Delta\tau_0^\epsilon(E, L, m_1) - \Delta\tau_0^\epsilon(E, -L, m_1)]. \end{aligned} \quad (25)$$

We note that the infrared divergences in Δt_0^ϵ and $\Delta\tau_0^\epsilon$ cancel in these expressions because the divergences are independent of L ; see the Supplemental Material [74].

First law from the S-matrix.—In this section, we put our first law in the broader context of the quantum S-matrix description of the classical two-body problem [81]. Given the two-body initial state of well-separated massive point particles $|\Psi_{\text{in}}\rangle = |p_1 p_2\rangle$ of mass m_1 and m_2 , the action of the unitary S-matrix operator,

$$\hat{S} = \mathcal{T} \exp \left(-\frac{i}{\hbar} \int dt H_{\text{int}}(t) \right), \quad (26)$$

describes the time evolution of the state in terms of the interaction Hamiltonian H_{int} , with $\mathcal{T}$ denoting time ordering. The classical two-body scattering dynamics in the asymptotic $\hbar \rightarrow 0$ limit is equivalently obtained by evaluating the action, and therefore H_{int} , on-shell.

Motivated by (26), we define the exponential representation $\hat{S} = \exp(i\hat{N}/\hbar)$ [37, 38], where $\hat{N}$ is a Hermitian operator. We then study the *real-valued* two-body matrix element

$$N(\mathbb{E}, q, m_1, m_2) := \langle p'_1 p'_2 | \hat{N} | p_1 p_2 \rangle, \quad (27)$$

where we defined $|\Psi_{\text{out}}\rangle = |p'_1 p'_2\rangle$, the initial *total* energy $\mathbb{E}$ of both particles, and the exchanged momentum $q^\mu = p'_1{}^\mu - p_1{}^\mu = p_2{}^\mu - p'_2{}^\mu$. To make contact with the incoming total angular momentum $\mathbb{L} = (m_1 m_2 \sqrt{\gamma^2 - 1} b)/\mathbb{E}$, where $\gamma := \frac{p_1^\mu + p_1'^\mu}{2} \frac{p_{2\mu} + p_{2\mu}'}{2}$ and b is the impact parameter in the center of mass (CM) frame, we perform the Fourier transform

$$\begin{aligned} N^{>,\epsilon}(\mathbb{E}, \mathbb{L}, \{m_a\}) \\ = \frac{1}{4m_1 m_2 \sqrt{\gamma^2 - 1}} \int \frac{d^{2+2\epsilon} q}{(2\pi)^{2+2\epsilon}} e^{-i \frac{b(\mathbb{L}) \cdot q}{\hbar}} N(\mathbb{E}, q, \{m_a\}), \end{aligned} \quad (28)$$

using dimensional regularization with $d = 4 + 2\epsilon$. Infrared, $1/\epsilon$ divergences arise due to the long-range nature of gravity, but their analytic structure is understood [82].

In complete generality, the expectation value (27) is a function of the kinematic data $(\mathbb{E}, \mathbb{L}, \{m_a\})$, and its variation in the phase space is

$$\delta N^{>,\epsilon} = c_{\mathbb{L}}^\epsilon \delta \mathbb{L} + c_{\mathbb{E}}^\epsilon \delta \mathbb{E} + \sum_{a=1,2} c_{m_a}^\epsilon \delta m_a, \quad (29)$$

where $c_{\mathbb{L}}^\epsilon$, $c_{\mathbb{E}}^\epsilon$, and $c_{m_a}^\epsilon$ are gauge-invariant coefficients.

Using insights from the PM Hamiltonian description [30] and the relation with the radial action [38], we now identify the coefficients with the observables $\mathcal{B}^>$. First, by matching the scattering angle χ in the CM frame, it was shown that the matrix element (27) in the conservative case agrees with the radial action up to a constant proportional to $\mathbb{L}$ [38]:

$$N^{>,\epsilon} \Big|_{\text{cons}} = \underbrace{\int_{\mathcal{C}_r^{>,\epsilon}} dr \tilde{p}_{r,\text{c.m.}}(r; \mathbb{E}, \mathbb{L}, \{m_a\})}_{\mathbb{I}_r^{>,\epsilon}} + \pi \mathbb{L}, \quad (30)$$

where $\tilde{p}_{r,\text{c.m.}}$ is the radial relative momentum in the CM frame and $\mathcal{C}_r^{>,\epsilon}$ is the contour of integration for scattering orbits, which implicitly includes a regulator ϵ inherited from the dimensional regularization. In the Supplemental Material [74], using the PM conservative Hamiltonian and its symmetries in the CM frame [19, 30], we then provide a proof of the following conservative first law for the two-body scattering problem,

$$\delta N^{>,\epsilon} = -\chi \delta \mathbb{L} + \Delta t^\epsilon \delta \mathbb{E} - \sum_{a=1,2} \Delta \tau_a^\epsilon \delta m_a, \quad (31)$$

where $\Delta \tau_a^\epsilon$ is the elapsed proper time of particle a and Δt^ϵ is the elapsed global time.

If we appeal to $N^{>,\epsilon} = \mathbb{I}_r^{>,\epsilon} + \pi \mathbb{L}$ and restrict to variations with $\delta m_2 = 0$ [83], then we see Eq. (31) is structurally identical to our previous first law (21). However, the quantities in these laws might differ. Even in the conservative sector, the two-body incoming energy $\mathbb{E}$ and angular momentum $\mathbb{L}$ might not agree with the renormalized one-body, SF counterparts E_{ren} and L_{ren} . Moreover, while $N^{>,\epsilon}$ is computed here in the CM frame, GSF calculations might be in the initial rest frame of the heavy BH [84–86] or in any ‘nearby’ frame (including the CM frame); the frame of a GSF calculation is implicitly determined by the choice of gauge for the metric perturbations. However, at 0SF order, we can trivially identify $\mathbb{E}$ with $E + m_2$, $\mathbb{L}$ with L , and the CM frame with the heavy BH frame, as the relative radial momentum $\tilde{p}_{r,\text{c.m.}}$ coincides with the geodesic one $p_{r,0}$. Then (31) identically matches (9), and we have $\chi^\epsilon \rightarrow \chi_0^\epsilon$, $\Delta t^\epsilon \rightarrow \Delta t_0^\epsilon$, and $\Delta \tau_1^\epsilon \rightarrow \Delta \tau_0^\epsilon$. At n SF order, the matching with our SF first law is more challenging as the dynamics of the heavy BH (as well as the choice of frame) is encoded in a nontrivial way in the metric perturbations [87, 88]. We leave study of this to future work.

Incorporating dissipation in this framework is possible by combining the in-in expectation value [81] with the exponential representation of $\hat{S}$ [38]: for every observable $\mathcal{O}$ [89],

$$\begin{aligned} \langle \Delta \mathcal{O} \rangle &= \langle \Psi_{\text{in}} | \hat{S}^\dagger \mathcal{O} \hat{S} | \Psi_{\text{in}} \rangle - \langle \Psi_{\text{in}} | \mathcal{O} | \Psi_{\text{in}} \rangle \\ &= \sum_{j=1}^{+\infty} \frac{(-i)^j}{\hbar^j j!} \underbrace{[\hat{N}, [\hat{N}, \dots, [\hat{N}, \mathcal{O}] \dots]]}_{j \text{ times}}, \end{aligned} \quad (32)$$

$$\begin{aligned} \delta N^{>,\epsilon}(\mathbb{E}, \mathbb{L}, \{m_a\}) &= -\chi \delta \mathbb{L} + \Delta t^\epsilon \delta \mathbb{E} - \sum_{a=1,2} \Delta \tau_a^\epsilon \delta m_a \\ &\downarrow \delta N^{<}(\mathbb{E}, \mathbb{L}, \{m_a\}) = \delta N^{>,\epsilon}(\mathbb{E}, \mathbb{L}, \{m_a\}) - \delta N^{>,\epsilon}(\mathbb{E}, -\mathbb{L}, \{m_a\}) \\ \delta N^{<}(\mathbb{E}, \mathbb{L}, \{m_a\}) &= -\Delta \Phi \delta \mathbb{L} + \frac{2\pi}{\Omega_r} \delta \mathbb{E} - \sum_{a=1,2} \frac{2\pi}{\Omega_r} \langle z_a \rangle \delta m_a \end{aligned}$$

FIG. 1. Relation between the basis of scattering observables (coordinate time difference Δt^ϵ , elapsed proper time $\Delta \tau^\epsilon$, and deflection angle χ) and the corresponding bound-orbit ones (radial frequency Ω_r , averaged redshift $\langle z \rangle$, and periastron advance $\Delta \Phi$) for the classical gravitational two-body problem.

where now also the $\hat{N}$ matrix elements with on-shell gravitons are relevant. This suggests a physical principle to connect the coefficients ($c_{\mathbb{L}}^\epsilon, c_{\mathbb{E}}^\epsilon, c_{m_a}^\epsilon$) to observables at all orders, with dissipation included; see for example Eq. (3.48) of [38].

Conclusion.—In recent years, the study of unbound orbits through the S-matrix formalism has transformed the gravitational two-body problem. In this Letter, we developed a powerful new tool for such studies: an extension of the first law of binary BH mechanics to the unbound (and dissipative) case. Our derivation at all SF orders utilizes a novel version of the pseudo-Hamiltonian formalism [58, 67, 69]. More generally, we showed how the first law can be derived from the variation of the classical S-matrix in the phase space of the kinematic data $P_i = (\mathcal{E}, L, \{m_a\})$; see Fig. 1. In that context, the S-matrix can be interpreted as a generating functional of classical observables. Among those observables, we have highlighted the elapsed proper time $\Delta \tau^\epsilon$ as a new, core element of the (regularized) basis of scattering observables $\mathcal{B}^> = (\chi, \Delta t^\epsilon, \Delta \tau^\epsilon)$.

Using the relation between the scattering and bound radial action, we also established a full correspondence (25) between the bases of scattering observables $\mathcal{B}^>$ and bound observables $\mathcal{B}^<$ at 0SF order, as again summarized in Fig. 1. This extends the well-known map between the deflection angle and periastron advance.

Given the first law’s varied applications for bound orbits [62, 63, 90–111], we expect our work to open many new avenues for scattering calculations. We particularly encourage self-force scattering calculations [70, 78, 112–114] of the observables Δt^ϵ and $\Delta \tau^\epsilon$. For bound orbits, the 1SF conservative Hamiltonian can be calculated directly from the averaged redshift $\langle z \rangle$, and post-adiabatic waveform models [62, 115, 116] can be written in a gauge-invariant form with $\langle z \rangle$ as an invariant building block [80, 117]. This implies that if the analytic continuation between $\Delta \tau^\epsilon$ and $\langle z \rangle$ can be extended to 1SF order, then scattering calculations of $\Delta \tau^\epsilon$ can provide direct inputs to bound-orbit self-force waveform models.

Natural extensions also present themselves. First, one might consider scattering orbits of spinning BHs [55, 60, 118]. Second, we considered only two-body matrix ele-

ments, but nothing prevents us from studying the variation of matrix elements involving on-shell graviton states, which should be related to the gravitational waveform. Finally, we hope that linking the first laws for scattering and bound orbits beyond OSF can shed light on the tail effects that have limited the applicability of scatter-to-bound maps [43, 49].

Acknowledgments. We thank Alex Le Tiec for comments on a draft of this Letter. RG would like to thank A. Ilderton, M. Poratti, K. Rajeev and C. Shi for useful discussions. JL and AP gratefully acknowledge helpful conversations with S. Isoyama and T. Tanaka about pseudo-Hamiltonian dynamics and with L. Barack, C. Kavanagh, R. Porto, D. Usseglio, and M. van de Meent about scatter-to-bound maps. RG is grateful to the Mainz Institute of Theoretical Physics (MITP) of the Cluster of Excellence PRISMA+ (Project ID 390831469), for its hospitality and its partial support during the completion of this work. JL acknowledges the support of an STFC studentship. AP acknowledges the support of a Royal Society University Research Fellowship and a UKRI Frontier Research Grant (as selected by the ERC) under the Horizon Europe Guarantee scheme [grant number EP/Y008251/1].

* rgonzo@ed.ac.uk

† J.E.Lewis@soton.ac.uk

‡ A.Pound@soton.ac.uk

- [1] J. Luo *et al.* (TianQin), *TianQin: a space-borne gravitational wave detector*, *Class. Quant. Grav.* **33** (2016) 035010, [arXiv:1512.02076 \[astro-ph.IM\]](#).
- [2] W.-H. Ruan, Z.-K. Guo, R.-G. Cai, and Y.-Z. Zhang, *Taiji program: Gravitational-wave sources*, *Int. J. Mod. Phys. A* **35** (2020) 2050075, [arXiv:1807.09495 \[gr-qc\]](#).
- [3] D. Reitze *et al.*, *Cosmic Explorer: The U.S. Contribution to Gravitational-Wave Astronomy beyond LIGO*, *Bull. Am. Astron. Soc.* **51** (2019) 035, [arXiv:1907.04833 \[astro-ph.IM\]](#).
- [4] M. Maggiore *et al.*, *Science Case for the Einstein Telescope*, *JCAP* **03** (2020) 050, [arXiv:1912.02622 \[astro-ph.CO\]](#).
- [5] P. A. Seoane *et al.* (LISA), *Astrophysics with the Laser Interferometer Space Antenna*, *Living Rev. Rel.* **26** (2023) 2, [arXiv:2203.06016 \[gr-qc\]](#).
- [6] M. Pürrer and C.-J. Haster, *Gravitational waveform accuracy requirements for future ground-based detectors*, *Phys. Rev. Res.* **2** (2020) 023151, [arXiv:1912.10055 \[gr-qc\]](#).
- [7] A. Dhani, S. Völkel, A. Buonanno, H. Estelles, J. Gair, H. P. Pfeiffer, L. Pompili, and A. Toubiana, *Systematic Biases in Estimating the Properties of Black Holes Due to Inaccurate Gravitational-Wave Models*, (2024), [arXiv:2404.05811 \[gr-qc\]](#).
- [8] T. Baumgarte and S. Shapiro, *Numerical Relativity: Solving Einstein's Equations on the Computer* (Cambridge University Press, 2010).
- [9] L. Barack and A. Pound, *Self-force and radiation reaction in general relativity*, *Rept. Prog. Phys.* **82** (2019) 016904, [arXiv:1805.10385 \[gr-qc\]](#).
- [10] L. Blanchet, *Gravitational Radiation from Post-Newtonian Sources and Inspiralling Compact Binaries*, *Living Rev. Rel.* **17** (2014) 2, [arXiv:1310.1528 \[gr-qc\]](#).
- [11] T. Damour, *High-energy gravitational scattering and the general relativistic two-body problem*, *Phys. Rev. D* **97** (2018) 044038, [arXiv:1710.10599 \[gr-qc\]](#).
- [12] A. Buonanno and T. Damour, *Effective one-body approach to general relativistic two-body dynamics*, *Phys. Rev. D* **59** (1999) 084006, [arXiv:gr-qc/9811091](#).
- [13] A. Buonanno and T. Damour, *Transition from inspiral to plunge in binary black hole coalescences*, *Phys. Rev. D* **62** (2000) 064015, [arXiv:gr-qc/0001013](#).
- [14] T. Damour, *Gravitational scattering, post-Minkowskian approximation and Effective One-Body theory*, *Phys. Rev. D* **94** (2016) 104015, [arXiv:1609.00354 \[gr-qc\]](#).
- [15] M. Khalil, A. Buonanno, J. Steinhoff, and J. Vines, *Energetics and scattering of gravitational two-body systems at fourth post-Minkowskian order*, *Phys. Rev. D* **106** (2022) 024042, [arXiv:2204.05047 \[gr-qc\]](#).
- [16] A. Buonanno, G. U. Jakobsen, and G. Mogull, *Post-Minkowskian theory meets the spinning effective-one-body approach for two-body scattering*, *Phys. Rev. D* **110** (2024) 044038, [arXiv:2402.12342 \[gr-qc\]](#).
- [17] A. Buonanno, G. Mogull, R. Patil, and L. Pompili, *Post-Minkowskian Theory Meets the Spinning Effective-One-Body Approach for Bound-Orbit Waveforms*, (2024), [arXiv:2405.19181 \[gr-qc\]](#).
- [18] G. Kälín and R. A. Porto, *From Boundary Data to Bound States*, *JHEP* **01** (2020) 072, [arXiv:1910.03008 \[hep-th\]](#).
- [19] G. Kälín and R. A. Porto, *From boundary data to bound states. Part II. Scattering angle to dynamical invariants (with twist)*, *JHEP* **02** (2020) 120, [arXiv:1911.09130 \[hep-th\]](#).
- [20] M. V. S. Saketh, J. Vines, J. Steinhoff, and A. Buonanno, *Conservative and radiative dynamics in classical relativistic scattering and bound systems*, *Phys. Rev. Res.* **4** (2022) 013127, [arXiv:2109.05994 \[gr-qc\]](#).
- [21] G. Cho, G. Kälín, and R. A. Porto, *From boundary data to bound states. Part III. Radiative effects*, *JHEP* **04** (2022) 154, [Erratum: *JHEP* **07**, 002 (2022)], [arXiv:2112.03976 \[hep-th\]](#).
- [22] R. Gonzo and C. Shi, *Boundary to bound dictionary for generic Kerr orbits*, *Phys. Rev. D* **108** (2023) 084065, [arXiv:2304.06066 \[hep-th\]](#).
- [23] T. Adamo, R. Gonzo, and A. Ilderton, *Gravitational bound waveforms from amplitudes*, *JHEP* **05** (2024) 034, [arXiv:2402.00124 \[hep-th\]](#).
- [24] Z. Bern, C. Cheung, R. Roiban, C.-H. Shen, M. P. Solon, and M. Zeng, *Black Hole Binary Dynamics from the Double Copy and Effective Theory*, *JHEP* **10** (2019) 206, [arXiv:1908.01493 \[hep-th\]](#).
- [25] N. E. J. Bjerrum-Bohr, P. H. Damgaard, L. Plante, and P. Vanhove, *The SAGEX review on scattering amplitudes Chapter 13: Post-Minkowskian expansion from scattering amplitudes*, *J. Phys. A* **55** (2022) 443014, [arXiv:2203.13024 \[hep-th\]](#).
- [26] A. Buonanno, M. Khalil, D. O'Connell, R. Roiban, M. P. Solon, and M. Zeng, *Snowmass White Paper: Gravitational Waves and Scattering Amplitudes*, in *Snowmass 2021* (2022) [arXiv:2204.05194 \[hep-th\]](#).

- [27] D. A. Kosower, R. Monteiro, and D. O’Connell, *The SAGEX review on scattering amplitudes Chapter 14: Classical gravity from scattering amplitudes*, *J. Phys. A* **55** (2022) 443015, [arXiv:2203.13025 \[hep-th\]](#).
- [28] P. Di Vecchia, C. Heissenberg, R. Russo, and G. Veneziano, *The gravitational eikonal: From particle, string and brane collisions to black-hole encounters*, *Phys. Rept.* **1083** (2024) 1, [arXiv:2306.16488 \[hep-th\]](#).
- [29] D. Neill and I. Z. Rothstein, *Classical Space-Times from the S Matrix*, *Nucl. Phys. B* **877** (2013) 177, [arXiv:1304.7263 \[hep-th\]](#).
- [30] C. Cheung, I. Z. Rothstein, and M. P. Solon, *From Scattering Amplitudes to Classical Potentials in the Post-Minkowskian Expansion*, *Phys. Rev. Lett.* **121** (2018) 251101, [arXiv:1808.02489 \[hep-th\]](#).
- [31] Z. Bern, C. Cheung, R. Roiban, C.-H. Shen, M. P. Solon, and M. Zeng, *Scattering Amplitudes and the Conservative Hamiltonian for Binary Systems at Third Post-Minkowskian Order*, *Phys. Rev. Lett.* **122** (2019) 201603, [arXiv:1901.04424 \[hep-th\]](#).
- [32] Z. Bern, J. Parra-Martinez, R. Roiban, M. S. Ruf, C.-H. Shen, M. P. Solon, and M. Zeng, *Scattering Amplitudes and Conservative Binary Dynamics at $\mathcal{O}(G^4)$* , *Phys. Rev. Lett.* **126** (2021) 171601, [arXiv:2101.07254 \[hep-th\]](#).
- [33] Z. Bern, J. Parra-Martinez, R. Roiban, M. S. Ruf, C.-H. Shen, M. P. Solon, and M. Zeng, *Scattering Amplitudes, the Tail Effect, and Conservative Binary Dynamics at $\mathcal{O}(G^4)$* , *Phys. Rev. Lett.* **128** (2022) 161103, [arXiv:2112.10750 \[hep-th\]](#).
- [34] P. Di Vecchia, C. Heissenberg, R. Russo, and G. Veneziano, *The eikonal approach to gravitational scattering and radiation at $\mathcal{O}(G^3)$* , *JHEP* **07** (2021) 169, [arXiv:2104.03256 \[hep-th\]](#).
- [35] P. Di Vecchia, C. Heissenberg, R. Russo, and G. Veneziano, *Radiation Reaction from Soft Theorems*, *Phys. Lett. B* **818** (2021) 136379, [arXiv:2101.05772 \[hep-th\]](#).
- [36] N. E. J. Bjerrum-Bohr, P. H. Damgaard, L. Planté, and P. Vanhove, *The amplitude for classical gravitational scattering at third Post-Minkowskian order*, *JHEP* **08** (2021) 172, [arXiv:2105.05218 \[hep-th\]](#).
- [37] P. H. Damgaard, L. Plante, and P. Vanhove, *On an exponential representation of the gravitational S-matrix*, *JHEP* **11** (2021) 213, [arXiv:2107.12891 \[hep-th\]](#).
- [38] P. H. Damgaard, E. R. Hansen, L. Planté, and P. Vanhove, *Classical observables from the exponential representation of the gravitational S-matrix*, *JHEP* **09** (2023) 183, [arXiv:2307.04746 \[hep-th\]](#).
- [39] G. Kälin, Z. Liu, and R. A. Porto, *Conservative Dynamics of Binary Systems to Third Post-Minkowskian Order from the Effective Field Theory Approach*, *Phys. Rev. Lett.* **125** (2020) 261103, [arXiv:2007.04977 \[hep-th\]](#).
- [40] C. Dlapa, G. Kälin, Z. Liu, and R. A. Porto, *Conservative Dynamics of Binary Systems at Fourth Post-Minkowskian Order in the Large-Eccentricity Expansion*, *Phys. Rev. Lett.* **128** (2022) 161104, [arXiv:2112.11296 \[hep-th\]](#).
- [41] F. Febres Cordero, M. Kraus, G. Lin, M. S. Ruf, and M. Zeng, *Conservative Binary Dynamics with a Spinning Black Hole at $\mathcal{O}(G^3)$ from Scattering Amplitudes*, *Phys. Rev. Lett.* **130** (2023) 021601, [arXiv:2205.07357 \[hep-th\]](#).
- [42] C. Dlapa, G. Kälin, Z. Liu, J. Neef, and R. A. Porto, *Radiation Reaction and Gravitational Waves at Fourth Post-Minkowskian Order*, *Phys. Rev. Lett.* **130** (2023) 101401, [arXiv:2210.05541 \[hep-th\]](#).
- [43] C. Dlapa, G. Kälin, Z. Liu, and R. A. Porto, *Local in Time Conservative Binary Dynamics at Fourth Post-Minkowskian Order*, *Phys. Rev. Lett.* **132** (2024) 221401, [arXiv:2403.04853 \[hep-th\]](#).
- [44] G. U. Jakobsen and G. Mogull, *Linear response, Hamiltonian, and radiative spinning two-body dynamics*, *Phys. Rev. D* **107** (2023) 044033, [arXiv:2210.06451 \[hep-th\]](#).
- [45] G. U. Jakobsen, G. Mogull, J. Plefka, B. Sauer, and Y. Xu, *Conservative Scattering of Spinning Black Holes at Fourth Post-Minkowskian Order*, *Phys. Rev. Lett.* **131** (2023) 151401, [arXiv:2306.01714 \[hep-th\]](#).
- [46] G. U. Jakobsen, G. Mogull, J. Plefka, and B. Sauer, *Dissipative Scattering of Spinning Black Holes at Fourth Post-Minkowskian Order*, *Phys. Rev. Lett.* **131** (2023) 241402, [arXiv:2308.11514 \[hep-th\]](#).
- [47] M. Driesse, G. U. Jakobsen, G. Mogull, J. Plefka, B. Sauer, and J. Usovitsch, *Conservative Black Hole Scattering at Fifth Post-Minkowskian and First Self-Force Order*, *Phys. Rev. Lett.* **132** (2024) 241402, [arXiv:2403.07781 \[hep-th\]](#).
- [48] A. Brandhuber, G. Chen, G. Travaglini, and C. Wen, *Classical gravitational scattering from a gauge-invariant double copy*, *JHEP* **10** (2021) 118, [arXiv:2108.04216 \[hep-th\]](#).
- [49] K. Cho, K. Kim, and K. Lee, *Binary black holes and quantum off-shell recursion*, *JHEP* **05** (2024) 050, [arXiv:2311.01284 \[hep-th\]](#).
- [50] D. Akpinar, F. Febres Cordero, M. Kraus, M. S. Ruf, and M. Zeng, *Spinning Black Hole Scattering at $\mathcal{O}(G^3S^2)$: Casimir Terms, Radial Action and Hidden Symmetry*, (2024), [arXiv:2407.19005 \[hep-th\]](#).
- [51] A. Le Tiec, *The Overlap of Numerical Relativity, Perturbation Theory and Post-Newtonian Theory in the Binary Black Hole Problem*, *Int. J. Mod. Phys. D* **23** (2014) 1430022, [arXiv:1408.5505 \[gr-qc\]](#).
- [52] D. Bini, T. Damour, and A. Gericco, *Novel approach to binary dynamics: application to the fifth post-Newtonian level*, *Phys. Rev. Lett.* **123** (2019) 231104, [arXiv:1909.02375 \[gr-qc\]](#).
- [53] A. Le Tiec, L. Blanchet, and B. F. Whiting, *The First Law of Binary Black Hole Mechanics in General Relativity and Post-Newtonian Theory*, *Phys. Rev. D* **85** (2012) 064039, [arXiv:1111.5378 \[gr-qc\]](#).
- [54] S. E. Gralla and A. Le Tiec, *Thermodynamics of a Black Hole with Moon*, *Phys. Rev. D* **88** (2013) 044021, [arXiv:1210.8444 \[gr-qc\]](#).
- [55] L. Blanchet, A. Buonanno, and A. Le Tiec, *First law of mechanics for black hole binaries with spins*, *Phys. Rev. D* **87** (2013) 024030, [arXiv:1211.1060 \[gr-qc\]](#).
- [56] A. Le Tiec, *A Note on Celestial Mechanics in Kerr Spacetime*, *Class. Quant. Grav.* **31** (2014) 097001, [arXiv:1311.3836 \[gr-qc\]](#).
- [57] A. Le Tiec, *First Law of Mechanics for Compact Binaries on Eccentric Orbits*, *Phys. Rev. D* **92** (2015) 084021, [arXiv:1506.05648 \[gr-qc\]](#).
- [58] R. Fujita, S. Isoyama, A. Le Tiec, H. Nakano, N. Sago, and T. Tanaka, *Hamiltonian Formulation of the Conservative Self-Force Dynamics in the Kerr Geometry*, *Class. Quant. Grav.* **34** (2017) 134001, [arXiv:1612.02504 \[gr-qc\]](#).

- [59] L. Blanchet and A. Le Tiec, *First Law of Compact Binary Mechanics with Gravitational-Wave Tails*, *Class. Quant. Grav.* **34** (2017) 164001, [arXiv:1702.06839 \[gr-qc\]](#).
- [60] P. Ramond and A. Le Tiec, *First law of mechanics for spinning compact binaries: Dipolar order*, *Phys. Rev. D* **106** (2022) 044057, [arXiv:2202.09345 \[gr-qc\]](#).
- [61] J. L. Friedman, K. Uryu, and M. Shibata, *Thermodynamics of binary black holes and neutron stars*, *Phys. Rev. D* **65** (2002) 064035, [Erratum: *Phys. Rev. D* 70, 129904 (2004)], [arXiv:gr-qc/0108070](#).
- [62] B. Wardell, A. Pound, N. Warburton, J. Miller, L. Durkan, and A. Le Tiec, *Gravitational Waveforms for Compact Binaries from Second-Order Self-Force Theory*, *Phys. Rev. Lett.* **130** (2023) 241402, [arXiv:2112.12265 \[gr-qc\]](#).
- [63] A. Pound, B. Wardell, N. Warburton, and J. Miller, *Second-Order Self-Force Calculation of Gravitational Binding Energy in Compact Binaries*, *Phys. Rev. Lett.* **124** (2020) 021101, [arXiv:1908.07419 \[gr-qc\]](#).
- [64] T. Damour, A. Nagar, and S. Bernuzzi, *Improved effective-one-body description of coalescing nonspinning black-hole binaries and its numerical-relativity completion*, *Phys. Rev. D* **87** (2013) 084035, [arXiv:1212.4357 \[gr-qc\]](#).
- [65] D. Bini and T. Damour, *Analytical determination of the two-body gravitational interaction potential at the fourth post-Newtonian approximation*, *Phys. Rev. D* **87** (2013) 121501, [arXiv:1305.4884 \[gr-qc\]](#).
- [66] N. Afshordi *et al.* (LISA Consortium Waveform Working Group), *Waveform Modelling for the Laser Interferometer Space Antenna*, (2023), [arXiv:2311.01300 \[gr-qc\]](#).
- [67] S. Isoyama, R. Fujita, H. Nakano, N. Sago, and T. Tanaka, *“Flux-balance formulae” for extreme mass-ratio inspirals*, *PTEP* **2019** (2019) 013E01, [arXiv:1809.11118 \[gr-qc\]](#).
- [68] F. M. Blanco and E. E. Flanagan, *Particle Motion under the Conservative Piece of the Self-Force is Hamiltonian*, *Phys. Rev. Lett.* **130** (2023) 051201, [arXiv:2205.01667 \[gr-qc\]](#).
- [69] R. Gonzo, J. Lewis, and A. Pound, *to appear*.
- [70] L. Barack and O. Long, *Self-force correction to the deflection angle in black-hole scattering: A scalar charge toy model*, *Phys. Rev. D* **106** (2022) 104031, [arXiv:2209.03740 \[gr-qc\]](#).
- [71] B. Carter, *Global structure of the Kerr family of gravitational fields*, *Phys. Rev.* **174** (1968) 1559.
- [72] W. Schmidt, *Celestial mechanics in Kerr spacetime*, *Class. Quant. Grav.* **19** (2002) 2743, [arXiv:gr-qc/0202090](#).
- [73] We could equivalently define the radial action as the Hadamard finite part of Eq. (6), which is the coefficient of ϵ^0 in the integral’s Laurent series around the $\epsilon = 0$ [119]. The scattering angle and elapsed times then inherit this Hadamard regularization.
- [74] See Supplemental Material at [URL will be inserted by publisher] for a full derivation of the conservative first law for scattering orbits from the S-matrix formalism, the closed-form and PM-expanded formulas of the regularized scattering observables (scattering angle, time delay and elapsed proper time) at 0SF order and their analytic continuation to the corresponding bound observables.
- [75] E. Poisson, A. Pound, and I. Vega, *The Motion of point particles in curved spacetime*, *Living Rev. Rel.* **14** (2011) 7, [arXiv:1102.0529 \[gr-qc\]](#).
- [76] S. L. Detweiler and B. F. Whiting, *Selfforce via a Green’s function decomposition*, *Phys. Rev. D* **67** (2003) 024025, [arXiv:gr-qc/0202086](#).
- [77] A. Pound, *Motion of small objects in curved spacetimes: An introduction to gravitational self-force*, *Fund. Theor. Phys.* **179** (2015) 399, [arXiv:1506.06245 \[gr-qc\]](#).
- [78] O. Long, C. Whittall, and L. Barack, *Black hole scattering near the transition to plunge: Self-force and resummation of post-Minkowskian theory*, (2024), [arXiv:2406.08363 \[gr-qc\]](#).
- [79] A. Pound, *Self-consistent gravitational self-force*, *Phys. Rev. D* **81** (2010) 024023, [arXiv:0907.5197 \[gr-qc\]](#).
- [80] J. Lewis, T. Kakehi, A. Pound, and T. Tanaka, *Post-adiabatic dynamics and waveform generation in self-force theory: an invariant pseudo-Hamiltonian framework*, (2025), [arXiv:2507.08081 \[gr-qc\]](#).
- [81] D. A. Kosower, B. Maybee, and D. O’Connell, *Amplitudes, Observables, and Classical Scattering*, *JHEP* **02** (2019) 137, [arXiv:1811.10950 \[hep-th\]](#).
- [82] S. Weinberg, *Infrared photons and gravitons*, *Phys. Rev.* **140** (1965) B516.
- [83] In the GSF context, we would expect the coefficient of such a variation to be related to the surface gravity of the primary BH [53, 54], while in the PM limit it should asymptote to the elapsed proper time of the BH [108, 120].
- [84] S. E. Gralla and K. Lobo, *Self-force effects in post-Minkowskian scattering*, *Class. Quant. Grav.* **39** (2022) 095001, [Erratum: *Class. Quant. Grav.* 41, 179501 (2024)], [arXiv:2110.08681 \[gr-qc\]](#).
- [85] C. Cheung, J. Parra-Martinez, I. Z. Rothstein, N. Shah, and J. Wilson-Gerow, *Effective Field Theory for Extreme Mass Ratio Binaries*, *Phys. Rev. Lett.* **132** (2024) 091402, [arXiv:2308.14832 \[hep-th\]](#).
- [86] C. Cheung, J. Parra-Martinez, I. Z. Rothstein, N. Shah, and J. Wilson-Gerow, *Gravitational Scattering and Beyond from Extreme Mass Ratio Effective Field Theory*, (2024), [arXiv:2406.14770 \[hep-th\]](#).
- [87] S. L. Detweiler and E. Poisson, *Low multipole contributions to the gravitational selfforce*, *Phys. Rev. D* **69** (2004) 084019, [arXiv:gr-qc/0312010](#).
- [88] R. Bonetto, A. Pound, and Z. Sam, *Deformed Schwarzschild horizons in second-order perturbation theory: Mass, geometry, and teleology*, *Phys. Rev. D* **105** (2022) 024048, [arXiv:2109.09514 \[gr-qc\]](#).
- [89] Notice that in the limit $\hbar \rightarrow 0$, the commutators $(-i/\hbar)[\cdot, \cdot]$ are equivalent to suitable Dirac brackets $\{\cdot, \cdot\}$ which generate the time evolution in the classical phase space [118, 121].
- [90] P. Jaranowski and G. Schafer, *Towards the 4th post-Newtonian Hamiltonian for two-point-mass systems*, *Phys. Rev. D* **86** (2012) 061503, [arXiv:1207.5448 \[gr-qc\]](#).
- [91] P. Jaranowski and G. Schäfer, *Dimensional regularization of local singularities in the 4th post-Newtonian two-point-mass Hamiltonian*, *Phys. Rev. D* **87** (2013) 081503, [arXiv:1303.3225 \[gr-qc\]](#).
- [92] P. Jaranowski and G. Schäfer, *Derivation of local-in-time fourth post-Newtonian ADM Hamiltonian for spinless compact binaries*, *Phys. Rev. D* **92** (2015) 124043, [arXiv:1508.01016 \[gr-qc\]](#).
- [93] T. Damour, P. Jaranowski, and G. Schäfer, *Nonlocal-in-time action for the fourth post-Newtonian con-*

- servative dynamics of two-body systems*, *Phys. Rev. D* **89** (2014) 064058, [arXiv:1401.4548 \[gr-qc\]](#).
- [94] T. Damour, P. Jaranowski, and G. Schäfer, *Conservative dynamics of two-body systems at the fourth post-Newtonian approximation of general relativity*, *Phys. Rev. D* **93** (2016) 084014, [arXiv:1601.01283 \[gr-qc\]](#).
- [95] L. Bernard, L. Blanchet, A. Bohé, G. Faye, and S. Marsat, *Fokker action of nonspinning compact binaries at the fourth post-Newtonian approximation*, *Phys. Rev. D* **93** (2016) 084037, [arXiv:1512.02876 \[gr-qc\]](#).
- [96] L. Bernard, L. Blanchet, A. Bohé, G. Faye, and S. Marsat, *Energy and periastron advance of compact binaries on circular orbits at the fourth post-Newtonian order*, *Phys. Rev. D* **95** (2017) 044026, [arXiv:1610.07934 \[gr-qc\]](#).
- [97] L. Bernard, L. Blanchet, A. Bohé, G. Faye, and S. Marsat, *Dimensional regularization of the IR divergences in the Fokker action of point-particle binaries at the fourth post-Newtonian order*, *Phys. Rev. D* **96** (2017) 104043, [arXiv:1706.08480 \[gr-qc\]](#).
- [98] A. Le Tiec, E. Barausse, and A. Buonanno, *Gravitational Self-Force Correction to the Binding Energy of Compact Binary Systems*, *Phys. Rev. Lett.* **108** (2012) 131103, [arXiv:1111.5609 \[gr-qc\]](#).
- [99] S. Akcay, L. Barack, T. Damour, and N. Sago, *Gravitational self-force and the effective-one-body formalism between the innermost stable circular orbit and the light ring*, *Phys. Rev. D* **86** (2012) 104041, [arXiv:1209.0964 \[gr-qc\]](#).
- [100] S. Isoyama, L. Barack, S. R. Dolan, A. Le Tiec, H. Nakano, A. G. Shah, T. Tanaka, and N. Warburton, *Gravitational Self-Force Correction to the Innermost Stable Circular Equatorial Orbit of a Kerr Black Hole*, *Phys. Rev. Lett.* **113** (2014) 161101, [arXiv:1404.6133 \[gr-qc\]](#).
- [101] M. van de Meent, *Self-force corrections to the periastron advance around a spinning black hole*, *Phys. Rev. Lett.* **118** (2017) 011101, [arXiv:1610.03497 \[gr-qc\]](#).
- [102] E. Barausse, A. Buonanno, and A. Le Tiec, *The complete non-spinning effective-one-body metric at linear order in the mass ratio*, *Phys. Rev. D* **85** (2012) 064010, [arXiv:1111.5610 \[gr-qc\]](#).
- [103] S. Akcay and M. van de Meent, *Numerical computation of the effective-one-body potential q using self-force results*, *Phys. Rev. D* **93** (2016) 064063, [arXiv:1512.03392 \[gr-qc\]](#).
- [104] D. Bini, T. Damour, and A. Gericco, *Confirming and improving post-Newtonian and effective-one-body results from self-force computations along eccentric orbits around a Schwarzschild black hole*, *Phys. Rev. D* **93** (2016) 064023, [arXiv:1511.04533 \[gr-qc\]](#).
- [105] D. Bini, T. Damour, and A. Gericco, *New gravitational self-force analytical results for eccentric orbits around a Schwarzschild black hole*, *Phys. Rev. D* **93** (2016) 104017, [arXiv:1601.02988 \[gr-qc\]](#).
- [106] D. Bini, T. Damour, and A. Gericco, *Spin-dependent two-body interactions from gravitational self-force computations*, *Phys. Rev. D* **92** (2015) 124058, [Erratum: *Phys. Rev. D* **93**, 109902 (2016)], [arXiv:1510.06230 \[gr-qc\]](#).
- [107] M. Colleoni, L. Barack, A. G. Shah, and M. van de Meent, *Self-force as a cosmic censor in the Kerr overspinning problem*, *Phys. Rev. D* **92** (2015) 084044, [arXiv:1508.04031 \[gr-qc\]](#).
- [108] A. Zimmerman, A. G. M. Lewis, and H. P. Pfeiffer, *Redshift factor and the first law of binary black hole mechanics in numerical simulations*, *Phys. Rev. Lett.* **117** (2016) 191101, [arXiv:1606.08056 \[gr-qc\]](#).
- [109] A. Le Tiec and P. Grandclément, *Horizon Surface Gravity in Corotating Black Hole Binaries*, *Class. Quant. Grav.* **35** (2018) 144002, [arXiv:1710.03673 \[gr-qc\]](#).
- [110] L. Barack, M. Colleoni, T. Damour, S. Isoyama, and N. Sago, *Self-force effects on the marginally bound zoom-whirl orbit in Schwarzschild spacetime*, *Phys. Rev. D* **100** (2019) 124015, [arXiv:1909.06103 \[gr-qc\]](#).
- [111] A. Albertini, A. Nagar, A. Pound, N. Warburton, B. Wardell, L. Durkan, and J. Miller, *Comparing second-order gravitational self-force, numerical relativity, and effective one body waveforms from inspiralling, quasicircular, and nonspinning black hole binaries*, *Phys. Rev. D* **106** (2022) 084061, [arXiv:2208.01049 \[gr-qc\]](#).
- [112] L. Barack *et al.*, *Comparison of post-Minkowskian and self-force expansions: Scattering in a scalar charge toy model*, *Phys. Rev. D* **108** (2023) 024025, [arXiv:2304.09200 \[hep-th\]](#).
- [113] C. Whittall and L. Barack, *Frequency-domain approach to self-force in hyperbolic scattering*, *Phys. Rev. D* **108** (2023) 064017, [arXiv:2305.09724 \[gr-qc\]](#).
- [114] D. Bini, A. Gericco, C. Kavanagh, A. Pound, and D. Usseglio, *Post-Minkowskian self-force in the low-velocity limit: scalar field scattering*, (2024), [arXiv:2406.15878 \[gr-qc\]](#).
- [115] A. Pound and B. Wardell, *Black hole perturbation theory and gravitational self-force*, (2021), [10.1007/978-981-15-4702-7_38 - 1](#), [arXiv:2101.04592 \[gr-qc\]](#).
- [116] J. Mathews and A. Pound, *Post-adiabatic waveform-generation framework for asymmetric precessing binaries*, (2025), [arXiv:2501.01413 \[gr-qc\]](#).
- [117] Z. Nasipak, *Metric reconstruction and the Hamiltonian for eccentric, precessing binaries in the small-mass-ratio limit*, (2025), [arXiv:2507.07746 \[gr-qc\]](#).
- [118] R. Gonzo and C. Shi, *Scattering and bound observables for spinning particles in Kerr spacetime with generic spin orientations*, (2024), [arXiv:2405.09687 \[hep-th\]](#).
- [119] L. Blanchet and G. Faye, *Hadamard regularization*, *J. Math. Phys.* **41** (2000) 7675, [arXiv:gr-qc/0004008](#).
- [120] S. N. Albalat, A. Zimmerman, M. Giesler, and M. A. Scheel, *Redshift factor and the small mass-ratio limit in binary black hole simulations*, *Phys. Rev. D* **106** (2022) 044006, [arXiv:2203.04893 \[gr-qc\]](#).
- [121] J.-H. Kim, J.-W. Kim, and S. Lee, *Massive twistor worldline in electromagnetic fields*, *JHEP* **08** (2024) 080, [arXiv:2405.17056 \[hep-th\]](#).