

GATOS – XI. Excess dust heating in the narrow-line regions of nearby AGN revealed with *JWST*/MIRI

Houda Haidar^{1b,1*}, David J. Rosario,¹ Ismael García-Bernete,² Almudena Alonso-Herrero,² Anelise Audibert,^{3,4} Steph Campbell^{1b}, Chris M. Harrison^{1b}, Tiago Costa,¹ Laura Hermosa Muñoz^{1b},² Françoise Combes,⁵ Dimitra Rigopoulou^{1b},⁶ Claudio Ricci,^{7,8} Cristina Ramos Almeida^{1b},^{3,4} Enrica Bellocchi^{1b},^{9,10} Peter Boorman,^{11,12} Andrew Bunker,⁶ Richard Davies^{1b}, Daniel Delaney,^{13,14} Tanio Díaz Santos,^{15,16} Federico Esposito,¹⁷ Victoria A. Fawcett^{1b},¹ Poshak Gandhi^{1b},¹⁸ Santiago García-Burillo,¹⁷ Omaira González-Martín,¹⁹ Erin K. S. Hicks,^{13,14,20} Sebastian F. Hönig,¹⁸ Alvaro Labiano,²¹ Nancy A. Levenson,²² Enrique Lopez-Rodriguez^{1b},^{23,24} Chris Packham,^{20,25} Miguel Pereira-Santaella,²⁶ Rogemar A. Riffel^{1b},^{27,28} Alberto Rodríguez Ardila^{1b},^{29,30} John Schneider,²⁰ T. Taro Shimizu^{1b}, Marko Stalevski^{1b},^{31,32} Montserrat Villar Martín^{1b},²⁷ Martin Ward,³³ Lulu Zhang^{1b},²⁰ Gillian Leeds²³ and Fergus R. Donnan^{1b,34}

¹*School of Mathematics, Statistics and Physics, Newcastle University, Newcastle upon Tyne NE1 7RU, UK*

²*Centro de Astrobiología (CAB), CSIC–INTA, Camino Bajo del Castillo s/n, E-28692 Villanueva de la Cañada, Madrid, Spain*

³*Instituto de Astrofísica de Canarias, Calle Via Láctea s/n, E-38205 La Laguna, Tenerife, Spain*

⁴*Departamento de Astrofísica, Universidad de La Laguna, E-38206 La Laguna, Tenerife, Spain*

⁵*LUX, Observatoire de Paris, Collège de France, PSL University, CNRS, Sorbonne University, F-75014, Paris, France*

⁶*Department of Physics, University of Oxford, Keble Road, Oxford OX1 3RH, UK*

⁷*Department of Astronomy, University of Geneva, Chemin d'Ecogia 16, CH-1290 Versoix, Switzerland*

⁸*Instituto de Estudios Astrofísicos, Facultad de Ingeniería y Ciencias, Universidad Diego Portales, Av. Ejército Libertador 441, Santiago, Chile*

⁹*Departamento de Física de la Tierra y Astrofísica, Facultad de CC Físicas, Universidad Complutense de Madrid, E-28040 Madrid, Spain*

¹⁰*Instituto de Física de Partículas y del Cosmos (IPARCOS), Facultad de CC Físicas, Universidad Complutense de Madrid, E-28040 Madrid, Spain*

¹¹*Cahill Center for Astrophysics, California Institute of Technology, 1216 East California Boulevard, Pasadena, CA 91125, USA*

¹²*Max-Planck-Institut für Extraterrestrische Physik, Gießenbachstraße 1, D-85748 Garching, Germany*

¹³*Department of Physics & Astronomy, University of Alaska Anchorage, Anchorage, AK 99508-4664, USA*

¹⁴*Department of Physics, University of Alaska, Fairbanks, AK 99775-5920, USA*

¹⁵*Institute of Astrophysics, Foundation for Research and Technology–Hellas (FORTH), Heraklion GR-70013, Greece*

¹⁶*School of Sciences, European University Cyprus, Diogenes Street, Engomi, 1516 Nicosia, Cyprus*

¹⁷*Observatorio Astronómico Nacional (OAN–IGN), Observatorio de Madrid, Alfonso XII 3, E-28014 Madrid, Spain*

¹⁸*Department of Physics & Astronomy, University of Southampton, Highfield, Southampton SO17 1BJ, UK*

¹⁹*Instituto de Radioastronomía y Astrofísica (IRyA), Universidad Nacional Autónoma de México, Antigua Carretera a Pátzcuaro #8701, Colonia ExHda. San José de la Huerta, C.P. 58089 Morelia, Michoacán, Mexico*

²⁰*Department of Physics and Astronomy, The University of Texas at San Antonio (UTSA), 1 UTSA Circle, San Antonio, TX 78249-0600, USA*

²¹*Telespazio UK for ESA, ESAC, Camino Bajo del Castillo s/n, E-28692 Villanueva de la Cañada, Spain*

²²*Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA*

²³*Department of Physics & Astronomy, University of South Carolina, Columbia, SC 29208, USA*

²⁴*Kavli Institute for Particle Astrophysics & Cosmology (KIPAC), Stanford University, Stanford, CA 94305, USA*

²⁵*National Astronomical Observatory of Japan, National Institutes of Natural Sciences (NINS), 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan*

²⁶*Instituto de Física Fundamental (IFF), CSIC, Calle Serrano 123, E-28006 Madrid, Spain*

²⁷*Centro de Astrobiología (CAB), CSIC–INTA, Ctra. de Ajalvir km 4, Torrejón de Ardoz, E-28850 Madrid, Spain*

²⁸*Departamento de Física, CCNE, Universidade Federal de Santa Maria, Av. Roraima 1000, 97105-900 Santa Maria, RS, Brazil*

²⁹*Laboratório Nacional de Astrofísica (LNA/MCTI), 37530-000 Itajubá, MG, Brazil*

³⁰*Observatório Nacional, Rua General José Cristino 77, 20921-400 São Cristóvão, Rio de Janeiro, RJ, Brazil*

³¹*Astronomical Observatory, Volgina 7, 11060 Belgrade, Serbia*

³²*Sterrenkundig Observatorium, Universiteit Gent, Krijgslaan 281-S9, B-9000 Gent, Belgium*

³³*Centre for Extragalactic Astronomy, Department of Physics, Durham University, South Road, Durham DH1 3LE, UK*

³⁴*Department of Astronomy & Astrophysics, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA*

ABSTRACT

We present *James Webb Space Telescope*/Mid-Infrared Instrument imaging of eight nearby active galactic nuclei (AGN) from the GATOS (Galactic Activity, Torus, and Outflow Survey) survey to investigate the physical conditions of extended dust in their narrow-line regions (NLRs). In four galaxies (ESO 428–G14, NGC 4388, NGC 3081, and NGC 5728), we detect spatially resolved dust structures extending ~ 100 – 200 pc along the NLR. In these systems, we find a strong link between the morphology of the dust, the radio ejecta, and the coronal [Si VI] emission, implying that dust carries imprints of the processes shaping the NLR. Using spatially resolved spectral energy distributions, we show that dust in the NLR has systematically steeper slopes than star-forming clumps. This dust emits at temperatures in the range 150–220 K, at a distance of ~ 150 pc from the nucleus. Using simple models, we show that, even under optimistic assumptions of grain size and AGN luminosity, the excess mid-infrared emission cannot be explained by AGN illumination alone. We interpret this excess heating as *in situ*. We show that shocks with velocities $v_{\text{shock}} \sim 200$ – 400 km s $^{-1}$ in dense gas can close this gap, and in some cases even account for the total observed emission. This, combined with multiple lines of evidence for shocks in these regions, supports a scenario in which shocks not only coexist with dust but may be playing a key role in heating it. Our findings reveal shocks may be an important and previously overlooked driver of extended dust emission in the central hundreds of parsecs in AGN.

Key words: methods: miscellaneous – methods: observational – galaxies: nuclei – galaxies: Seyfert.

1 INTRODUCTION

Interstellar dust lies at the heart of galaxy evolution, serving as a key driver of the thermodynamics and chemistry of galaxies (e.g. E. E. Salpeter 1977; D. C. B. Whittet & C. M. Leung 1993; J. C. Weingartner & B. T. Draine 2001; B. T. Draine 2003, 2011). Dust shapes the spectral energy distributions (SEDs) of galaxies, by absorbing and scattering optical and ultraviolet (UV) photons, then re-emitting this energy predominantly as infrared (IR) radiation (e.g. B. T. Draine & H. M. Lee 1984; B. T. Draine & N. Anderson 1985; D. Calzetti 2001; A. Li & B. T. Draine 2001; A. G. G. M. Tielens 2005). Additionally, grain surfaces serve as a medium where essential chemical reactions form molecules (e.g. D. Hollenbach & E. E. Salpeter 1971). At the same time, these grains also act as the primary reservoir in which refractory elements are stored, while regulating their availability in the gas phase through accretion and destruction processes, giving rise to a diverse and complex chemistry (e.g. D. C. B. Whittet & C. M. Leung 1993; J. C. Weingartner & B. T. Draine 2001).

Dust proves especially important in the context of active galactic nuclei (AGN). These luminous systems, powered by the accretion onto supermassive black holes, are responsible for regulating star formation in their host galaxies through feedback processes driven by powerful winds and jets (e.g. G. Tabor & J. Binney 1993; J. Silk & M. J. Rees 1998; A. C. Fabian 2012; A. King & K. Pounds 2015; C. M. Harrison & C. Ramos Almeida 2024). In many AGN, strong optical extinction has been attributed to the presence of an optically obscuring thick structure, dubbed as ‘torus’ (e.g. J. H. Krolik & M. C. Begelman 1988; W. Jaffe et al. 2004; M. Nenkova et al. 2008; C. Ramos Almeida & C. Ricci 2017). In the AGN unification model, this dusty torus has been held responsible for the dichotomy between type 1 and type 2 AGN (R. R. J. Antonucci & J. S. Miller 1985; C. M. Urry & P. Padovani 1995). The dusty torus has now been imaged in the submillimetre with Atacama Large Millimeter/submillimetre Array (ALMA) and in the near- and mid-IR with Very Large Telescope Interferometer with spatial resolutions spanning from a few to tens of parsecs (J. F. Gallimore

et al. 2016; S. García-Burillo et al. 2016, 2019, 2021; A. Alonso-Herrero et al. 2018, 2019, 2023; F. Combes et al. 2019; GRAVITY Collaboration 2020; V. Gámez Rosas et al. 2022).

In addition to this, substantial effort in high-resolution IR astronomy over the past decades has revealed compelling evidence of a significant amount of dust located beyond the torus, and extending from a few parsecs to tens of parsecs along the narrow-line region (NLR) in a number of nearby AGN (e.g. S. F. Hönig et al. 2012, 2013; K. R. W. Tristram et al. 2014; J. H. Leftley et al. 2018). In some cases, this has been interpreted as a dusty wind originating from the inner edges of the torus (see S. F. Hönig 2019, for a review). For dust to be launched from the torus itself, radiation pressure must exceed gravitational forces, as theoretically demonstrated by A. C. Fabian, A. Celotti & M. C. Erlund (2006) and also M. Venzani, S. Hönig & D. Williamson (2020).

The presence of dust along outflowing regions in Seyferts has long been recognized (e.g. J. J. Bock et al. 2000; J. T. Radomski et al. 2003), with early studies already highlighting its role as a key component of the NLR (e.g. G. M. MacAlpine 1985; D. E. Osterbrock 1989). The latter is subject to outflows, driven by winds or jets, which can alter the distribution of the embedded dusty gas. Dust can carry imprints of such processes, making it a direct tracer of how AGN impact their surroundings. While AGN photoionization is considered the primary radiation source powering emission in the NLR, several works have shown that it alone may not be enough, and that contributions from shocks cannot be neglected (e.g. S. M. Viegas-Aldrovandi & M. Contini 1989; J. A. Morse, J. C. Raymond & A. S. Wilson 1996; J. W. Ferguson, K. T. Korista & G. J. Ferland 1997). In many Seyferts, shocks have been shown to be a viable mechanism for shaping the spatial and velocity distribution of gas in the NLR, particularly in regions where the jet or wind interacts with the interstellar medium (ISM; e.g. H. Falcke, A. S. Wilson & C. Simpson 1998; I. Evans et al. 1999; M. Whittle & A. S. Wilson 2004; R. Morganti, C. N. Tadhunter & T. A. Oosterloo 2005; J. F. Gallimore et al. 2006; D. J. Rosario et al. 2010a, b; B. MijWSTngo et al. 2012; D. R. A. Williams et al. 2017).

Recent *James Webb Space Telescope* (*JWST*)/Mid-Infrared Instrument (MIRI) imaging of the nearby Seyfert ESO 428–G14 has successfully resolved parsec-scale dust along the NLR (H. Haidar

* E-mail: h.haidar2@newcastle.ac.uk

et al. 2024, fig. 1 therein), extending up to 100 pc from the nucleus. This dust bears a striking morphological resemblance to the radio and [Si VI] emission, coinciding with regions where shocks are prevalent (H. Falcke et al. 1996; R. A. Riffel et al. 2006; D. May et al. 2018). By comparing AGN illumination, dusty winds, and shocks as potential sources of dust heating, we argued in H. Haidar et al. (2024) that the radiative output from fast shocks can reproduce the observed dust temperatures better than heating from AGN radiation fields alone. This puts forward a scenario in which shocks play a central role in driving the IR emission within the central few hundred parsecs of AGN.

Finding dust to be cospatial with shocked regions raises a critical question about its survival. It is well established that processes such as sputtering and shattering in shocks are the key drivers of dust destruction in the ISM (e.g. A. G. G. M. Tielens et al. 1994; A. P. Jones et al. 1994; A. P. Jones, A. G. G. M. Tielens & D. J. Hollenbach 1996). Sputtering, in particular, releases refractory elements, such as iron (Fe), that were previously locked within dust grains, back into the gas phase. Since [Fe II] is also an efficient coolant in shocked gas (e.g. P. Hartigan, J. Raymond & R. Pierson 2004), enhanced [Fe II] emission is commonly used as a signature of shocks (B. T. Draine & E. E. Salpeter 1979). Several hypotheses and models have been put forward to explain how dust endures the passage of a shock: shielding by cool clouds, longer destruction time-scale, resistance through coagulation, or consistent replenishment processes at play (M. Villar-Martin et al. 2001; F. D. Priestley et al. 2022; J. Chastenot et al. 2024; F. Kirchschrager et al. 2024; H. M. Richie et al. 2024; S. Y. Dedikov & E. O. Vasiliev 2025). Additionally, the detection of molecular outflows suggests that dust may survive for similar reasons, such as shielding by dense clumps and rapid reformation mechanisms (D. S. N. Rupke, K. Gültekin & S. Veilleux 2017; A. J. Richings & C.-A. Faucher-Giguère 2018a, b; R. J. Farber & M. Gronke 2022; M. Otsuki & H. Hirashita 2024).

Building on our pilot study of ESO 428–G14, we expand this analysis to the full sample from the *JWST* Cycle 1 programme ‘*Dust in the Wind*’ (ID: 2064, PI: D. Rosario) within the ambit of the Galactic Activity, Torus, and Outflow Survey (GATOS) Collaboration (see also A. Alonso-Herrero et al. 2021; S. García-Burillo et al. 2021; I. García-Bernete et al. 2024a, b; L. Zhang et al. 2024; L. Fuller et al. 2025), which consists of eight nearby AGN (see Table 1) that show some evidence of extended dust from ground-based observations (D. Asmus, S. F. Hönig & P. Gandhi 2016). Our goal is to explore (1) the extent and morphology of resolved dust in the NLR, (2) whether this dust coexists with shocks, and (3) whether shock heating can contribute to the observed dust emission. This paper is organized as follows. The observations and data reduction are presented in Section 2. The results are described in Section 3 and discussed in Section 4. A summary of our results is given in Section 5.

2 OBSERVATIONS AND METHODS

2.1 *JWST*/MIRI imaging

All targets (see Table 1) were observed with the *JWST*/MIRI in five broad-band imaging filters: *F560W*, *F1000W*, *F1500W*, *F1800W*, and *F2100W* (corresponding to central wavelengths $\lambda_{\text{central}} = 5.6, 10, 15, 18,$ and $21 \mu\text{m}$, respectively). The angular resolution of the MIRI imaging is set by the point spread function (PSF), with full widths at half-maximum (FWHM) of 0.2, 0.32,

0.48, 0.59, and 0.67 arcsec for the filters listed above.¹ For key morphological comparisons discussed later in Section 3, we focus on the *F1000W* band, which corresponds to a physical PSF size of $\sim 23\text{--}90$ pc, depending on the target (see Table 1). The SUB256 subarray was used (28.2 arcsec on a side), along with a fast read-out mode and the shortest allowed number of groups ($N_{\text{groups}} = 5$) to minimize the saturation of the bright nuclear emission expected in the targets. A relatively large number of integrations ($N_{\text{ints}} = 50$) and four dithered exposures yielded a final exposure time of 358 s in each filter. The raw data were downloaded from the *Mikulski Archive for Space Telescopes (MAST)*,² and processed using the *JWST* pipeline PYTHON package version 10.2 with CRDS reference files `jwst_1097.pmap`.

Data reduction and subtraction of the PSF were performed as in H. Haidar et al. (2024). Given the bright nature of these galaxies in the MIR, most images had a few partially saturated pixels at the nucleus. These specific pixels were restored from up-the-ramp readouts by turning off the first-frame flagging step during Stage 1 of the *JWST* pipeline (see H. Haidar et al. 2024, for more details). This enables the flux from the first group to be included, yielding two unsaturated groups from which a reliable ramp slope can be estimated.³

In the specific case of NGC 3227, only a single good group was available. To address this, we turned off the `suppress_one_group` parameter in the ramp-fitting step of Stage 1, thus permitting ramp estimates on a single group. Further details on target selection, data processing, and the removal of the bright central point-source emission will be provided in Rosario et al. (in preparation).

2.2 VLT/SINFONI spectroscopy

K-band (1.9–2.5 μm) integral-field spectroscopic data are available for all our targets, obtained using the Spectrograph for Integral Field Observations in the Near Infrared (SINFONI) on the European Southern Observatory (ESO) Very Large Telescope (VLT; F. Eisenhauer et al. 2003; H. Bonnet et al. 2004). Data reduction was carried out with the SINFONI data reduction software.

For most targets, the observations were obtained with the 100 mas plate scale, except in the case of NGC 3227 and NGC 5135, which were observed with the 250 mas plate scale. After processing, the effective spatial sampling of the data cubes is half the plate scale, corresponding to $0.05 \text{ arcsec pixel}^{-1}$ for the *K100* data [field of view (FoV) $\sim 4 \text{ arcsec} \times 4 \text{ arcsec}$] and $0.125 \text{ arcsec pixel}^{-1}$ for the *K250* data (FoV $\sim 8 \text{ arcsec} \times 8 \text{ arcsec}$). All targets are obtained with adaptive optics (AO) assistance except in the case of NGC 3227 and NGC 5135. The AO-assisted targets achieve an average angular resolution of PSF FWHM ~ 0.17 arcsec, while the non-AO observations have a typical PSF of 0.41 arcsec. As we discuss later in Section 3, for the key targets exhibiting dusty NLRs (see Figs 2 and 5), these SINFONI observations provide resolutions that are comparable to, if not superior to, *JWST*’s *F1000W* filter (see Section 2.1).

The emission line flux maps for [Si VI] $\lambda 1.96 \mu\text{m}$ were generated using the custom IDL code LINEFIT (R. Davies et al. 2011), which fits the emission line of interest with an unresolved line profile (a

¹See also <https://jwst-docs.stsci.edu/mid-infrared-instrument/miri-observing-modes/miri-imaging#gsc.tab=0>.

²<https://mast.stsci.edu/portal/Mashup/Clients/Mast/Portal.html>

³See also *JWST* Technical Report JWST-STScI-008866, SM-12.

Table 1. Sample properties.

AGN	Type	Distance (Mpc)	Scale (pc arcmin ⁻¹)	log ₁₀ (L _{2-10 keV}) (erg s ⁻¹)	log ₁₀ (L _{bol}) (erg s ⁻¹)	PA _{NLR} (°)	PA _{radio} (°)	PA _{disc} (°)
NGC 4388	Sy 2	19.2	93	42.5	43.7	30	30	90
ESO 428–G14	Sy 2	23.2	112	41.55	42.74	130	127	62
NGC 3081	Sy 2	37	180	42.78	43.98	164	164	60
NGC 5728	Sy 2	39	190	42.83	44	120	127	53
NGC 3227	Sy 1.5	15	73	42.0	43.19	195	170	68
NGC 2992	Sy 1.5–2	30.9	150	41.94	43.12	125	154	90
NGC 7172	Sy 2	33.9	180	42.67	43.86	60	90	53
NGC 5135	Sy 2	64.8	281	43.25	44.45	30	–	25

Note. (1) Target name. (2) Type based on optical AGN classification from D. Asmus et al. (2014). (3) Redshift-independent distances are taken from NED. (4) The pixel scale is derived from the distance. (5) Hard X-ray luminosities (2–10 keV) are computed from the integrated *Swift*/BAT fluxes, unless stated otherwise (see Section 2.4). (6) Bolometric luminosities are obtained by applying a bolometric correction to the X-ray luminosities (see equation 1). (7 and 8) The NLR and radio position angles (PA) are taken from D. Asmus et al. (2016). (9) Disc inclinations are taken from HyperLeda.

sky line) convolved with a Gaussian, as well as a linear function to the line-free continuum. This single-component Gaussian fitting procedure was performed for each spaxel of the data cube. More details on the SINFONI reduction and analysis can be found in D. Delaney et al. (2025).

2.3 VLA radio continuum imaging

All radio maps for this study (except NGC 4388, for which the radio map was kindly provided by A. J. Sargent et al. 2024) were extracted from the National Radio Astronomy Observatory (NRAO) Very Large Array (VLA) [Archive Survey \(NVAS\) Images portal](#), which provides pipeline-processed radio continuum images for a major subset of the VLA archive up to 2005. We rely on the AIPS-processed pipeline outputs from NVAS, with no additional processing or cleaning. We select images of all targets with angular resolutions <1 arcsec and sufficient sensitivity to detect extended radio emission. For each galaxy, we adopt the radio band showing the most prominent extended structure along the NLR. These are: 4–8.5 GHz for ESO 428–G14, NGC 4388, NGC 5135, NGC 3227, and NGC 2992; and 1.5 GHz for NGC 3081, NGC 5728, and NGC 7172. The 4–8.5 GHz observations provide higher angular resolution, while the 1.5 GHz data have lower resolutions but are more sensitive to low surface brightness emission. This leads to the different spatial scales probed across the sample, as seen in Figs 5 and C1. The angular resolution of the radio images, as measured from the beams, is 0.44 arcsec × 0.20 arcsec for ESO 428–G14, 0.38 arcsec × 0.27 arcsec for NGC 4388, 2.25 arcsec × 1.14 arcsec for NGC 3081, and 2.02 arcsec × 1.16 arcsec for NGC 5728 (see middle panel in Fig. 5, bottom left for the beams). For ESO 428–G14 and NGC 4388, these radio maps provide sufficient resolution to compare with *JWST*'s *F1000W* filter (PSF FWHM ∼ 0.32 arcsec). For NGC 5728, while we cannot compare dust-radio morphologies, one could still constrain the orientation of the jet, as also presented in earlier studies (see e.g. T. T. Shimizu et al. 2019). In the case of NGC 3081, the radio map is unresolved at the available resolution, and no morphological conclusions are drawn in this case; the map is presented for completeness only. We present more details on this later in Section 3.4.

For the additional galaxies shown in Appendix C (Fig. C1), the corresponding beam sizes are 1.3 arcsec × 0.8 arcsec (NGC 5135), 3.5 arcsec × 1.1 arcsec (NGC 7172), 0.28 arcsec × 0.24 arcsec (NGC 3227), and 0.52 arcsec × 0.39 arcsec (NGC 2992).

2.4 Bolometric luminosities

To compute AGN bolometric luminosities ($L_{\text{bol,AGN}}$), we use estimates of the absorption-corrected 2–10 keV X-ray luminosity from C. Ricci et al. (2017), in which X-ray spectral fits have been performed for all AGN in the 70-month *Swift*/BAT all-sky survey. ESO 428–G14 and NGC 5135 are not covered by the survey, in which case, the X-ray luminosity is extracted from N. A. Levenson et al. (2006) and V. Singh et al. (2012), respectively. The X-ray luminosities are converted to $L_{\text{bol,AGN}}$ using the bolometric correction K_X from F. Duras et al. (2020):

$$K_X(L_{2-10\text{keV}}) = 15.33 \left[1 + \left(\frac{\log(L_{2-10\text{keV}}/L_\odot)}{11.8} \right)^{16.2} \right]. \quad (1)$$

2.5 Emission line contamination

In H. Haidar et al. (2024), we discussed in detail how *JWST*/MIRI broad-band imaging can be subject to contamination from strong emission lines, and as a result, these images may not exclusively trace dust emission. For example, the *F1000W* filter is mainly affected by contributions from the [S IV]λ10.51 μm and H₂ 0–0 S(2) [9.65 μm] lines. This can make it challenging to determine the morphology and extent of the dust, potentially attributing features to dust that may, in fact, be closely linked to ionized or molecular gas instead.

We developed a method to correct for contamination, using ESO 428–G14 as a case study (see H. Haidar et al. 2024), successfully producing a line-free image in the *F1000W* filter. The decontamination method was further extended by S. Campbell et al. (2025) to include the full Cycle 1 *JWST*/GATOS sample and all MIRI filter bands in the ‘Dust in the Wind’ programme. S. Campbell et al. (2025) leverage *JWST*/Medium-Resolution Spectrometer (MRS) spectroscopy available for three of the galaxies (NGC 5728, NGC 3081, and NGC 7172) to reproduce synthetic broad-band images and cross-match them with the MIRI imaging. By using this technique, they are able to accurately separate the dust continuum from the emission lines and correct the real images accordingly. In the absence of *JWST*/MRS, S. Campbell et al. (2025) find that the imaging-based only method presented in H. Haidar et al. (2024) systematically overestimates the emission line contribution by 5–10 per cent, making it a conservative but reliable measure for dust emission. The images presented in this paper are corrected using this improved method. We note that, while MRS spectroscopy provides an excellent tool to constrain

the contribution of emission lines to the MIRI filters, the spatial resolution is typically ~ 25 per cent worse than the imager, making the latter more suitable for resolving fine dust structures (S. Campbell et al. 2025). For a full treatment of the contamination correction method and its effects on our images, see S. Campbell et al. (2025).

For morphological studies, we exclusively use images in the *F1000W* filter, as this band offers the best compromise between spatial resolution with *JWST* and the prominence of the nuclear extended MIR emission. Longer wavelength images have progressively worse resolution from 15–21 μm , yielding lower detail in the extended emission. The *F560W* images offer the highest resolution, but our previous studies show that dust around the AGN does not emit strongly in this band (H. Haidar et al. 2024; S. Campbell et al. 2025).

3 RESULTS

3.1 Dust in the narrow-line region

We present the full *JWST*/MIRI imaging sample in Figs 1 and 2. The images correspond to the *F1000W* continuum (dust only) emission, following PSF subtraction and correction for emission line contamination (H. Haidar et al. 2024; S. Campbell et al. 2025). In each galaxy, the mid-infrared (MIR) emission reveals a range of distinct features, including star-forming discs, either seen face-on (e.g. NGC 5135, NGC 5728) or edge-on (e.g. NGC 7172, NGC 2992), as well as nuclear rings (e.g. NGC 3081, NGC 3227). Each galaxy displays a distinct morphology in the MIR with a wealth of unique features. At the same time, they are all characterized by the presence of a bright nuclear component that, in some cases, outshines the entire galaxy disc. This nuclear component may include emission from the unresolved torus region and will be investigated in detail in follow-up studies focused on individual galaxies. In what follows, we focus on the central 4 arcsec \times 4 arcsec region (~ 300 –800 pc, depending on the galaxy), which covers the circumnuclear region and matches SINFONI’s FoV that maps the ionized [Si VI] emission.

Below, we describe each galaxy in turn, highlighting whether the dust morphology aligns with the NLR axis as traced by ionized and molecular gas. Here, we define *coupling* as the spatial and morphological correspondence between dust and ionized and/or molecular outflows, implying that the dust may be entrained or otherwise influenced by the NLR.

3.1.1 Strong dust–NLR coupling

Starting with ESO 428–G14 (Fig. 1, top-left panel), the dust emission reveals an asymmetric morphology, extending along a position angle (PA, measured north to east) of $\sim 131^\circ$ from the north-west (NW) to the south-east (SE), with a total extent of ~ 250 pc. This emission follows closely the direction of the radio jet (H. Falcke et al. 1996, 1998) and known molecular and ionized outflows in this AGN (e.g. D. May et al. 2018; C. Feruglio et al. 2020). This indicates that dust is spatially coupled with the ionized NLR gas and the radio jet, as also demonstrated in H. Haidar et al. (2024).

NGC 4388 (Fig. 1, top-right panel) is another complex system, featuring dust structures that extend from north-east (NE) to south-west (SW) along a PA of $\sim 30^\circ$. The bulk of the extended dust emission is concentrated SW of the nucleus, reaching distances up to ~ 150 pc. Towards the NE, the dust structure splits

into two thin and faint strands, with the longer one extending up to ~ 150 pc and the shorter one to ~ 90 pc. Recent mid- to far-IR imaging with *SOFIA* (*Stratospheric Observatory for Infrared Astronomy*) further supports this picture, revealing extended 30–40 μm dust emission coincident with the NLR at a similar PA (L. Fuller et al. 2025). This orientation is also consistent with that of the NLR and the radio jet (e.g. A. Rodríguez-Ardila et al. 2017), indicating that the dust is well coupled with the NLR.

In NGC 3081 (Fig. 1, bottom-left panel), the morphology of the dust traces an S-like shape extending north to south, and along a PA similar to that of the NLR (PA $\sim 165^\circ$). To the NE of the AGN, the dust emission peaks prominently at ~ 100 pc, coinciding with the region where P. Ferruit, A. S. Wilson & J. Mulchaey (2000) previously reported an enhancement in the [O III]/H α ratio with values as bright as the nucleus itself. The dust overlaps with the ionized bipolar outflow detected by A. Schnorr-Müller et al. (2016) within the inner ~ 200 pc region, further supporting that the dust is coupled with the NLR.

Similarly, NGC 5728 (Fig. 1, bottom-right panel) also exhibits a symmetric biconal dust structure. The latter extends up to ~ 150 pc from the nucleus on both sides of the cone, oriented at a PA of $\sim 120^\circ$. The dust emission enhances the edges of the cones, taking on a shape that traces the molecular and ionized outflows propagating through the disc (T. T. Shimizu et al. 2019; R. Davies et al. 2024; I. García-Bernete et al. 2024b). To the NW, a prominent slab-like feature at ~ 330 pc appears facing the northern cone, also detected in the neon lines with *JWST*/MRS (R. Davies et al. 2024, their fig. 3). This further supports the case in which the dust is physically coupled with the ionized NLR in this galaxy.

3.1.2 Weak or unclear dust–NLR coupling

In contrast, the remaining galaxies show bright MIR nuclei and/or rings, but with no clear evidence that dust is linked to the NLR and/or outflows. In the case of NGC 7172 (Fig. 2, top-left panel), the MIR emission is dominated by the prominent edge-on dust lane. Within the inner 1 arcsec \times 1 arcsec, a faint hint of extended dust can be seen SW to the nucleus, but it is embedded in the PSF and detectable only after subtraction. While previous studies have established the presence of ionized cones and outflows along the NLR in this galaxy (A. D. Thomas et al. 2017; A. Alonso Herrero et al. 2023; L. Hermosa Muñoz et al. 2024), the nearly face-on orientation of the ionized cone, combined with the edge-on disc, makes NGC 7172 a case of weak coupling (L. Hermosa Muñoz et al. 2024). This further supports that the dust within the inner few hundred parsecs in this galaxy may not be well coupled with the NLR.

NGC 2992 (Fig. 2, top-right panel) shows a similar behaviour to NGC 7172, where the MIR emission is dominated by the disc (I. García-Bernete et al. 2015). While the inner 1 arcsec \times 1 arcsec shows a bright nuclear source, there is no evidence of extended dust along the NLR (PA $\sim 125^\circ$). Interestingly, as shown in Fig. A1, we detect very faint dust clumps along this PA at kpc scale distances from the nucleus, visible only after strong contrast stretching, which coincide with the outflowing CO clumps reported (M. V. Zanchettin et al. 2023, their fig. 6). This is in a region known to have powerful [O III] outflows (M. Guolo-Pereira et al. 2021; M. V. Zanchettin et al. 2023), providing tentative evidence that some dust may be entrained on kpc scales. In the central ~ 200 pc, the dust emission comes mainly from the disc, and the




Figure 1. *JWST/MIRI F1000W* images of the central regions of four galaxies (ESO 428–G14, NGC 4388, NGC 3081, and NGC 5728). North is up, and east is to the left. The images have been PSF-subtracted and decontaminated from emission lines. Each main panel shows the pure dust continuum emission in logarithmic scale, highlighting extended structures. The zoom-ins display the flux maps of [Si VI] emission observed with SINFONI, matched to a ~ 4 arcsec \times 4 arcsec FoV. Overlaid contours show the extended dust emission in the main panels. Where relevant, arrows mark regions where shocks are prevalent, based on [Fe II] enhancement.

[Si VI] emission appears compact, with no clear evidence that dust and the NLR are connected at these scales.

Another special case is NGC 3227 (Fig. 2, bottom-left panel), which is nearly face-on and hosts a circumnuclear stellar ring (F. K. B. Barbosa et al. 2006). The ring dominates the MIR emission with PA of $\sim -30^\circ$, as also detected in the near-IR (R. I. Davies et al. 2006). The NLR is defined by an ionization cone oriented along a PA of $\sim 30^\circ$, with evidence of ionized and molecular outflows (T. C. Fischer et al. 2013; A. Alonso-Herrero et al. 2019; G. E. Polack et al. 2024). The cone is visible only on the NE side in previous studies (G. E. Polack et al. 2024), but it is not traced in our [Si VI] map (Fig. 2), where no distinct ionization structure is apparent. The cone is almost perpendicular to the nuclear ring, indicating that the dust detected in *JWST* is not participating in the outflow.

Finally, NGC 5135 (Fig. 2, bottom-right panel) is a strongly barred spiral galaxy, with a central morphology resembling the nuclear ring in *Hubble Space Telescope* imaging (G. E. Polack et al. 2024). Although the galaxy hosts an extended ionized NLR and outflows (A. G. Bedregal et al. 2009), the MIR emission arises primarily from the ring with no spatial evidence of it being associated or coupled with the NLR. This is in agreement with G. Sabatini et al. (2018), where they report that the molecular gas

kinematics are dominated by a barred, star-forming (SF) structure.

This diversity in the sample illustrates that the NLR is a multiphase medium. All eight galaxies in this programme have been selected based on prior evidence of extended MIR emission on scales of tens to hundreds of parsecs from ground-based observations (D. Asmus et al. 2016). However, such extended emission is not clearly detected in all cases. In ESO 428–G14, NGC 4388, NGC 3081, and NGC 5728, the morphology of the dust appears to be closely coupled with the ionized NLR, possibly being entrained in the outflow. On the other hand, NGC 7172, NGC 2992, NGC 3227, and NGC 5153 show weak or inconclusive coupling between the dust and the ionized and/or molecular phases. It is possible that any dust associated with the NLR or outflow in these galaxies lies on smaller physical scales (≤ 50 pc) and therefore remains unresolved by *JWST/MIRI*. In general, these galaxies also do not show PAs that favour strong interactions between the NLR and the disc (see Table 1), which would otherwise allow the outflow to easily carry material off the disc.

For the remainder of the paper, we focus on the first four galaxies (ESO 428–G14, NGC 4388, NGC 3081, and NGC 5728) because they provide the clearest morphological evidence for




Figure 2. Similar to Fig. 1, but for NGC 7172, NGC 2992, NGC 3227, and NGC 5135.

dust–NLR coupling out to scales of a few hundred parsecs from the nucleus. This spatial correlation between dust, ionized, and molecular gas suggests that the dust may be dynamically linked to the NLR, allowing us to investigate its physical properties, survival, and role in AGN feedback.

3.2 MIR regions of interest

To assess whether the dust in the NLR is characterized by properties different from other MIR-emitting regions, we define several regions of interest (ROIs) across the full sample. From these ROIs, we compute photometric SEDs for each galaxy, as shown in Figs B1–B8. All apertures are chosen with diameters larger than the PSF FWHM of the longest wavelength filter ($F2100W$, FWHM = 0.67 arcsec) to minimize the need for wavelength-dependent aperture corrections (see also H. Haidar et al. 2024). For all ROIs, we adopt an aperture diameter of 0.7 arcsec, corresponding to one FWHM of the lowest resolution data set (i.e. $F2100W$). We note that going through each ROI and SED galaxy-by-galaxy is beyond the scope of this paper, and we defer this analysis to future work.

Fig. 3 (left) shows the median SEDs across all ROIs and galaxies, separating dusty NLRs from SF clumps. As shown in Section 3.1, only four galaxies have resolved dust along the NLR (ESO 428–G14, NGC 4388, NGC 5728, and NGC 3081). As a result, we restrict the NLR SEDs to these targets. Across all galaxies, we find a systematic difference between SEDs of dust

in the NLR and those of SF clumps. On average, SF clumps are characterized by flatter and hotter SED distributions, while dusty NLRs produce SEDs that are steeper by a factor of 2 (e.g. $F_{18\mu\text{m}, \text{NLR}}/F_{10\mu\text{m}, \text{NLR}} = 4$). While each ROI in each galaxy shows its own unique distribution, some common trends can still be identified across the sample. For example, SF clumps consistently produce higher $5.6\mu\text{m}$ contributions (e.g. NGC 7172, NGC 3227, and NGC 2992). This rise towards the shortest wavelengths may correspond to a low level of stellar light in the MIR. For dusty NLRs, a slight flux enhancement at $5.6\mu\text{m}$ is likely due to this stellar component (≤ 10 per cent; Rosario et al., in preparation). In general, both NLR dust and SF clumps produce SEDs that peak at around at $18\text{--}21\mu\text{m}$. In SF clumps, a modest excess in the $F1800W$ band over an otherwise flat SED shape could instead arise from polycyclic aromatic hydrocarbon emission (I. García-Bernete et al. 2022).

Overall, we find the dust in the NLR region emits strongly in the MIR and is physically distinct from SF clumps. This suggests that these two regions may be subject to different radiation fields and/or may host distinct dust grain populations.

3.3 Temperature of dusty NLRs

We have shown that the NLRs of ESO 428–G14, NGC 4388, NGC 3081, and NGC 5728 are characterized by dusty, distinct morphologies that emit strongly in the MIR. In this section,




Figure 3. Left: SEDs taken over five filters (*F560W*, *F1000W*, *F1500W*, *F1800W*, and *F2100W*) normalized by the *F1500W* filter. Different colours represent different galaxies. Each curve shows the median SED of all ROIs for that galaxy. For each galaxy, the dashed lines represent the median SED of SF regions. Where relevant, we plot the median for regions covering dust in the NLR, represented as solid lines. The latter have been shifted up in the Y-axis for clarity. The shaded area represents the 16th–84th percentile. Right: Galaxy median SEDs for dust in the NLR (solid) and for star-forming clumps (dashed).




Figure 4. Monte Carlo blackbody fits for the dusty NLRs of ESO 428–G14, NGC 4388 (top row), NGC 3081, and NGC 5728 (bottom row). Each panel shows an SED extracted with an aperture diameter of 0.7 arcsec at ~ 150 pc from the nucleus, where shocks dominate (see Fig. 1). Apertures are marked as orange circles in the upper left of each panel. Red stars are the photometric points. The solid lines represent the median of the fits, and the shaded bands show the 16th–84th percentiles. For each galaxy, we show the median temperature T_{med} , and the 16th–84th values (T_{low} , T_{high}).

we quantify this emission by constraining the dust temperature along their NLRs. To do this, we extract SEDs from ROIs located within the NLRs at a common distance from the nucleus. A fixed distance allows us to quantify whether AGN properties have an impact on the observed dust temperature. We adopt a distance $d_{\text{AGN}} = 150$ pc from the nucleus. This distance is large enough such that, in the more distant galaxies, it exceeds the $F2100W$ PSF FWHM (0.65 arcsec), yet not so large that, in the nearer systems, we do not risk missing regions where NLR dust is still detected. All apertures are taken with a diameter $d = 0.7$ arcsec, as also done in Section 3.2. We note that these regions also overlap with known shock spots from enhanced [Fe II] emission (see later in Section 3.4.1).

Given that we only have five photometric points, it is challenging to constrain the dust temperature from a single blackbody (BB) fit. As such, we perform a Monte Carlo (MC) fitting approach, where, in each of the 1000 draws, we perturb the fluxes within their uncertainties, randomly dropping one of the photometric points and fit a single-temperature BB to the remaining four. The fits are performed on absolute fluxes, treating the $F560W$ band as an upper limit to account for any potential stellar contribution to the dust emission (Rosario et al., in preparation). Additionally, we adopt the absolute $F560W$ flux uncertainty as a uniform conservative error across all bands, preventing unrealistically tight temperature constraints given the very small per band errors.

The solid lines in Fig. 4 show the medians of the MC BB fits, with the shaded areas representing the 16th–84th percentile distribution. On average, all dusty NLRs produce temperatures within the range of ~ 150 – 200 K, consistent with warm dust that peaks in the MIR ($\lambda \sim 15$ – 19 μm). The grain population that this represents would depend on the strength of the radiation field heating the dust (e.g. B. S. Hensley & B. T. Draine 2023).

Recently, E. Lopez-Rodriguez et al. (2025) conducted a *JWST*/MRS continuum study on six nearby galaxies, including NGC 3081, NGC 5728, and NGC 7172. They report a relatively constant dust temperature of ~ 130 K (peaking at 22 μm) within the inner 500 pc, which is lower than the values we derive, albeit measured over a different spatial region. Nevertheless, their study also finds that dust temperatures are broadly consistent across galaxies and show no evident correlation with AGN properties, in agreement with our results.

3.4 A dusty–radio–coronal link?

We compare the morphologies of dust, radio, and coronal emission to investigate whether dust near AGN is spatially associated with other AGN-driven components. In Fig. 5, the images for each galaxy are astrometrically aligned and reprojected onto the largest pixel scale for a common pixel grid. We show the images in their native angular resolutions, rather than convolving to a common PSF, in order to preserve the finer details that each data set presents. The different angular resolutions probed by *JWST*, SINFONI, and the VLA are discussed in detail in Section 2. In most cases, the angular resolution achieved with SINFONI (~ 0.17 arcsec) is finer than that of *JWST*'s $F1000W$ filter (~ 0.32 arcsec). For the radio maps where a morphological comparison is relevant (ESO 428–G14: 0.44 arcsec \times 0.20 arcsec and NGC 4388: 0.38 arcsec \times 0.27 arcsec), the angular resolution is comparable to that of *JWST* (see Section 2.3). For the purposes of our qualitative morphological inspection, these differences in resolution are not expected to strongly affect our conclusions on

the relative extents, nor the orientations, of the emission along the NLR.

We find that AGN with dusty NLRs (i.e. ESO 428–G14, NGC 4388, NGC 3081, and NGC 5728) show a strong spatial correlation with the radio emission, as shown in Fig. 5. Specifically, the radio maps for ESO 428–G14 (H. Falcke et al. 1996, 1998) and NGC 4388 (A. J. Sargent et al. 2024) exhibit a striking, spatially resolved correspondence with the dust. In both cases, the dust and radio structures extend along the NLR, and their morphologies closely trace one another, as shown in the first two panels of Fig. 5.

For NGC 5728, the 1.5 GHz radio emission is prominent along a PA consistent with the NLR and the dust emission, but extends primarily towards the NW (A. S. Wilson et al. 1993), reaching several hundred parsecs beyond the observed dusty bicone. A fainter contribution from the disc is also present, though less prominent than the jet/NLR emission. NGC 3081, on the other hand, shows only a tentative hint of the 1.5 GHz radio emission along the NLR, but the radio map is not well resolved and is aligned with the direction of the beam (2.25 arcsec \times 1.14 arcsec) as also demonstrated in N. M. Nagar et al. (1999) and C. G. Mundell et al. (2009). As such, we do not draw any conclusions from the map presented in Fig. 5. Very recently, A. J. Sargent et al. (2025) present new VLA 6 GHz imaging of NGC 3081, revealing spatially resolved radio emission that extends well beyond the beam and up to ~ 170 pc from the nucleus on both sides (see their fig. 1). The emission reported in A. J. Sargent et al. (2025) is extended along a similar PA to the dust probed in our study, further supporting a dust–radio link in NGC 3081.

In Fig. C1, we show that several galaxies with weak dust–NLR coupling nevertheless present interesting large-scale radio features. In particular, NGC 2992 exhibits a striking ‘figure-eight’ in the 4.8 GHz radio map. This has been interpreted in the literature as radio bubbles, likely linked to stellar activity (S. C. Chapman et al. 2000). Overall, we find no spatial correspondence between the radio bubbles and the dust emission. NGC 5135 is another unique case, hosting a radio hotspot at ~ 1 kpc from the nucleus, associated with a supernova (SN) shock (A. G. Bedregal et al. 2009; L. Colina et al. 2012), which is evident in the 4–8 GHz map as a large radio hotspot. While the radio emission traces the dust morphology nicely at larger scales, within the central 4 arcsec \times 4 arcsec the galaxy shows no clear link between the dust and the radio along the NLR. Overall, this work motivates future homogenous radio observations for these key targets.

Another compelling connection is that between the dust and the coronal line emission. In Fig. 5, we systematically find that the morphology of the dust closely mirrors that of the [Si VI] emission, regardless of whether there is a tight dust–radio correlation or not. This is also true for galaxies with weak dust–NLR coupling, as shown in Fig. C1, where both the dust and [Si VI] emissions appear compact within the inner 4 arcsec \times 4 arcsec region.

Previous studies have shown that AGN photoionization is generally responsible for the observed coronal line emission in Seyferts. These are highly ionized species, with high ionization potentials (IPs > 100 eV), which makes them insensitive to star formation and can therefore act as direct tracers of AGN activity (e.g. M. A. Prieto, O. Marco & J. Gallimore 2005; A. Rodríguez-Ardila et al. 2006). However, when these coronal lines are observed out to several hundreds of parsecs from the nucleus, as in ESO 428–G14, NGC 4388, NGC 3081, and NGC 5728, AGN photoionization alone is unlikely to account for this. This is because




Figure 5. *JWST*/MIRI ‘Dust in the Wind’ sample. Left: Extended dust emission revealed by the *JWST*/MIRI *F1000W* image, shown after PSF subtraction and decontamination. Middle: Radio continuum emission from VLA observations. Right: Comparison between the dust morphology (image) and the radio emission (shown as white contours). The zoomed-in 4 arcsec \times 4 arcsec region displays the [Si VI] coronal line emission from SINFONI.

the high-energy photon flux becomes too low at large distances. Therefore, the detection of extended coronal lines can be used as a tracer for an additional *in situ* excitation process, such as shocks (e.g. J. W. Ferguson et al. 1997).

More recently, A. Rodríguez-Ardila et al. (2025) investigated the extent of the coronal emission in ESO 428–G14, NGC 5728, and NGC 3081, among other Seyferts, and found a strong positive trend between the size of the coronal line emission and the power of the radio jet. Their work suggests that radio jets propagating through the ambient gas drive shocks that produce coronal lines extending from several hundred parsecs to kiloparsec scales.

While it remains debated whether the radio emission in Seyferts traces a jet or a wind (e.g. H. Falcke et al. 1996; G. V. Bicknell et al. 1998; T. C. Fischer et al. 2023), this distinction is not critical here. Regardless of the nature of the radio emission, AGN hosting outflows are capable of driving shocks (e.g. M. Pereira-

Santaella et al. 2022; V. A. Fawcett et al. 2023, 2025; R. A. Riffel et al. 2025). In addition to the excitation of the coronal line, an important shock tracer is an enhancement in [Fe II]/Pa β and [Fe II]/[P II] ratios, which has already been reported in the literature within the central few hundred parsecs of these galaxies. We highlight in Fig. 1 the regions where these shocks are likely to be important. In what follows, we summarize some key observational signatures of shocks within the central few hundred parsecs of our main AGN: ESO 428–G14, NGC 4388, NGC 3081, and NGC 5728.

3.4.1 Evidence from [Fe II] emission

The enhancement in [Fe II] 1.644 μ m emission relative to a non-refractory species, such as [P II], and evaluated by means of flux ratios [Fe II]/[P II] or [Fe II]/Pa β , offer a key diagnostic to mea-

sure the relative contribution of photoionization and shocks (e.g. E. Oliva et al. 2001; C. Ramos Almeida et al. 2006; C. Ramos Almeida, A. M. Pérez García & J. A. Acosta-Pulido 2009; T. Storchi-Bergmann et al. 2009; R. A. Riffel et al. 2014). Maps using these line ratios have also been found to correlate well with radio emission, further supporting that shock excitation may be the key ingredient in the production of [Fe II] (e.g. D. A. Forbes & M. J. Ward 1993; M. Blietz et al. 1994).

(i) In ESO 428–G14, R. A. Riffel et al. (2006) report on the detection of $1.27 \leq [\text{Fe II}]/\text{Pa } \beta \leq 2.73$ ratios that are cospatial with the radio emission. The strongest [Fe II]/Pa β values coincide with the radio hotspots towards the NW, approximately ~ 120 pc away from the nucleus.

(ii) In NGC 4388, [Fe II]/Pa β peaks SW to the nucleus at ~ 150 pc (R. A. Knop et al. 2001; A. Rodríguez-Ardila et al. 2017). In A. Rodríguez-Ardila et al. (2017), they also report on the detection of double peak [Fe II] lines in the same region, further supporting an additional excitation source other than photoionization to be at play.

(iii) In NGC 5728, M. Durré & J. Mould (2018) use [Fe II]/[P II] to constrain the ionization source in the NLR. Ratios with values higher than 2 would indicate shock contribution. They compute [Fe II]/[P II] along the cone and find ratios of ~ 5.8 SW to the AGN and ~ 3.9 in the NW.

(iv) In NGC 3081, J. Reunanen, J. K. Kotilainen & M. A. Prieto (2003) leverage the [Fe II]/Br γ ratio, which can also be sensitive to the excitation mechanism, and report a weak value of 0.6. However, the coronal line emission in NGC 3081 appears more spatially complex. M. A. Prieto et al. (2005) resolved the [Si VII] 2.48 μm emission into a compact source associated with the AGN, and a fainter blob located ~ 120 pc NW to the AGN. This same blob coincides with the second brightest region in H α and [O III], previously reported by P. Ferruit et al. (2000) (see their fig. 11). M. A. Prieto et al. (2005) propose that the concentric shell-like structures seen in H α and [O III] could be indicative of propagating shock fronts. We mark this region as a tentative shock site in Fig. 1.

Taken together, the close correspondence between dust, radio, and coronal line structures, along with evidence of strong enhancement in [Fe II] emission, supports the scenario in which shocks play a key role in shaping the NLRs of these AGN. However, the extent to which shocks contribute to the IR emission, as opposed to direct AGN illumination, remains unclear. Below, we investigate the origin of dust heating in the NLR, now that the broader picture of its morphology and excitation has been established.

4 DISCUSSION

4.1 What heats the dust in the MIR?

Given the direct influence of the central source on the NLR, the simplest and most logical interpretation would be that dust in the NLR is heated and illuminated by AGN radiation fields, as previously reported in the literature for a number of Seyferts. The challenge here, however, is not whether photons are energetic enough, but whether the AGN radiation fields can sustain dust temperatures in the range of 150–200 K at distances of several hundred parsecs from the nucleus. To determine this, we constrain the expected AGN radiation field for these galaxies and

Table 2. Expected AGN radiation field U_{AGN} and dust temperatures (assuming classical ISM grain sizes) derived at a distance $d_{\text{AGN}} = 150$ pc from the nucleus.

AGN	$\log_{10} U_{\text{AGN}}$	$T_{\text{dust}, a=0.005 \mu\text{m}}$ (K)	$T_{\text{dust}, a=0.25 \mu\text{m}}$ (K)
ESO 428–G14	3.12	64	29
NGC 4388	4.08	100	46
NGC 5728	4.38	115	53
NGC 3081	4.36	114	52
NGC 7172	4.24	108	49
NGC 2992	3.50	77	35
NGC 3227	3.57	80	36
NGC 5135	4.84	142	65

assess whether it alone is able to reproduce the dust temperatures presented in Section 3.3.

Following A. G. G. M. Tielens (2010) (chapter 5, equation 5.43), and as also done in H. Haidar et al. (2024), we calculate the AGN radiation field at a distance d_{AGN} from the nucleus in terms of the expected anisotropic radiation field expressed in units of the Habing field:

$$U_{\text{AGN}} = 2.1 \times 10^4 \left(\frac{L_{\text{bol,AGN}}}{10^4 L_{\odot}} \right) \left(\frac{0.1 \text{ pc}}{d_{\text{AGN}}} \right)^2, \quad (2)$$

where $L_{\text{bol,AGN}}$ is the bolometric AGN luminosity (see Section 2.4). Subsequently, for a given grain size a , the dust temperature can be expressed as

$$T_{\text{d}} \approx 33.5 \left(\frac{1 \mu\text{m}}{a} \right)^{0.2} \left(\frac{U_{\text{AGN}}}{10^4} \right)^{0.2} \text{ K}, \quad (3)$$

following A. G. G. M. Tielens (2010) (chapter 5, equation 5.44). Note that this assumes that the intrinsic AGN SED that heats the dust peaks strongly in the UV, roughly like the local Habing field, which corresponds to $1.6 \times 10^{-3} \text{ erg cm}^{-2} \text{ s}^{-1}$ (H. J. Habing 1968). We note that AGN can also show an enhanced soft X-ray component; however, this would be inefficient at heating dust, as grain absorption efficiencies drop steeply at X-ray energies (e.g. A. Laor & B. T. Draine 1993).

We compute the AGN radiation field at a fixed distance of $d_{\text{AGN}} = 150$ pc, the same as that adopted in Section 3.3. This allows us to constrain whether the temperatures expected from AGN heating are able to reproduce those derived from the observations. We present the results for the full sample in Table 2. On average, we find that the AGN radiation field across the sample is $\log_{10} U_{\text{AGN}} \sim 4$. This corresponds to dust temperatures of $T_{\text{dust}} \sim 100$ K for the smallest grains (i.e. $a = 0.005 \mu\text{m}$) and $T_{\text{dust}} \sim 50$ K for larger grains (i.e. $a = 0.25 \mu\text{m}$), consistent with the canonical grain size distribution proposed by J. S. Mathis, W. Ruml & K. H. Nordsieck (1977).

We summarize our results in Fig. 6. The orange stars represent the dust temperatures computed from the SEDs of the shocked regions indicated in Fig. 4, and the shaded area covers the lowest and highest possible temperatures across all four galaxies $T_{\text{dust}} \sim [150, 220]$ K. The blue column represents the possible dust temperatures over a range of grain sizes $a = [0.005, 0.25] \mu\text{m}$ expected from AGN heating.

In general, for a fixed distance from the nucleus, a higher bolometric luminosity leads to higher dust temperatures. However, even when accounting for the smallest dust grain sizes, which tend to heat more efficiently, the temperature expected from AGN heating still does not reach those probed with *JWST*.




Figure 6. Comparison between the median dust temperatures from MC BB fits (orange stars) and predicted temperatures from AGN illumination (blue bars). The shaded orange area represents the minimum and maximum possible dust temperatures across the four galaxies. The colour gradient within each blue bar represents the range of predicted temperatures resulting from variations in the assumed grain size $a = [0.005, 0.25] \mu\text{m}$. For all four galaxies, additional heating mechanisms must be present, as AGN illumination alone cannot account for the observed dust temperatures.

To achieve a minimum temperature of $T \sim 150 \text{ K}$, one would require AGN luminosities in the range of $L_{\text{AGN,bol}} = [3.8 \times 10^{44}, 2 \times 10^{46}] \text{ erg s}^{-1}$ for grains $a = [0.005, 0.25] \mu\text{m}$, respectively. Values $\geq 10^{45} \text{ erg s}^{-1}$ enter the quasar regime and are atypical of local Seyferts. We thus interpret this as excess heating that cannot be accounted for by AGN illumination alone, indicating that additional *in situ* heating may be at play. Below, we discuss possible processes in more detail.

4.2 *In situ* dust heating

Several processes can, in principle, contribute to heating the dust in the NLR. Such processes include: *in situ* heating from ongoing star formation, radiation-pressure-driven polar wind, radiative shocks from the post-shock region, or simply, collisional heating from hot plasma.

We have shown in Fig. 3 that dusty NLRs produce SEDs that are systematically steeper than those of SF clumps. As such, we can rule out star formation as a dominant source of radiation in the NLR. Polar dusty winds, while effective on parsec to tens of parsec scales (S. F. Hönl 2019), cannot propagate to the hundreds of parsecs scale probed here, consistent with interferometric results and radiative transfer models (e.g. S. F. Hönl 2019; A. Alonso-Herrero et al. 2021).

Given the strong spatial correspondence between dust in the NLR and the radio emission (Fig. 5), jet and/or wind–ISM interactions, such as shocks, could contribute to the observed MIR emission. In fact, B. T. Draine (1981) reports that for fast shocks with velocities ($v_{\text{shock}} \geq 200 \text{ km s}^{-1}$), dust grains become an important coolant, radiating up to 17 per cent of the shock energy in the IR. As presented in Section 3.4, all AGN with dusty NLRs (ESO 428–G14, NGC 4388, NGC 3081, and NGC 5728) have direct evidence of shocks in the central few hundred parsecs (see

regions highlighted in Fig. 1). This further establishes shocks as a strong candidate for additional dust heating.

By adopting the M. A. Dopita & R. S. Sutherland (1995) formalism for fast radiative shocks, we demonstrated in H. Haidar et al. (2024) that the post-shock cooling region can generate radiation fields strong enough to heat the dust to MIR-emitting temperatures. As an exercise, we apply the same approach here to test which combinations of shock velocity and ISM density are needed to reproduce the minimum dust temperatures observed in our sample. The total radiative flux per unit shock surface is given by

$$F_{\text{T}} = 2.28 \times 10^{-3} \left(\frac{v_{\text{s}}}{100 \text{ km s}^{-1}} \right)^{3.0} \left(\frac{n}{\text{cm}^{-3}} \right) \text{ erg cm}^{-2} \text{ s}^{-1}, \quad (4)$$

where v_{s} is the velocity of the shock and n is the particle number density. Substituting this flux into equation (3), scaling it by the Habing field, and solving for $T_{\text{dust}} = 150 \text{ K}$ and $a = 0.005 \mu\text{m}$, implies $v_{\text{shock}} \sim 400 \text{ km s}^{-1}$ for $n \sim 10^3 \text{ cm}^{-3}$, $v_{\text{shock}} \sim 230 \text{ km s}^{-1}$ for $n \sim 5 \times 10^3 \text{ cm}^{-3}$, and $v_{\text{shock}} \sim 180 \text{ km s}^{-1}$ for $n \sim 10^4 \text{ cm}^{-3}$. These estimates show that shock velocities in the range of $v_{\text{shock}} = 200\text{--}400 \text{ km s}^{-1}$ propagating through dense enough media ($n \sim 10^{3\text{--}4} \text{ cm}^{-3}$) could already account for dust temperatures of $\sim 150 \text{ K}$.

Recent *JWST*/MRS studies compared shock models with MIR line ratio diagnostics for a number of AGN, including NGC 3081 and NGC 5728, and found significant overlap between the predictions of photoionization and shock models, making it difficult to set robust constraints on shock velocities and pre-densities (e.g. L. Zhang et al. 2025; R. A. Riffel et al. 2026). Nevertheless, such conditions are reasonable and have already been reported for the low-luminosity AGN ESO 428–G14 (R. A. Riffel et al. 2006; D. May et al. 2018). This makes shock heating not only a plausible additional source, but also a mechanism that could potentially by itself account for the observed dust temperatures. Future work will explore shock modelling tailored specifically for *JWST*/MRS observations.

Another interesting process to consider is collisional heating of dust grains by hot plasma. This is especially relevant given the strong morphological resemblance between the dust and the coronal line emission (see Fig. 1), which traces hot gas in the range $T \sim 10^{5\text{--}6} \text{ K}$ (e.g. E. Oliva et al. 1994). In such hot plasma, collisions of dust grains with electrons/ions can drive dust to higher effective temperatures (e.g. B. T. Draine & E. E. Salpeter 1979; E. Dwek 1987; M. Bocchio et al. 2013; S. A. Drozdov & Y. A. Shchekinov 2019). This process, however, strongly depends on grain size, as smaller grains undergo large temperature fluctuations but are also quickly destroyed by charging and sputtering, whereas larger ones could survive long enough to contribute to the dust emission (B. T. Draine & E. E. Salpeter 1979; E. Dwek 1987).

4.3 A surviving population of dust grains?

The presence of dust in regions dominated by shocks (Fig. 1), as indicated by strong enhancement in [Fe II] emission (Section 3.4.1), raises important questions about dust survival under such conditions. Theoretically, shocks are expected to destroy dust grains, releasing refractory elements such as [Fe II], previously locked within them, back into the gas phase (P. M. Roully & L. Spitzer 1952; L. L. Cowie 1978; A. P. Jones et al. 1994, 1996; A. G. G. M. Tielens et al. 1994; J. W. Ferguson et al. 1997). However, in a number of systems, observations show that dust is preva-

lent where shock destruction processes such as sputtering and/or shattering are expected (e.g. C. N. Tadhunter, S. M. Scarrott & C. D. Rolph 1990; D. Alloin et al. 2000; M. Villar-Martín et al. 2001; M. Shahbandeh et al. 2023; H. Haidar et al. 2024). The survival of dust in extreme conditions has attracted considerable attention in recent years, with models predicting factors such as longer destruction time-scales, grain size distribution and composition, coagulation, as well as shock velocity to play key roles in supporting dust survival (M. Bocchio, A. P. Jones & J. D. Slavin 2014; D. S. N. Rupke et al. 2017; R. J. Farber & M. Gronke 2022; Y. Dubois et al. 2024; M. Otsuki & H. Hirashita 2024; H. M. Richie et al. 2024).

The survival of dust has direct implications on the survival of H_2 molecular gas. The latter has a dissociation potential of only ~ 5 eV and would not survive direct AGN irradiation without dust shielding. This has led some models to suggest that molecules may instead form inside the outflows, with dust acting as the necessary catalyst (e.g. K. Zubovas & A. R. King 2014; A. J. Richings & C.-A. Faucher-Giguère 2018a, b). Grain growth is possible in the outflow through metal accretion and/or coagulation, each leaving a specific imprint on the extinction curve (e.g. H. Hirashita 2012; H. Hirashita & T. Nozawa 2017). On the other hand, complete dust formation (i.e. production of grain seeds) in the outflows is much more difficult as it requires gas conditions that are similar to asymptotic giant branch (AGB) stars or SN ejecta, where very dense, metal-rich, and rapidly cooling gas would enable the nucleation of stable dust precursors (e.g. M. Elvis, M. Marengo & M. Karovska 2002; A. Sarangi, E. Dwek & D. Kazanas 2019).

Dust processing also strongly depends on grain material and composition. In particular, sputtering and shattering destruction thresholds are set by material bond properties (A. G. G. M. Tielens et al. 1994). For example, in contrast to carbonaceous grains, silicate grains are thought to be difficult to reform under ISM conditions and are therefore believed to originate in AGB stars. As such, silicate grains should be preserved in the passage of shocks (A. P. Jones & J. A. Nuth 2011). In supernova (SN) shocks, M. Bocchio et al. (2014) find that carbonaceous grains are quickly destroyed, even when exposed to slow shocks $v_{\text{shock}} \sim 50 \text{ km s}^{-1}$, while silicate grains demonstrate greater resilience to shock processing. As such, a possible hypothesis would be that shocks preferentially destroy carbonaceous grains, leaving a population dominated by silicates in post-shock regions. This picture is further complicated by the fact that sputtering yields are higher for silicates than for graphite (e.g. A. G. G. M. Tielens et al. 1994), which would suggest the opposite trend. However, survival in shocks depends on more than sputtering yields alone, including grain charging, shattering, and the ability of different materials to reform in the ISM. This gap remains poorly explored and highlights the need for more realistic dust modelling in shocked environments to constrain how grains endure the passage of a shock.

5 SUMMARY AND CONCLUSION

We investigated the presence and origin of dust in the NLRs of eight nearby Seyferts from the *JWST*/MIRI imaging programme ‘*Dust in the Wind*’, part of the GATOS survey. Our key findings can be summarized as follows:

(i) Among the eight galaxies, ESO 428–G14, NGC 4388, NGC 3081, and NGC 5728 show extended dust structures within the inner $r \sim 2$ arcsec region, aligned with the NLR and exhibiting morphologies comparable to the ionized gas (see Figs 1 and 2).

(ii) For galaxies with dusty NLRs, we find a tight spatial correspondence between the morphology of the dust, the radio, and the [Si VI] emissions (see Fig. 5). Indeed, all of these galaxies host NLRs oriented at inclinations (see Table 1) that favour coupling between the extended dust, the ionized gas, and the outflows.

(iii) We extract photometric SEDs across several ROIs and find that dust in the NLR produces systematically steeper SEDs than SF clumps, with $F_{18\mu\text{m}}/F_{10\mu\text{m}}$ ratios of ~ 4 versus ~ 2 , respectively, averaged across the full sample (Fig. 3). This suggests that dust in the NLR is subject to different grain processing and/or harder radiation fields than the surrounding regions.

(iv) We compute the temperature of the dust in shock-dominated regions within the NLR, at 150 pc from the nucleus, and find consistent values within the range of 150–200 K, characteristic of warm dust peaking in the MIR (Fig. 4). Such temperatures can be easily achieved if the emission is dominated by small grains ($a = 0.005 \mu\text{m}$).

(v) Using simple models, we investigate the origin of the observed dust emission and find that, even under optimistic assumptions of grain size and AGN luminosity, direct AGN heating alone is not able to reproduce the observed dust temperature. We propose that the excess heating is instead produced *in situ*. We show that fast radiative shocks with velocities $v \sim 200\text{--}400 \text{ km s}^{-1}$ propagating through dense enough media ($n \sim 10^{3\text{--}4} \text{ cm}^{-3}$) could account for the observed temperatures.

The evolution and processing of dust in shocks depends on several parameters, such as grain size, composition, local density, and shock velocity, that are often not well constrained in observations. This highlights the need for shock and dust evolution models to evaluate how efficiently these properties contribute to the observed dust in AGN-driven outflows. Future progress will come from combining such dust models with constraints from *JWST* and ALMA, providing multiphase diagnostics needed to build a complete picture on dust survival. Already, the tentative detection of faint warm dusty clumps in NGC 2992 (see Fig. A1 and Appendix A) that can be cross-matched with outflowing CO clumps (M. V. Zanchettin et al. 2023) provides a glimpse of how the warm and cold phases interconnect in outflows, a link that would be critical to fully understand dust survival and its role in AGN feedback.

ACKNOWLEDGEMENTS

We thank the referee for a constructive report that helped improve the paper. HH thanks Martin Elvis and Brandon Hensley for valuable discussions on dust processing, Maria Vitoria Zanchettin for providing the NGC 2992 ALMA maps, and Travis Fischer and the A. J. Sargent et al. (2024) team for sharing the NGC 4388 radio maps. HH is grateful to ESA/ESAC for hosting their research visit in Madrid, with special thanks to Alvaro, Ismael, Almudena, and Laura for their support during the visit. HH acknowledges support from ESA through the Science Faculty (Funding reference ESA-SCI-E-LE-183), and funding from STFC and Newcastle University.

DJR and SC acknowledge the support of the UK STFC through grant ST/X001105/1. AA-H, LHM, and MVM acknowledge support from grant PID2021-124665NB-I00 funded by the Spanish Ministry of Science and Innovation and the State Agency of Research MCIN/AEI/10.13039/501100011033 and ERDF A way of making Europe. AA acknowledges funding from the European Union (WIDERA ExGal-Twin, GA 101158446). MP-S acknowledges support under grants RYC2021-

033094-I, CNS2023-145506, and PID2023-146667NB-I00 funded by MCIN/AEI/10.13039/501100011033 and the European Union NextGenerationEU/PRTR. IG-B was supported by the Programa Atracción de Talento Investigador ‘César Nombela’ via grant 2023-T1/TEC-29030 funded by the Community of Madrid. CR acknowledges support from SNSF Consolidator grant F01-13252, FONDECYT Regular grant 1230345, ANID BASAL project FB210003 and the China–Chile joint research fund. MS acknowledges support by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia (MSTDIRS) through contract no. 451-03-66/2024-03/200002 with the Astronomical Observatory (Belgrade). CRA and AA acknowledge support from the Agencia Estatal de Investigación of the Ministerio de Ciencia, Innovación y Universidades (MCIU/AEI) under the grant ‘Tracking active galactic nuclei feedback from parsec to kiloparsec scales’, with reference PID2022-141105NB-I00 and the European Regional Development Fund (ERDF). AB acknowledges funding from the ‘FirstGalaxies’ Advanced Grant from the European Research Council (ERC) under the European Union’s Horizon 2020 Framework Programme (grant agreement no. 789056). DR acknowledges support from STFC through grants ST/S000488/1 and ST/W000903/1. CP, JS, LZ, EKSH, acknowledge grant support from the Space Telescope Science Institute (ID: JWST-GO-02064.002). FE and SG-B acknowledge support from the Spanish grant PID2022-138560NB-I00, funded by MCIN/AEI/10.13039/501100011033/FEDER, EU. CMH acknowledges funding from a United Kingdom Research and Innovation (UKRI) grant (code: MR/V022830/1). OG-M acknowledges financial support from PAPIIT UNAM project IN109123 and ‘Ciencia de Frontera’ CONAHCyT project CF-2023-G-100. ARA acknowledges Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for partial support to this work through grant 313739/2023-4. EB acknowledges support from the Spanish grants PID2022-138621NB-I00 and PID2021-123417OB-I00, funded by MCIN/AEI/10.13039/501100011033/FEDER, EU. MW acknowledges support from a Leverhulme Emeritus Fellowship, EM-2021-064. RAR acknowledges the support from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq; Projects 303450/2022-3 and 403398/2023-1), the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES; Project 88887.894973/2023-00), and Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul (FAPERGS). SFH acknowledges support through UK Research and Innovation (UKRI) under the UK government’s Horizon Europe Funding Guarantee (EP/Z533920/1, selected in the 2023 ERC Advanced Grant round) and an STFC Small Award (ST/Y001656/1).

This work is based on observations made with the NASA/ESA/CSA *James Webb Space Telescope*. This work makes use of several PYTHON packages: NUMPY (C. R. Harris et al. 2020), MATPLOTLIB (J. D. Hunter 2007), and ASTROPY (Astropy Collaboration 2013, 2018).

DATA AVAILABILITY

All data are publicly available and can be extracted as described in Section 2. The reduced *JWST* images are available upon request; otherwise can also be extracted from *MAST* according to Section 2.

REFERENCES

Alloin D., Pantin E., Lagage P. O., Granato G. L., 2000, *A&A*, 363, 926

- Alonso-Herrero A. et al., 2018, *ApJ*, 859, 144
 Alonso-Herrero A. et al., 2019, *A&A*, 628, A65
 Alonso-Herrero A. et al., 2021, *A&A*, 652, A99
 Alonso-Herrero A. et al., 2023, *A&A*, 675, A88
 Antonucci R. R. J., Miller J. S., 1985, *ApJ*, 297, 621
 Asmus D., Hönig S. F., Gandhi P., Smette A., Duschl W. J., 2014, *MNRAS*, 439, 1648
 Asmus D., Hönig S. F., Gandhi P., 2016, *ApJ*, 822, 109
 Astropy Collaboration, 2013, *A&A*, 558, A33
 Astropy Collaboration, 2018, *AJ*, 156, 123
 Barbosa F. K. B., Storchi-Bergmann T., Cid Fernandes R., Winge C., Schmitt H., 2006, *MNRAS*, 371, 170
 Bedregal A. G., Colina L., Alonso-Herrero A., Arribas S., 2009, *ApJ*, 698, 1852
 Bicknell G. V., Dopita M. A., Tsvetanov Z. I., Sutherland R. S., 1998, *ApJ*, 495, 680
 Blietz M., Cameron M., Drapatz S., Genzel R., Krabbe A., van der Werf P., Sternberg A., Ward M., 1994, *ApJ*, 421, 92
 Bocchio M., Jones A. P., Verstraete L., Xilouris E. M., Micelotta E. R., Bianchi S., 2013, *A&A*, 556, A6
 Bocchio M., Jones A. P., Slavin J. D., 2014, *A&A*, 570, A32
 Bock J. J. et al., 2000, *AJ*, 120, 2904
 Bonnet H. et al., 2004, *The Messenger*, 117, 17
 Calzetti D., 2001, *PASP*, 113, 1449
 Campbell S. et al., 2025, *MNRAS*, 544, 648
 Chapman S. C., Morris S. L., Alonso-Herrero A., Falcke H., 2000, *MNRAS*, 314, 263
 Chastenot J. et al., 2024, *A&A*, 690, A348
 Colina L., Pereira-Santaella M., Alonso-Herrero A., Bedregal A. G., Arribas S., 2012, *ApJ*, 749, 116
 Combes F. et al., 2019, *A&A*, 623, A79
 Cowie L. L., 1978, *ApJ*, 225, 887
 Davies R. I. et al., 2006, *ApJ*, 646, 754
 Davies R. et al., 2011, *ApJ*, 741, 69
 Davies R. et al., 2024, *A&A*, 689, A263
 Dedikov S. Y., Vasiliev E. O., 2025, *New Astron.*, 114, 102293
 Delaney D., Berger C., Hicks E., Burtscher L., Rosario D., Müller-Sánchez F., Malkan M., 2025, *ApJ*, 984, 163
 Dopita M. A., Sutherland R. S., 1995, *ApJ*, 455, 468
 Draine B. T., 1981, *ApJ*, 245, 880
 Draine B. T., 2003, *ARA&A*, 41, 241
 Draine B. T., 2011, *Physics of the Interstellar and Intergalactic Medium*. Princeton Univ. Press, Princeton, NJ
 Draine B. T., Anderson N., 1985, *ApJ*, 292, 494
 Draine B. T., Lee H. M., 1984, *ApJ*, 285, 89
 Draine B. T., Salpeter E. E., 1979, *ApJ*, 231, 438
 Drozdov S. A., Shchekinov Y. A., 2019, *Astrophysics*, 62, 540
 Dubois Y. et al., 2024, *A&A*, 687, A240
 Duras F. et al., 2020, *A&A*, 636, A73
 Durré M., Mould J., 2018, *ApJ*, 867, 149
 Dwek E., 1987, *ApJ*, 322, 812
 Eisenhauer F. et al., 2003, in Iye M., Moorwood A. F. M., eds, *Proc. SPIE Conf. Vol. 4841, Instrument Design and Performance for Optical/Infrared Ground-based Telescopes*. SPIE, Bellingham, p. 1548
 Elvis M., Marengo M., Karovska M., 2002, *ApJ*, 567, L107
 Evans I., Koratkar A., Allen M., Dopita M., Tsvetanov Z., 1999, *ApJ*, 521, 531
 Fabian A. C., 2012, *ARA&A*, 50, 455
 Fabian A. C., Celotti A., Erlund M. C., 2006, *MNRAS*, 373, L16
 Falcke H., Wilson A. S., Simpson C., Bower G. A., 1996, *ApJ*, 470, L31
 Falcke H., Wilson A. S., Simpson C., 1998, *ApJ*, 502, 199
 Farber R. J., Gronke M., 2022, *MNRAS*, 510, 551
 Fawcett V. A. et al., 2023, *MNRAS*, 525, 5575
 Fawcett V. A. et al., 2025, *MNRAS*, 537, 2003
 Ferguson J. W., Korista K. T., Ferland G. J., 1997, *ApJS*, 110, 287
 Ferruit P., Wilson A. S., Mulchaey J., 2000, *ApJS*, 128, 139

- Feruglio C., Fabbiano G., Bischetti M., Elvis M., Travascio A., Fiore F., 2020, *ApJ*, 890, 29
- Fischer T. C., Crenshaw D. M., Kraemer S. B., Schmitt H. R., 2013, *ApJS*, 209, 1
- Fischer T. C., Johnson M. C., Secrest N. J., Crenshaw D. M., Kraemer S. B., 2023, *ApJ*, 953, 87
- Forbes D. A., Ward M. J., 1993, *ApJ*, 416, 150
- Fuller L. et al., 2025, *ApJS*, 276, 64
- Gallimore J. F., Axon D. J., O’Dea C. P., Baum S. A., Pedlar A., 2006, *AJ*, 132, 546
- Gallimore J. F. et al., 2016, *ApJ*, 829, L7
- Gómez Rosas V. et al., 2022, *Nature*, 602, 403
- García-Bernete I. et al., 2015, *MNRAS*, 449, 1309
- García-Bernete I. et al., 2022, *A&A*, 666, L5
- García-Bernete I. et al., 2024a, *A&A*, 681, L7
- García-Bernete I. et al., 2024b, *A&A*, 691, A162
- García-Burillo S. et al., 2016, *ApJ*, 823, L12
- García-Burillo S. et al., 2019, *A&A*, 632, A61
- García-Burillo S. et al., 2021, *A&A*, 652, A98
- GRAVITY Collaboration, 2020, *A&A*, 634, A1
- Guolo-Pereira M., Ruschel-Dutra D., Storchi-Bergmann T., Schnorr-Müller A., Cid Fernandes R., Couto G., Dametto N., Hernandez-Jimenez J. A., 2021, *MNRAS*, 502, 3618
- Habing H. J., 1968, *Bull. Astron. Inst. Netherlands*, 19, 421
- Haidar H. et al., 2024, *MNRAS*, 532, 4645
- Harris C. R. et al., 2020, *Nature*, 585, 357
- Harrison C. M., Ramos Almeida C., 2024, *Galaxies*, 12, 17
- Hartigan P., Raymond J., Pierson R., 2004, *ApJ*, 614, L69
- Hensley B. S., Draine B. T., 2023, *ApJ*, 948, 55
- Hermosa Muñoz L. et al., 2024, *A&A*, 690, A350
- Hirashita H., 2012, *MNRAS*, 422, 1263
- Hirashita H., Nozawa T., 2017, *Planet. Space Sci.*, 149, 45
- Hollenbach D., Salpeter E. E., 1971, *ApJ*, 163, 155
- Hönig S. F., 2019, *ApJ*, 884, 171
- Hönig S. F., Kishimoto M., Antonucci R., Marconi A., Prieto M. A., Tristram K., Weigelt G., 2012, *ApJ*, 755, 149
- Hönig S. F. et al., 2013, *ApJ*, 771, 87
- Hunter J. D., 2007, *Comput. Sci. Eng.*, 9, 90
- Jaffe W. et al., 2004, *Nature*, 429, 47
- Jones A. P., Nuth J. A., 2011, *A&A*, 530, A44
- Jones A. P., Tielens A. G. G. M., Hollenbach D. J., McKee C. F., 1994, *ApJ*, 433, 797
- Jones A. P., Tielens A. G. G. M., Hollenbach D. J., 1996, *ApJ*, 469, 740
- King A., Pounds K., 2015, *ARA&A*, 53, 115
- Kirschschlager F., Sartorio N. S., De Looze I., Barlow M. J., Schmidt F. D., Priestley F. D., 2024, *MNRAS*, 528, 5364
- Knop R. A., Armus L., Matthews K., Murphy T. W., Soifer B. T., 2001, *AJ*, 122, 764
- Krolik J. H., Begelman M. C., 1988, *ApJ*, 329, 702
- Laor A., Draine B. T., 1993, *ApJ*, 402, 441
- Leftley J. H., Tristram K. R. W., Hönig S. F., Kishimoto M., Asmus D., Gandhi P., 2018, *ApJ*, 862, 17
- Levenson N. A., Heckman T. M., Krolik J. H., Weaver K. A., Życki P. T., 2006, *ApJ*, 648, 111
- Li A., Draine B. T., 2001, *ApJ*, 554, 778
- Lopez-Rodriguez E. et al., 2025, *ApJ*, 994, 206
- MacAlpine G. M., 1985, in Miller J. S., ed., *Astrophysics of Active Galaxies and Quasi-Stellar Objects*. Oxford Univ. Press, Oxford, p. 259
- Mathis J. S., Rumpl W., Nordsieck K. H., 1977, *ApJ*, 217, 425
- May D., Rodríguez-Ardila A., Prieto M. A., Fernández-Ontiveros J. A., Diaz Y., Mazzalay X., 2018, *MNRAS*, 481, L105
- Mingo B., Hardcastle M. J., Croston J. H., Evans D. A., Kharb P., Kraft R. P., Lenc E., 2012, *ApJ*, 758, 95
- Morganti R., Tadhunter C. N., Oosterloo T. A., 2005, *A&A*, 444, L9
- Morse J. A., Raymond J. C., Wilson A. S., 1996, *PASP*, 108, 426
- Mundell C. G., Ferruit P., Nagar N., Wilson A. S., 2009, *ApJ*, 703, 802
- Nagar N. M., Wilson A. S., Mulchaey J. S., Gallimore J. F., 1999, *ApJS*, 120, 209
- Nenkova M., Sirocky M. M., Nikutta R., Ivezić Ž., Elitzur M., 2008, *ApJ*, 685, 160
- Oliva E., Salvati M., Moorwood A. F. M., Marconi A., 1994, *A&A*, 288, 457
- Oliva E. et al., 2001, *A&A*, 369, L5
- Osterbrock D. E., 1989, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei*. University Science Books, Mill Valley, CA
- Otsuki M., Hirashita H., 2024, *MNRAS*, 528, 5008
- Pereira-Santaella M. et al., 2022, *A&A*, 665, L11
- Polack G. E., Revalski M., Crenshaw D. M., Fischer T. C., Schmitt H. R., Kraemer S. B., Meena B., Rafelski M., 2024, *ApJ*, 975, 129
- Priestley F. D., Arias M., Barlow M. J., De Looze I., 2022, *MNRAS*, 509, 3163
- Prieto M. A., Marco O., Gallimore J., 2005, *MNRAS*, 364, L28
- Radomski J. T., Piña R. K., Packham C., Telesco C. M., De Buizer J. M., Fisher R. S., Robinson A., 2003, *ApJ*, 587, 117
- Ramos Almeida C., Ricci C., 2017, *Nat. Astron.*, 1, 679
- Ramos Almeida C., Pérez García A. M., Acosta-Pulido J. A., Rodríguez Espinosa J. M., Barrera R., Manchado A., 2006, *ApJ*, 645, 148
- Ramos Almeida C., Pérez García A. M., Acosta-Pulido J. A., 2009, *ApJ*, 694, 1379
- Reunanen J., Kotilainen J. K., Prieto M. A., 2003, *MNRAS*, 343, 192
- Ricci C. et al., 2017, *ApJS*, 233, 17
- Richie H. M., Schneider E. E., Abruzzo M. W., Torrey P., 2024, *ApJ*, 974, 81
- Richings A. J., Faucher-Giguère C.-A., 2018a, *MNRAS*, 474, 3673
- Richings A. J., Faucher-Giguère C.-A., 2018b, *MNRAS*, 478, 3100
- Riffel R. A., Storchi-Bergmann T., Winge C., Barbosa F. K. B., 2006, *MNRAS*, 373, 2
- Riffel R. A., Vale T. B., Storchi-Bergmann T., McGregor P. J., 2014, *MNRAS*, 442, 656
- Riffel R. A., Souza-Oliveira G. L., Costa-Souza J. H., Zakamska N. L., Storchi-Bergmann T., Riffel R., Bianchin M., 2025, *ApJ*, 982, 69
- Riffel R. A. et al., 2026, *A&A*, 705, A59
- Rodríguez-Ardila A., Prieto M. A., Viegas S., Gruenwald R., 2006, *ApJ*, 653, 1098
- Rodríguez-Ardila A. et al., 2017, *MNRAS*, 465, 906
- Rodríguez-Ardila A., Fonseca-Faria M. A., Dahmer-Hahn L. G., Prieto A., Riffel R., Riffel R. A., 2025, *MNRAS*, 538, 2800
- Rosario D. J., Whittle M., Nelson C. H., Wilson A. S., 2010a, *MNRAS*, 408, 565
- Rosario D. J., Whittle M., Nelson C. H., Wilson A. S., 2010b, *ApJ*, 711, L94
- Routly P. M., Spitzer L., Jr, 1952, *ApJ*, 115, 227
- Rupke D. S. N., Gültekin K., Veilleux S., 2017, *ApJ*, 850, 40
- Sabatini G., Gruppioni C., Massardi M., Giannetti A., Burkutean S., Cimatti A., Pozzi F., Talia M., 2018, *MNRAS*, 476, 5417
- Salpeter E. E., 1977, *ARA&A*, 15, 267
- Sarangi A., Dwek E., Kazanas D., 2019, *ApJ*, 885, 126
- Sargent A. J., Fischer T. C., Johnson M. C., van der Horst A. J., Secrest N. J., Shuvo O. I., Cigan P. J., Smith K. L., 2024, *ApJ*, 961, 230
- Sargent A. J., van der Horst A. J., Johnson M. C., Fischer T. C., Secrest N. J., Cigan P. J., Shuvo O. I., Smith K. L., 2025, *ApJ*, 986, 194
- Schnorr-Müller A., Storchi-Bergmann T., Robinson A., Lena D., Nagar N. M., 2016, *MNRAS*, 457, 972
- Schönell A. J., Storchi-Bergmann T., Riffel R. A., Riffel R., Bianchin M., Dahmer-Hahn L. G., Diniz M. R., Dametto N. Z., 2019, *MNRAS*, 485, 2054
- Shahbandeh M. et al., 2023, *MNRAS*, 523, 6048
- Shimizu T. T. et al., 2019, *MNRAS*, 490, 5860
- Silk J., Rees M. J., 1998, *A&A*, 331, L1
- Singh V., Risaliti G., Braito V., Shastri P., 2012, *MNRAS*, 419, 2089

- Storchi-Bergmann T., McGregor P. J., Riffel R. A., Simões Lopes R., Beck T., Dopita M., 2009, *MNRAS*, 394, 1148
- Tabor G., Binney J., 1993, *MNRAS*, 263, 323
- Tadhunter C. N., Scarrott S. M., Rolph C. D., 1990, *MNRAS*, 246, 163
- Thomas A. D. et al., 2017, *ApJS*, 232, 11
- Tielens A. G. G. M., 2005, *The Physics and Chemistry of the Interstellar Medium*. Cambridge Univ. Press, Cambridge
- Tielens A. G. G. M., 2010, *The Physics and Chemistry of the Interstellar Medium*. Cambridge Univ. Press, Cambridge
- Tielens A. G. G. M., McKee C. F., Seab C. G., Hollenbach D. J., 1994, *ApJ*, 431, 321
- Tristram K. R. W., Burtscher L., Jaffe W., Meizenheimer K., Hönig S. F., Kishimoto M., Schartmann M., Weigelt G., 2014, *A&A*, 563, A82
- Urry C. M., Padovani P., 1995, *PASP*, 107, 803
- Venanzi M., Hönig S., Williamson D., 2020, *ApJ*, 900, 174
- Viegas-Aldrovandi S. M., Contini M., 1989, *ApJ*, 339, 689
- Villar-Martín M., De Young D., Alonso-Herrero A., Allen M., Binette L., 2001, *MNRAS*, 328, 848
- Weingartner J. C., Draine B. T., 2001, *ApJ*, 548, 296
- Whittet D. C. B., Leung C. M., 1993, Status Report, 1 July 1992–30 April 1993, Rensselaer Polytechnic Institute, Troy, New York
- Whittle M., Wilson A. S., 2004, *AJ*, 127, 606
- Williams D. R. A. et al., 2017, *MNRAS*, 472, 3842
- Wilson A. S., Braatz J. A., Heckman T. M., Krolik J. H., Miley G. K., 1993, *ApJ*, 419, L61
- Zanchettin M. V. et al., 2023, *A&A*, 679, A88
- Zhang L. et al., 2024, *ApJ*, 974, 195
- Zhang L. et al., 2025, *ApJS*, 280, 65
- Zubovas K., King A. R., 2014, *MNRAS*, 439, 400

APPENDIX A: NGC 2992: WARM VERSUS COLD

M. V. Zanchettin et al. (2023) report on the detection of cold molecular clumps in NGC 2992 that are entrained with the outflow, with projected distances up to 1.7 kpc and velocities close to $\sim 200 \text{ km s}^{-1}$. In Fig. A1, we show that several of these clumps can also be detected in the MIR, albeit very faint. The *JWST* and ALMA images are astrometrically aligned, so the positions of these clumps in the *JWST* image are reliable, although they are of very low surface brightness and only visible under strong stretching of the image. We do not convolve the *JWST* image to the ALMA beam, as this would wash out the already very faint dusty clumps and prevent a visual inspection. The clumps are distributed along the NLR and also extend to kpc in scale away from the nucleus. In agreement with M. V. Zanchettin et al. (2023), we find no spatial correspondence between these clumps and the radio bubbles, which are too compact and fall below the location of these clumps. M. V. Zanchettin et al. (2023) propose that these clumps are likely linked to a previous AGN episode, which could explain their faint detection in *JWST*.

APPENDIX B: SEDS

We extract SEDs across various ROIs and for all galaxies as presented in Figs B1–B8. For ESO 428–G14, the SEDs are presented and extensively studied in H. Haidar et al. (2024) (fig. 3 therein). A case-by-case study of these SEDs is beyond the scope of this paper. Here we show them only for illustrative purposes to help interpret the medians presented in Fig. 3. Upcoming focused target studies from the GATOS programme will provide detailed case-by-case analyses of these SEDs.

APPENDIX C: DUST–RADIO LINK FOR THE REST OF THE SAMPLE

Just as done in Fig. 5, we compare the morphology of the dust with the radio and coronal line emissions for galaxies that show weak coupling between the dust and the ionized NLR (NGC 5135, NGC 7172, NGC 3227, and NGC 2992). We find that, on larger scales, the radio emission generally traces well the dust, except for NGC 2992, for which 4.8 GHz emission is dominated by the radio bubbles. Other interesting features include the SN shock in NGC 5135 (L. Colina et al. 2012, their fig. 1), which




Figure A1. Comparison between the ALMA CO(2–1) flux map (courtesy of M. Zanchettin) and the *JWST*/MIRI *F1000W* image (bottom) of NGC 2992, in logarithmic stretch. North is up, and east to the left. Cyan circles mark clumps identified in CO(2–1) by M. V. Zanchettin et al. (2023) (see their fig. 6), several of which coincide with MIR clumps detected in the *JWST* image.

appears to be cospatial with dust in the galaxy disc. L. Colina et al. (2012) estimate that the total energy produced in the SN shock is $\sim 2.7 \times 10^{42} \text{ erg s}^{-1}$. This highlights the strength of shocks in this region and their potential role in shaping the dust in the ISM. NGC 3227 shows some extended radio emission that may be

associate with the nuclear ring, which also shines bright in the MIR. A. J. Schönell et al. (2019) found that this ring is shocked with high $[\text{Fe II}]/\text{Pa } \beta$ ratios (≥ 4) throughout the ring (see their fig. 4). This dust–shock relation, for both NCG 5135 and NGC 3227, further supports that dust can survive in these regions.




Figure B1. Left: SEDs extracted from individual apertures on the PSF-subtracted and decontaminated images in NGC 5728. Red curves correspond to regions covering emission along the NLR, while blue curves correspond to SF clumps in the disc. Right: Illustrative *JWST*/MIRI three-colour image (R: *F560W*, G: *F1000W*, B: *F1500W*), produced from the original (non-PSF-subtracted) images, with ROIs overlaid to indicate the regions used for SED extraction.




Figure B2. Same as Fig. B1, but for NGC 4388.




Figure B3. Same as Fig. B1, but for NGC 3081.




Figure B4. Same as Fig. B1, but for NGC 7172.




Figure B5. Same as Fig. B1, but for NGC 3227.




Figure B6. Same as Fig. B1, but for NGC 5135.




Figure B7. Same as Fig. B1, but for NGC 2992.




Figure B8. Same as Fig. B1, but for ESO 428-G14, also presented in figs 1 and 3 in H. Haidar et al. (2024).

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.




Figure C1. Same as Fig. 5, but for the remaining four galaxies in the sample.