

# **Focusing of light by a nano-hole array**

*Nikolay Zheludev*

*Optoelectronics Research Centre, University of Southampton, SO17 1BJ, UK*

Here for the first time we demonstrate experimentally and theoretically that the Talbot effect on a quasi-periodic array of nano-holes may be used to achieve sub-wavelength field localizations and well-isolated 'hot spots' of high electromagnetic energy concentration.

It is commonly believed that far-field optical resolution is fundamentally limited by diffraction at the wavelength level because evanescent waves carrying sub-wavelength-scale information from an object dissipate in the far field. A recent remarkable theoretical discovery suggests that evanescent fields may not be needed to achieve sub-wavelength resolution: Berry and Popescu 2006 predicted that a grating structure could create sub-wavelength localizations of light that propagate further into the far field than more familiar evanescent waves. They relate this effect to the fact that band-limited functions are able to oscillate arbitrarily faster than the highest Fourier components they contain, a phenomenon called super-oscillation.

Here we report on how a quasi-periodic array of holes creates sub-wavelength localizations of light without evanescent waves and how the fields created by such diffraction fall into the class of super-oscillating fields [1]. This effect offers a new way to achieve sub-wavelength imaging in the far field.

## **References**

- [1] Focusing of light by a nano-hole array. F.Huang, Y.Chen, G.De Abajo, N.Zheludev. *APPLIED PHYSICS LETT.* 90, 091119 (2007)